



## Singular Cardinals and the pcf Theory

Thomas Jech

*The Bulletin of Symbolic Logic*, Vol. 1, No. 4. (Dec., 1995), pp. 408-424.

Stable URL:

<http://links.jstor.org/sici?sici=1079-8986%28199512%291%3A4%3C408%3ASCATPT%3E2.0.CO%3B2-5>

*The Bulletin of Symbolic Logic* is currently published by Association for Symbolic Logic.

---

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/asl.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

---

The JSTOR Archive is a trusted digital repository providing for long-term preservation and access to leading academic journals and scholarly literature from around the world. The Archive is supported by libraries, scholarly societies, publishers, and foundations. It is an initiative of JSTOR, a not-for-profit organization with a mission to help the scholarly community take advantage of advances in technology. For more information regarding JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## SINGULAR CARDINALS AND THE PCF THEORY

THOMAS JECH

**§1. Introduction.** Among the most remarkable discoveries in set theory in the last quarter century is the rich structure of the arithmetic of singular cardinals, and its deep relationship to large cardinals. The problem of finding a complete set of rules describing the behavior of the continuum function  $2^{\aleph_\alpha}$  for singular  $\aleph_\alpha$ 's, known as the *Singular Cardinals Problem*, has been attacked by many different techniques, involving forcing, large cardinals, inner models, and various combinatorial methods. The work on the singular cardinals problem has led to many often surprising results, culminating in a beautiful theory of Saharon Shelah called the pcf theory ("pcf" stands for "possible cofinalities"). The most striking result to date states that if  $2^{\aleph_n} < \aleph_\omega$  for every  $n = 0, 1, 2, \dots$ , then  $2^{\aleph_\omega} < \aleph_{\omega_4}$ .

In this paper we present a brief history of the singular cardinals problem, the present knowledge, and an introduction into Shelah's pcf theory. In Sections 2, 3 and 4 we introduce the reader to cardinal arithmetic and to the singular cardinals problems. Sections 5, 6, 7 and 8 describe the main results and methods of the last 25 years and explain the role of large cardinals in the singular cardinals problem. In Section 9 we present an outline of the pcf theory.

**§2. The arithmetic of cardinal numbers.** Cardinal numbers were introduced by Cantor in the late 19th century and problems arising from investigations of rules of arithmetic of cardinal numbers led to the birth of set theory. The operations of addition, multiplication and exponentiation of infinite cardinal numbers are a natural generalization of such operations on integers. Addition and multiplication of infinite cardinals turns out to be simple: when at least one of the numbers  $\kappa, \lambda$  is infinite then both  $\kappa + \lambda$  and  $\kappa \cdot \lambda$  are equal to  $\max\{\kappa, \lambda\}$ . In contrast with  $+$  and  $\cdot$ , exponentiation presents fundamental problems. In the simplest nontrivial case,  $2^\kappa$  represents the cardinal number of the power set  $P(\kappa)$ , the set of all subsets of  $\kappa$ . (Here we adopt the usual convention of set theory that the number  $\kappa$  is identified with a set of cardinality  $\kappa$ , namely the set of all ordinal numbers

---

Received August 16, 1995; revised September 4, 1995.

Supported in part by NSF grant DMS-9401275 and by NRC COBASE grant while visiting the Center for Theoretical Study in Czech Republic.

smaller than  $\kappa$ . In this representation, the cardinal number  $\aleph_\alpha$  is the same as the ordinal number  $\omega_\alpha$ .) By a celebrated theorem of Cantor,  $2^\kappa > \kappa$  holds for all cardinals  $\kappa$ , and therefore  $2^{\aleph_\alpha} \geq \aleph_{\alpha+1}$  for every infinite cardinal  $\aleph_\alpha$ . In [6], Cantor conjectured that  $2^{\aleph_0} = \aleph_1$ , which became known as the *Continuum Hypothesis* (and the similar conjecture  $2^{\aleph_\alpha} = \aleph_{\alpha+1}$  as the Generalized Continuum Hypothesis or GCH).

It has soon become apparent that if GCH were true, one could completely describe rules for all of cardinal arithmetic, including infinite sums and products ( $\sum_{i \in I} \kappa_i$  and  $\prod_{i \in I} \kappa_i$ ) of cardinal numbers. Despite efforts of Cantor himself and others, the question whether GCH, or even CH, is true, remained unanswered until the emergence of methods of modern logic.

For a long time, the only source of inequalities in cardinal arithmetic was König's Theorem [37]. The theorem states that if  $\{\kappa_i : i \in I\}$  and  $\{\lambda_i : i \in I\}$  are two indexed families of cardinal numbers such that  $\kappa_i < \lambda_i$  for all  $i$ , then

$$\sum_{i \in I} \kappa_i < \prod_{i \in I} \lambda_i.$$

Note that Cantor's Theorem itself is a special case of this: letting  $\kappa_i = 1$  and  $\lambda_i = 2$  for all  $i < \kappa$ , we get  $\kappa = \sum_{i < \kappa} 1 < \prod_{i < \kappa} 2 = 2^\kappa$ .

König's Theorem also provides important information about singular cardinals. Let us recall that the *cofinality* of an infinite cardinal  $\aleph_\alpha$  is the least cardinal  $\aleph_\gamma = \text{cf } \aleph_\alpha$  such that  $\aleph_\alpha$  is the supremum of an increasing  $\aleph_\gamma$ -sequence of ordinal numbers. A cardinal  $\aleph_\alpha$  is *regular* if  $\text{cf } \aleph_\alpha = \aleph_\alpha$  and *singular* if  $\text{cf } \aleph_\alpha < \aleph_\alpha$ . As  $\text{cf}(\text{cf } \aleph_\alpha) = \text{cf } \aleph_\alpha$ , the cofinality is always a regular cardinal. Each successor cardinal  $\aleph_{\alpha+1}$  is regular, as is the countable cardinal  $\aleph_0$ .

One consequence of König's Theorem is that for every  $\alpha$  and every  $\beta$ ,  $\text{cf } \aleph_\alpha^{\aleph_\beta} > \aleph_\beta$ . It follows easily that when  $\aleph_\beta \geq \text{cf } \aleph_\alpha$  then  $\aleph_\alpha^{\aleph_\beta} \neq \aleph_\alpha$ , and therefore  $\aleph_\alpha^{\aleph_\beta} > \aleph_\alpha$ . Using this as well as rather elementary rules for cardinal exponentiation due to Hausdorff and Tarski, one obtains a complete set of rules for exponentiation under the assumption of GCH:

$$\aleph_\alpha^{\aleph_\beta} = \begin{cases} \aleph_\alpha & \text{if } \aleph_\beta < \text{cf } \aleph_\alpha \\ \aleph_{\alpha+1} & \text{if } \text{cf } \aleph_\alpha \leq \aleph_\beta < \aleph_\alpha \\ \aleph_{\beta+1} & \text{if } \aleph_\alpha \leq \aleph_\beta \end{cases}$$

Note that even in the presence of GCH, singular cardinals exhibit different behavior: the second clause does not apply if  $\aleph_\alpha$  is regular.

**§3. Consistency and independence of the generalized continuum hypothesis.** The continuum problem remained open until 1939 when a significant progress came from Gödel who showed in [25] that GCH is consistent with the axioms of set theory ZFC (Zermelo-Fraenkel's axioms with the axiom of

choice). Gödel produced the model  $L$  of *constructible sets* and proved that GCH holds in the model  $L$ . In addition to the consistency proof of GCH, Gödel's method introduced the important concept of *inner models*; an interested reader can learn more about this subject in the article [35], in this issue of the Bulletin.

In 1963, Cohen proved the independence of the continuum hypothesis [7], [8]. Cohen constructed a model of ZFC in which CH fails. Moreover, Cohen's method of *forcing* proved to be a powerful tool for obtaining other independence results, and in particular was used to show that cardinal arithmetic of *regular* cardinals can behave arbitrarily, within the limits imposed by König's Theorem. Shortly after Cohen's breakthrough, Solovay showed that  $2^{\aleph_0}$  can take any value not excluded by König's theorem, i.e., one can have  $2^{\aleph_0} = \aleph_\alpha$  for any  $\alpha$  as long as  $\text{cf } \aleph_\alpha > \aleph_0$ . Then Easton produced a model [13] in which the function  $2^{\aleph_\alpha}$ , for regular  $\aleph_\alpha$ , can behave in any prescribed way consistent with König's theorem.

The natural question arose whether the freedom enjoyed by the function  $2^{\aleph_\alpha}$  on regular cardinals extends as well to singular cardinals. In particular, can a singular cardinal be the least cardinal at which GCH fails? Or, specifically, is it possible to have  $2^{\aleph_n} = \aleph_{n+1}$  for every  $n$  while  $2^{\aleph_w} = \aleph_{w+2}$ ? These questions (first asked by Solovay in the mid-sixties, see [43]) are a part of what has become known as the *Singular Cardinals Problem*.

At first the consensus among set theorists was that an improvement in Cohen's method will lead to a general consistency result along the lines of Easton's theorem. The truth however turned out to be much more interesting.

**§4. The singular cardinals problem.** By Easton's theorem, the only rules for the continuum function  $2^{\aleph_\alpha}$  on regular cardinals that are provable in ZFC state that  $2^{\aleph_\alpha} \leq 2^{\aleph_\beta}$  if  $\alpha \leq \beta$ , and that  $\text{cf } 2^{\aleph_\alpha} > \aleph_\alpha$ . The situation is different for singular cardinals. If  $\aleph_\alpha$  is singular then an easy use of cardinal arithmetic shows that  $2^{\aleph_\alpha} = (\sum_{\gamma < \alpha} 2^{\aleph_\gamma})^{\text{cf } \aleph_\alpha}$ . As a consequence, if  $\aleph_\alpha$  is singular and if  $2^{\aleph_\gamma}$  has the same value  $2^{\aleph_{\gamma_0}}$  for eventually all  $\gamma < \alpha$  (i.e.,  $\gamma_0 \leq \gamma < \alpha$ ) then  $2^{\aleph_\alpha} = 2^{\aleph_{\gamma_0}}$  as well. Thus additional rules are needed if one hopes for an extension of Easton's theorem. This was observed, e.g., in [3] and [27].

For cardinal arithmetic on singular cardinals, it is not the continuum function  $2^{\aleph_\alpha}$  alone that determines all exponentiation. It turns out that it is the function  $\aleph_\alpha^{\text{cf } \aleph_\alpha}$  that is crucial for cardinal arithmetic. (Note that if  $\aleph_\alpha$  is regular then  $\aleph_\alpha^{\text{cf } \aleph_\alpha} = \aleph_\alpha^{\aleph_\alpha} = 2^{\aleph_\alpha}$ ; for singular cardinals we only have  $\aleph_\alpha^{\text{cf } \aleph_\alpha} \leq 2^{\aleph_\alpha}$ .)

The knowledge of the function  $\aleph_\alpha^{\text{cf } \aleph_\alpha}$  is sufficient for all cardinal exponentiation. That all cardinal arithmetic, including infinite products  $\prod$ , can be

reduced to the function  $\aleph_\alpha^{\text{cf } \aleph_\alpha}$  was already known to Gödel who in [26] so remarked pointing to Tarski's work [69].

The continuum function is determined as follows: If  $\aleph_\alpha$  is regular then  $2^{\aleph_\alpha} = \aleph_\alpha^{\text{cf } \aleph_\alpha}$ . So let  $\aleph_\alpha$  be singular. If  $2^{\aleph_\gamma} = 2^{\aleph_{\gamma_0}}$  for eventually all  $\gamma < \alpha$ , then  $2^{\aleph_\alpha} = 2^{\aleph_{\gamma_0}}$ . If the continuum function  $2^{\aleph_\gamma}$  is not eventually constant below  $\aleph_\alpha$  then  $2^{\aleph_\alpha} = \kappa^{\text{cf } \kappa}$  where  $\kappa = \sum_{\gamma < \alpha} 2^{\aleph_\gamma}$ .

The exponentiation  $\aleph_\alpha^{\aleph_\beta}$  is then determined from the continuum function and the function  $\kappa^{\text{cf } \kappa}$  inductively, the crucial case being when  $\text{cf } \aleph_\alpha \leq \aleph_\beta < \aleph_\alpha$  and  $\sum_{\gamma < \alpha} \aleph_\gamma^{\aleph_\beta} = \aleph_\alpha$ . Then we have  $\aleph_\alpha^{\aleph_\beta} = \aleph_\alpha^{\text{cf } \aleph_\alpha}$ .

Thus the key to cardinal arithmetic is the function  $\kappa^{\text{cf } \kappa}$ . A consequence of König's theorem is that  $\kappa^{\text{cf } \kappa} \neq \kappa$  and so  $\kappa^{\text{cf } \kappa} \geq \kappa^+$ . Of course if  $2^{\text{cf } \kappa} \geq \kappa$  then  $\kappa^{\text{cf } \kappa} = 2^{\text{cf } \kappa}$ . Hence what is decisive (in addition to the continuum function on regulars) is the behavior of  $\kappa^{\text{cf } \kappa}$  for those  $\kappa$  for which  $2^{\text{cf } \kappa} < \kappa$ . The simplest possibility is when

$$2^{\text{cf } \kappa} < \kappa \quad \text{implies} \quad \kappa^{\text{cf } \kappa} = \kappa^+.$$

This is known as the *Singular Cardinal Hypothesis* (SCH).

Assuming SCH, the continuum function on regular cardinals itself determines cardinal exponentiation: if  $\aleph_\alpha$  is singular then  $2^{\aleph_\alpha}$  is either  $\kappa$  or  $\kappa^+$  where  $\kappa = \sum_{\gamma < \alpha} 2^{\aleph_\gamma}$ , depending on whether  $2^{\aleph_\gamma}$  is or is not eventually constant. Similarly for  $\aleph_\alpha^{\aleph_\beta}$  where  $\text{cf } \aleph_\alpha \leq \aleph_\beta < \aleph_\alpha$ .

With this analysis of cardinal arithmetic it is now clear that the fundamental question related to the singular cardinals problem is whether SCH can fail. In the simplest case mentioned in Section 3, we can ask whether SCH can fail for  $\aleph_\omega$ : can we have  $2^{\aleph_0} < \aleph_\omega$  and at the same time  $\aleph_\omega^{\aleph_0} \geq \aleph_{\omega+2}$ ? (This is a somewhat finer question than whether  $\aleph_\omega$  can be a strong limit cardinal while  $2^{\aleph_\omega} \geq \aleph_{\omega+2}$ ; if  $2^{\aleph_n} < \aleph_\omega$  for all  $\aleph_n$  then  $2^{\aleph_\omega} = \aleph_\omega^{\aleph_0}$ .)

Using basic cardinal arithmetic and König's theorem, it is possible to derive several additional rules for the behavior of the function  $\kappa^{\text{cf } \kappa}$  (see [28]). For instance, if  $\kappa$  is strong limit, then  $\text{cf}(\kappa^{\text{cf } \kappa}) > \kappa$ ; or if  $\kappa \leq \lambda^{\text{cf } \lambda}$  for some  $\lambda < \kappa$  with  $\text{cf } \lambda \geq \text{cf } \kappa$ , then  $\kappa^{\text{cf } \kappa} \leq \lambda^{\text{cf } \lambda}$ . It turns out however that more dramatic restrictions are in store for arithmetic of singular cardinals.

**§5. Silver's theorem and Jensen's covering theorem.** Until 1974 most set theorists believed that the restriction to regular cardinals in Easton's theorem was due to the weakness of its proof and that analogous results for singular cardinals would be forthcoming. In particular, it was expected that it was possible for a singular cardinal to be the least counterexample to GCH.

This changed dramatically when Silver proved the following theorem [67]:

*If  $\kappa$  is a singular cardinal of uncountable cofinality, and if  $2^\lambda = \lambda^+$  for all  $\lambda < \kappa$ , then  $2^\kappa = \kappa^+$ .*

Following Silver's result, several theorems appeared that further restricted the behavior of the function  $2^\kappa$  for singular cardinals of uncountable cofinality, most notably the theorem of Galvin and Hajnal [15].

*If  $\aleph_\lambda$  is a strong limit cardinal of uncountable cofinality then  $2^{\aleph_\lambda} < \aleph_{(2^{|\lambda|})^+}$ .*

In another direction, Jensen was able to combine Silver's ideas with his previous work [34] on the fine structure of  $L$ , to prove his remarkable Covering Theorem [10]:

*If  $0^\#$  does not exist then every uncountable set of ordinals can be covered by a constructible set of the same cardinality.*

(The existence of  $0^\#$  is a large cardinal axiom that we shall return to in the next section.) An easy argument using the Covering Theorem shows that unless  $0^\#$  exists,  $2^{\text{cf } \kappa} < \kappa$  implies  $\kappa^{\text{cf } \kappa} = \kappa^+$ , i.e., SCH holds. Thus in order to violate SCH we need large cardinals.

A related corollary of the Covering Theorem states that if  $0^\#$  does not exist then for every  $\lambda \geq \aleph_2$ , if  $\lambda$  is a regular cardinal in  $L$  then  $\text{cf } \lambda = |\lambda|$ . In particular, a regular cardinal cannot be changed into a singular cardinal in the absence of large cardinals. (The assumption  $\lambda \geq \aleph_2$  is necessary, as Bukovský [4] and Namba [46] produced a forcing extension of  $L$ —in which  $0^\#$  cannot exist—where  $|\omega_2^L| = \aleph_1$  and  $\text{cf } \omega_2^L = \omega$ .)

The assumption of Silver's Theorem that  $\kappa$  has uncountable cofinality figures prominently in the proof. To let the reader appreciate the significance of uncountable cofinality, I shall give a brief outline of the methods used in Silver's and Galvin-Hajnal's theorems.

For simplicity, let us consider  $\aleph_{\omega_1}$ . For Silver's Theorem, assume that  $2^{\aleph_\alpha} = \aleph_{\alpha+1}$  for all  $\alpha < \omega_1$ . The main idea of the proof is that there exists a generic extension  $V[G]$  of the universe, in which there exists a *normal* ultrafilter  $U$  on  $P^V(\omega_1)$ . Normality means, as usual, that the diagonal is the least nonconstant function; here the fact that  $\omega_1$  is uncountable is essential. Working in  $V[G]$ , we can calculate the size of  $(2^{\aleph_{\omega_1}})^V$  as follows. Consider the ultrapower  $M$  of  $V$  by  $U$ , and the elementary embedding  $j: V \rightarrow M$ .  $M$  is not well founded, but thanks to normality, the  $M$ -ordinal  $\kappa$  represented by the function  $\alpha \mapsto \aleph_\alpha$  is the supremum of the  $M$ -ordinals  $j(\aleph_\gamma)$ ,  $\gamma < \omega_1$ . It follows that the linearly ordered set  $(\kappa^+)^M$  has size at most  $((\aleph_{\omega_1})^V)^+$ . A further argument gives  $|P^V(\aleph_{\omega_1})| \leq |P^M(\kappa)|$ , and since  $M \models 2^\kappa = \kappa^+$ , we have  $|(2^{\aleph_{\omega_1}})^V| \leq ((\aleph_{\omega_1})^V)^+$ . This is calculated in  $V[G]$ , but since the forcing extension is "mild," the same inequality holds in  $V$ .

Silver's proof has been reworked in [1], replacing the forcing technique by a purely combinatorial argument. A further improvement of the combinatorial method led to the above mentioned theorem of Galvin and Hajnal. In the Galvin-Hajnal theorem, the uncountability of the cofinality is again

essential. Roughly speaking, the generic ultrapower of Silver is replaced by a well-founded partial ordering of ordinal-valued functions. If  $f$  and  $g$  are ordinal functions on  $\omega_1$ , let  $f < g$  denote the relation

$$\{ \alpha < \omega_1 : f(\alpha) < g(\alpha) \} \text{ contains a club.}$$

Due to normality of the club filter, the relation  $<$  is a well-founded partial ordering. Hence every ordinal function  $f$  on  $\omega_1$  can be assigned its rank in  $<$ , the *Galvin-Hajnal norm*  $\|f\|$  of  $f$ . The essence of Galvin-Hajnal's proof is that the size of  $2^{\aleph_{\omega_1}}$  (if  $\aleph_{\omega_1}$  is strong limit) is related to the Galvin-Hajnal norm of the function  $\alpha \mapsto 2^{\aleph_\alpha}$ . The analysis of this relationship yields the upper bound  $2^{\aleph_{\omega_1}} < \aleph_{(2^{\aleph_1})^+}$ .

**§6. Large cardinals.** As the singular cardinals problem is so closely related to large cardinal axioms, we shall now review the basics of the theory of large cardinals.

An uncountable cardinal number  $\kappa$  is *inaccessible* if it is regular and a strong limit cardinal. An immediate consequence of inaccessibility is that  $V_\kappa$ , the collection of all sets of rank less than  $\kappa$ , is a model of ZFC; another immediate consequence is that  $\kappa = \aleph_\kappa$  is a fixed point of the aleph sequence. By Gödel's 2nd Incompleteness Theorem it follows that the existence of inaccessible cardinals is unprovable in ZFC. In fact, a slightly more involved argument shows that the relative consistency of inaccessible cardinals is unprovable. Thus the existence of inaccessibles is to ZFC as the existence of an infinite set is to Peano arithmetic. For that reason, large cardinal axioms are sometimes referred to as *strong axioms of infinity*.

Modern large cardinal theory recognizes a substantial number of large cardinal axioms. Interestingly enough, these axioms form a linearly ordered scale, on which the relation of a stronger axiom to the weaker theories is just as described above in the case of inaccessible cardinals and ZFC. This scale of large cardinals serves as a measure of consistency strength of various set theoretic assumptions (including assumptions on singular cardinals).

One of the most prominent large cardinals is a measurable cardinal. Measurable cardinals were introduced by Ulam while working on the *measure problem*, the question whether some set  $E$  can carry a nontrivial countably additive measure such that every subset of  $E$  is measurable. An uncountable cardinal  $\kappa$  is *measurable* if  $\kappa$  carries an atomless  $\kappa$ -additive 2-valued measure; equivalently, if  $\kappa$  carries a nonprincipal  $\kappa$ -complete ultrafilter. (The terms " $\kappa$ -additive" and " $\kappa$ -complete" refer to unions and intersections of fewer than  $\kappa$  sets.) In [50], Scott applied to measurable cardinals the method of ultraproducts, thus laying the groundwork for the modern large cardinal theory. One of the basic observations is that the existence of measurable


cardinals is equivalent to the existence of a nontrivial elementary embedding  $j: V \rightarrow M$  where  $M$  is some transitive model. Every measurable cardinal is not only inaccessible, but on the scale of large cardinals is above Mahlo, weakly compact and Ramsey cardinals, to mention just the most important categories of large cardinals (see picture).

Scott’s theorem [50] states that measurable cardinals don’t exist in  $L$ . The subsequent work of Gaifman, Rowbottom [49] and Silver [66] established the strength of measurable cardinals *vis à vis* constructible sets, isolating the principle “ $0^\#$  exists.” The existence of  $0^\#$  (between weakly compact and Ramsey on our scale) states that there aren’t many constructible sets, and the truth definition for the model  $L$  exists. Equivalently, there exists a nontrivial elementary embedding  $j: L \rightarrow L$ . One consequence of  $0^\#$  is that every uncountable cardinal (in  $V$ ) is inaccessible in  $L$ . In particular,  $\aleph_\omega$  is regular in  $L$ ; recall that by the Covering Theorem, if  $0^\#$  does not exist then  $\aleph_\omega$  (and every singular cardinal) must be singular in  $L$ .

Among large cardinals stronger than measurable, *supercompact* cardinals are most prominent. There exists at present a good classification of large cardinals between measurables and *Woodin cardinals*, using the theory of *inner models* (we refer the reader to Jensen’s article [35], in this Bulletin). In particular, measurable cardinals are classified by their *order*  $o(\kappa)$ , where  $o(\kappa) = 1$  means that  $\kappa$  is measurable,  $o(\kappa) = 2$  means that  $\kappa$  carries a normal measure in which almost all  $\alpha < \kappa$  are measurable, and so on, up to  $o(\kappa) = \kappa^{++}$ .

Above supercompact and huge cardinals, the scale approaches its end with the existence of an elementary embedding  $j: V_\lambda \rightarrow V_\lambda$ , as by a theorem of Kunen,  $j: V \rightarrow V$  is inconsistent.

For the benefit of nonspecialists, as well as a reference for the forthcoming sections we present a scale of the more prominent large cardinals:



**§7. Large cardinals and the singular cardinals problem.** Prior to Jensen’s Covering Theorem there had been scattered results indicating that there might be some link between the behavior of singular cardinals and the large cardinal axioms. For instance, Solovay proved in [68] that if  $\kappa$  is a supercompact cardinal and  $\lambda > \kappa$  is singular then  $\lambda^{\aleph_0} \leq \lambda^+$ . This means that the SCH holds above the least supercompact cardinal. On the other hand, Prikry introduced in [47] a method of forcing that changes the cofinality of a measurable cardinal to  $\omega$  without collapsing it. The measurable cardinal  $\kappa$

remains a cardinal but has  $\text{cf } \kappa = \omega$  in the extension. Subsequently, Silver devised a forcing method that, starting with a supercompact cardinal  $\kappa$ , produced a model in which  $\kappa$  is measurable and  $2^\kappa > \kappa^+$ . This, combined with the Prikry forcing, yields the consistency of the negation of SCH, relative to a supercompact cardinal: a model of ZFC in which  $\kappa$  is a strong limit cardinal,  $\text{cf } \kappa = \omega$ , and  $2^\kappa > \kappa^+$ .

About the same time that Jensen established, by the Covering Theorem, the necessity of large cardinals for the negation of the SCH, Magidor obtained the first in a series of consistency results using large cardinals. In [40], he proved the consistency of  $2^{\aleph_\omega} > \aleph_{\omega+1}$  (and  $\aleph_\omega$  strong limit) from a supercompact cardinal, and in [41] the consistency of  $2^{\aleph_\omega} = \aleph_{\omega+2}$  along with GCH below  $\aleph_\omega$ , from a 2-huge cardinal. When it soon became clear that large cardinals are indeed necessary, Magidor's method was refined to yield other consistency results. The general idea in all these proofs is as follows: start with a large cardinal  $\kappa$ , blow up  $2^\kappa$ , change the cofinality of  $\kappa$  by adjoining a new set of cardinals  $C$  cofinal in  $\kappa$ , and destroy all cardinals below  $\kappa$  that are not in the set  $C$ . The technique employed to change the cofinality evolved from Prikry's method, progressing through [42] to [48]. More recently, a new method for building models for the negation of SCH was developed in [20] and [22].

In Magidor's model [40],  $2^{\aleph_\omega}$  could be as large as  $\aleph_{\omega+\omega+1}$ , which was improved by Shelah [54] as to have  $2^{\aleph_\omega} = \aleph_{\alpha+1}$  for any  $\alpha < \omega_1$ . Woodin [14] constructed a model in which  $2^\kappa = \kappa^{++}$  for all infinite cardinals  $\kappa$ , and Cummings [9] found a model in which GCH fails exactly at all limit cardinals.

As for the consistency strength of the failure of SCH, a series of results proved it to be exactly a measurable cardinal of order  $o(\kappa) = \kappa^{++}$ . Magidor's assumption of supercompactness was first weakened by Woodin, and eventually Gitik obtained a model of  $\neg\text{SCH}$  in [17] using  $o(\kappa) = \kappa^{++}$ . The other direction, namely that  $o(\kappa) = \kappa^{++}$  is a necessary assumption, is also due to Gitik ([18]), and uses the technique of inner models.

The technique of inner models is an outgrowth of Jensen's Covering Theorem. The first step up from  $0^\#$  was the core model  $K$  and the covering theorem for  $K$  ([11], [12]), followed by inner models and covering theorems for measurable cardinals [44] and beyond. Details can be found in Jensen's article [35]. This technique produced a number of results, notably [45], [21], [19] and [23] showing the necessity of large cardinal axioms for various violations of SCH.

A major open problem in cardinal arithmetic is whether  $2^{\aleph_\omega}$  can be greater than  $\aleph_{\omega_1}$  (while  $\aleph_\omega$  is strong limit). The analysis provided by the pcf theory indicates that an entirely new approach would be needed, and the inner model technique shows that its consistency strength is enormous.

**§8. Upper bounds.** Following Silver's Theorem [67], one of the main directions of research in the theory of singular cardinals has been the search for upper bounds on  $2^{\aleph_\alpha}$  for strong limit singular cardinals  $\aleph_\alpha$ ; or more precisely, on the function  $\aleph_\alpha^{\text{cf } \aleph_\alpha}$  for those cardinals that satisfy  $2^{\text{cf } \aleph_\alpha} < \aleph_\alpha$ . Let us recall that the Galvin-Hajnal Theorem gives a bound

$$2^{\aleph_{\omega_1}} < \aleph_{(2^{\aleph_1})^+}$$

if  $\aleph_{\omega_1}$  is a strong limit cardinal. In full generality, if  $\aleph_\delta$  is a strong limit cardinal of uncountable cofinality then  $2^{\aleph_\delta} < \aleph_{(\aleph_\delta^{\text{cf } \delta})^+}$ .

The theorem leaves open the question whether an upper bound exists for  $2^{\aleph_\omega}$  (and other cardinals of cofinality  $\omega$ ). The method does not give any information in this case, as it depends heavily on the use of closed unbounded sets.

The other question left open is the case when  $\aleph_\delta$  is a fixed point of the aleph function, i.e., when  $\delta = \aleph_\delta$ .

Several papers extended the Galvin-Hajnal result under large cardinal assumptions: Magidor [39] proved  $2^{\aleph_{\omega_1}} < \aleph_{\omega_2}$  from Chang's Conjecture (which has the consistency strength of approximately Ramsey cardinals); Jech and Prikry [31] proved  $2^{F(\aleph_1)} < F(\aleph_2)$ , where  $F(\aleph_\alpha)$  is the  $\aleph_\alpha$ th fixed point of the  $\aleph$  function, from a saturated ideal (approximately supercompact); and Shelah [52] proved the same result from Chang's Conjecture. In [55] and [56] Shelah eliminated the large cardinal assumptions, obtaining somewhat weaker bounds, such as  $2^{F(\aleph_1)} < F((2^{2^{\aleph_1}})^+)$ . Interestingly, his method broke down for the  $\omega$ th iteration of the fixed point operation; eventually it turned out that in this case no a priori bounds are provable.

To obtain upper bounds for  $2^{\aleph_\omega}$ , a different approach is needed. The major breakthrough came when Shelah discovered the significance of cofinalities of ultraproducts  $\prod_{n=0}^{\infty} \aleph_n / D$ , where  $D$  is an ultrafilter on  $\omega$ . Making use of these, he obtained the analog of the Galvin-Hajnal Theorem: if  $\aleph_\omega$  is strong limit then

$$2^{\aleph_\omega} < \aleph_{(2^{\aleph_0})^+}$$

(cf. [53], Chapter XIII). The proof of this theorem led Shelah to a systematic study of cofinalities of reduced products of sets of cardinals. In a sequence of papers [52], [57], [58] and [61] (Chapters II, VIII and IX), he developed a beautiful theory that brought a number of unexpected results and yielded deep applications to cardinal arithmetic. The crowning achievement so far is the following result that we shall discuss in the following section: if  $2^{\aleph_0} < \aleph_\omega$ , then

$$\aleph_\omega^{\aleph_0} < \aleph_{\omega_4}.$$

**§9. Shelah's pcf theory.** In this last section we shall outline Shelah's theory of possible cofinalities (pcf). The central concept of the theory is the cofinality of an ultraproduct of a set of regular cardinals:

Let  $A$  be a set of regular cardinals, and let  $D$  be an ultrafilter on  $A$ . The ultraproduct  $\prod A/D$  consists of equivalence classes of ordinal functions  $f$  such that  $\text{dom}(f) = A$  and  $f(\alpha) \in \alpha$  for all  $\alpha \in A$ ; two functions  $f$  and  $g$  are equivalent mod  $D$  if  $\{\alpha : f(\alpha) = g(\alpha)\} \in D$ . The ultraproduct is linearly ordered by the relation  $f <_D g$  defined by  $\{\alpha : f(\alpha) < g(\alpha)\} \in D$ , and the *cofinality* of  $\prod A/D$  is the unique regular cardinal  $\kappa = \text{cof } \prod A/D$  such that  $(\prod A/D, <_D)$  has a cofinal subset of order type  $\kappa$ .

If  $\kappa = \text{cof } \prod A/D$  for some ultrafilter  $D$ , we say that  $\kappa$  is a *possible cofinality* (of  $\prod A$ ), and we let

$$\text{pcf } A = \left\{ \text{cof } \prod A/D : D \text{ an ultrafilter on } A \right\}$$

be the set of all possible cofinalities of  $\prod A$ .

As a typical case, let us consider the set  $A = \{\aleph_n\}_{n=0}^\infty$ . Of course, every  $\aleph_n$  is a possible cofinality, by a principal ultrafilter. If  $\kappa = \text{cof } \prod A/D$  where  $D$  is nonprincipal then clearly  $\kappa \geq \aleph_{\omega+1}$ , and since  $|\prod A| = \aleph_\omega^{\aleph_0}$ , we have  $\kappa \leq \aleph_\omega^{\aleph_0}$ . So unless  $\aleph_\omega^{\aleph_0} > \aleph_{\omega+1}$ , the set  $\text{pcf } A$  has only one element outside  $A$ , and so a meaningful theory of possible cofinalities requires the negation of the singular cardinals hypothesis. It turns out that the pcf theory is more fundamental than cardinal arithmetic.

In [51] Shelah proved, among others, the following result that was a precursor of the future pcf theory:

**THEOREM 9.1.** *For every singular cardinal  $\lambda$  of cofinality  $\omega$  there exists an increasing sequence  $\{\lambda_n\}_{n=0}^\infty$  of regular cardinals with limit  $\lambda$  such that  $\text{cof } \prod_{n=0}^\infty \lambda_n/D = \lambda^+$  for every nonprincipal ultrafilter  $D$ .*

This result shows, e.g., that  $\aleph_{\omega+1}$  is a possible cofinality of  $\prod_{n=0}^\infty \aleph_n$  regardless of the value of  $\aleph_\omega^{\aleph_0}$ .

Shortly thereafter, Shelah established the relation between possible cofinalities and cardinal arithmetic, and obtained an upper bound for  $2^{\aleph_\omega}$ , cf. [53]. We say that  $A$  is an *interval* of regular cardinals if whenever  $\alpha < \beta < \gamma$  are regular cardinals and  $\alpha, \gamma \in A$ , then  $\beta \in A$ . The following theorem reduces the simplest cases of the singular cardinals problem to the problem of possible cofinalities:

**THEOREM 9.2.**

- (a) *If  $A$  is an interval of regular cardinals such that  $|A| < \min A$  then  $\text{pcf } A$  is an interval.*
- (b) *If, moreover,  $(\min A)^{|A|} < \sup A$  then  $\kappa = |\prod A|$  is a possible cofinality.*

This immediately gives an upper bound on  $2^{\aleph_\omega}$  if  $\aleph_\omega$  is strong limit: since  $|\text{pcf}\{\aleph_n\}_{n=0}^\infty| \leq 2^{2^{\aleph_0}}$  (the number of ultrafilters on a countable set), and since we assume that  $2^{2^{\aleph_0}} < \aleph_\omega$ , we have  $2^{\aleph_\omega} = |\prod_{n=0}^\infty \aleph_n| < \aleph_{\aleph_\omega}$ . It also reduces the problem of evaluating  $\aleph_\omega^{\aleph_0}$  (and similar products) to finding the maximum of possible cofinalities of  $\prod_{n=0}^\infty \aleph_n$ . (There are analogous reductions for cases not covered by Theorem 9.2.) Notice another consequence of Theorem 9.2: as  $|\prod A|$  is a possible cofinality, it must be a regular cardinal. Hence if  $\aleph_\omega$  is a strong limit,  $2^{\aleph_\omega}$  is regular. This does not follow from König's Theorem.

If  $2^{\aleph_0} < \aleph_\omega$  then all regular cardinals between  $\aleph_{\omega+1}$  and  $\aleph_\omega^{\aleph_0}$  are possible cofinalities of  $\prod_{n=0}^\infty \aleph_n$ . It is interesting to see how they are attained in the models where  $2^{\aleph_\omega} > \aleph_{\omega+1}$ . A detailed analysis of this was done in [29] for Magidor's model [40]; a similar analysis is possible for other models. Let  $\kappa_n$ ,  $n < \omega$ , be the cardinals in the Prikry sequence; e.g., in the model where  $2^{\aleph_\omega} = \aleph_{\omega+2}$  these are  $\kappa_n = \aleph_{3n}$ . Then  $\text{cof} \prod_{n < \omega} \kappa_n^+ / D = \aleph_{\omega+1}$  and  $\text{cof} \prod_{n < \omega} \kappa_n^{++} / D = \aleph_{\omega+2}$ , for every nonprincipal ultrafilter. Hence  $\prod_n \aleph_{3n+1}$  has cofinality  $\aleph_{\omega+1}$  while  $\prod_n \aleph_{3n+2}$  has cofinality  $\aleph_{\omega+2}$ !

**Fundamentals of the pcf theory.** The theory developed by Shelah in [51], [57] and [58], and described in detail in the monograph [61] analyzes possible cofinalities of products  $\prod A$  where  $A$  is a set of regular cardinals with the property that  $|A| < \min A$ . (For rather trivial reasons, the general pcf theory breaks down in the case when  $\sup A$  is a fixed point of the aleph function.) We shall now present the main points of the theory; the reader interested in proofs of the theorems stated here and in the techniques involved might find either [5] or [30] helpful.

Let  $A$  be a set of regular cardinals and assume that  $|A| < \min A$ , and let

$$\text{pcf } A = \left\{ \text{cof} \prod A / D : D \text{ an ultrafilter on } A \right\}.$$

First we mention the trivial facts about pcf:

- $A \subseteq \text{pcf } A$ ,
- if  $A_1 \subseteq A_2$  then  $\text{pcf } A_1 \subseteq \text{pcf } A_2$ ,
- $\text{pcf}(A_1 \cup A_2) = \text{pcf } A_1 \cup \text{pcf } A_2$ .

Another not so difficult observation is that when  $|\text{pcf } A| < \min A$  then  $\text{pcf } \text{pcf } A = \text{pcf } A$ .

The key result of the pcf theory is the following:

**THEOREM 9.3 (Existence of generators).** *There exists a collection  $\{B_v : v \in \text{pcf } A\}$  of subsets of  $A$  such that for every  $v$ ,*

- (i)  $v = \max \text{pcf } B_v$ ,
- (ii)  $v \notin \text{pcf}(A - B_v)$ .

(The sets  $B_\nu$  are called *generators* of pcf  $A$ .)

In other words:

- (i) for every ultrafilter  $D$  on  $B_\nu$ ,  $\text{cof} \prod A/D \leq \nu$ ; and there exists some  $D$  on  $B_\nu$  such that  $\text{cof} \prod A/D = \nu$ ,
- (ii) for every  $D$ , if  $\text{cof} \prod A/D = \nu$  then  $B_\nu \in D$ .

Therefore, the cofinality of  $\prod A/D$  is determined by which generators belong to  $D$ :

$$\text{cof} \prod A/D = \text{the least } \lambda \text{ such that } B_\lambda \in D.$$

Let us note some consequences of Theorem 9.3:

- (iii)  $|\text{pcf } A| \leq 2^{|A|}$ .

This is because  $|\text{pcf } A|$  does not exceed the number of generators, which is at most  $2^{|A|}$ . An immediate consequence is a better bound on  $2^{\aleph_\omega}$  if  $\aleph_\omega$  is a strong limit:  $2^{\aleph_\omega} < \aleph_{(2^{\aleph_\omega})^+}$ .

- (iv) pcf  $A$  has a maximal element.

If not, then  $\{A - B_\nu : \nu \in \text{pcf } A\}$  has the finite intersection property; let  $D$  be an ultrafilter extending it, and let  $\lambda = \text{cof} \prod A/D$ . Then  $B_\lambda \in D$ , a contradiction.

The same argument proves the following important property of generators:

**THEOREM 9.4 (Compactness).** *For every  $X \subseteq A$  there exists a finite set of cardinals  $\nu_1, \dots, \nu_k \in \text{pcf } X$  such that  $X \subseteq B_{\nu_1} \cup \dots \cup B_{\nu_k}$ .*

**Transitive generators and the localization theorem.** Generators of pcf  $A$  are not necessarily unique, as (e.g.) changing  $B_\nu$  by a finite set won't affect properties (i) or (ii). The  $B_\nu$ 's are however unique up to the  $B_\eta$ 's for  $\eta < \nu$ . More precisely: for every  $\kappa \leq \max \text{pcf } A$ , let

$$J_\kappa = \text{the ideal (of subsets of } A) \text{ generated by the sets } B_\eta, \\ \text{for } \eta < \kappa \text{ and } \eta \in \text{pcf } A.$$

Properties (i) and (ii) imply that for every  $X \subseteq A$ ,

$$X \in J_\kappa \quad \text{if and only if} \quad \text{cof} \prod X/D < \kappa \text{ for every ultrafilter } D \text{ on } X,$$

and that each generator  $B_\nu$  is unique mod  $J_\nu$ .

An important feature of the pcf theory is that pcf  $A$  has *transitive* generators (incidentally, this theorem is highly nontrivial):

**THEOREM 9.5.** *There exist generators  $B_\nu$ ,  $\nu \in \text{pcf } A$ , such that whenever  $\eta \in B_\nu$  then  $B_\eta \subseteq B_\nu$ .*

Transitivity of generators is particularly useful when applied to sets  $A = \text{pcf } A$ . A fairly straightforward use of transitivity and compactness yields the following important result:

**THEOREM 9.6 (Localization Theorem).** *Let  $X \subseteq \text{pcf } A$  and let  $\lambda \in \text{pcf } X$ . Then  $X$  has a subset  $X_0$  such that  $|X_0| \leq |A|$ , and such that  $\lambda \in \text{pcf } X_0$ .*

**An upper bound for  $|\text{pcf } A|$ .** The strength of the Localization Theorem is best illustrated by its application providing an upper bound on the size of  $\text{pcf } A$ , and consequently, on the function  $2^{\aleph_\alpha}$  for singular  $\aleph_\alpha$ . We shall now outline this application of the pcf theory.

First we should mention that one application of the techniques used in the pcf theory is the following analog of Theorem 9.1 for uncountable cofinality:

**THEOREM 9.7.** *If  $\lambda$  is a singular cardinal of uncountable cofinality then there exists a closed unbounded set  $C \subseteq \lambda$  such that  $\text{cof } \prod_{\alpha \in C} \alpha^+ / D = \lambda^+$  for every ultrafilter  $D$  concentrating on end-segments of  $C$ .*

Now let  $A = \{\aleph_n\}_{n=0}^\infty$ . By theorem 9.2,  $\text{pcf } A$  is an interval of regular cardinals and, if  $2^{\aleph_0} < \aleph_\omega$  then  $\max \text{pcf } A = \aleph_\omega^{\aleph_0}$ . We claim that  $\max \text{pcf } A < \aleph_{\omega_4}$ , and therefore  $\aleph_\omega^{\aleph_0} < \aleph_{\omega_4}$ .

The structure of  $\text{pcf } A$  induces a closure operation on  $\Theta$ , where  $\aleph_{\Theta+1} = \max \text{pcf } A$ : if  $X \subseteq \Theta$ , we let

$$\alpha \in \bar{X} \quad \text{if and only if} \quad \aleph_{\alpha+1} \in \text{pcf} \{ \aleph_{\xi+1} : \xi \in X \}.$$

From Theorem 9.7 it follows that if  $\lambda < \Theta$  is an ordinal of uncountable cofinality then there exists a club  $C \subseteq \lambda$  such that  $\sup C = \lambda$ . From the Localization Theorem it follows that every set  $X \subseteq \Theta$  of order type  $\omega_1$  has a countable initial segment  $X_0$  such that  $\sup \bar{X}_0 \geq \sup X$ .

Shelah now proceeds to show that if  $\Theta \geq \aleph_4$  then no closure operation on  $\Theta$  with those two properties exists. Therefore  $\Theta < \aleph_4$ , and  $\max \text{pcf} \{ \aleph_n \}_{n=0}^\infty < \aleph_{\omega_4}$ !

There are some elements of the proof that are specific to  $\aleph_4$ , making it impossible at present to bring down the upper bound to, say,  $\aleph_{\omega_3}$ . The method definitely does not work for  $\aleph_{\omega_1}$ , cf. [33], but it is still an open problem whether  $2^{\aleph_\omega}$  can be greater than  $\aleph_{\omega_1}$  (with  $\aleph_\omega$  strong limit). In fact, it is still unknown whether  $|A| < |\text{pcf } A|$  is possible (together with  $|A| < \min A$ ).

**Reduced products of ordinals.** The main technique used in the analysis of possible cofinalities involves reduced products of ordinal numbers. We shall conclude the article with a brief description of this topic.

Let  $A$  be an infinite set, and let  $I$  be an ideal on  $A$ . For ordinal functions  $f, g$  on  $A$  we define:

$$f =_I g \quad \text{if} \quad \{a \in A : f(a) \neq g(a)\} \in I,$$

$$f <_I g \quad \text{if} \quad \{a \in A : f(a) \geq g(a)\} \in I.$$

Let  $\lambda_a$ ,  $a \in A$ , be limit ordinal numbers. The *reduced product*

$$\prod_{a \in A} \lambda_a / I$$

consists of equivalence classes of ordinal functions  $f \bmod =_I$ , such that  $f(a) \in \lambda_a$  for all  $a \in A$ .

Let  $\kappa$  be a regular cardinal. We say that  $P = \prod_{a \in A} \lambda_a / I$  is  $\kappa$ -directed if every subset  $X$  of  $P$  of size less than  $\kappa$  has an upper bound in  $P$ , i.e., a function  $f \in P$  such that  $g <_I f$  for every  $g \in X$ . We say that  $P$  has *true cofinality*  $\kappa$  if there exists an increasing sequence  $\langle f_\alpha : \alpha < \kappa \rangle$  cofinal in  $P$ , i.e.,  $f_0 <_I f_1 <_I \dots$ , and for every  $g \in P$  there is an  $\alpha$  such that  $g <_I f_\alpha$ . Note that if  $P$  has true cofinality  $\kappa$  then  $P$  is  $\kappa$ -directed.

We recall that for a given set  $A$  of regular cardinals,  $J_\kappa$  ( $\kappa \leq \max \text{pcf } A$ ) denotes the ideal on  $A$  generated by the  $B_\nu$ ,  $\nu < \kappa$ . The crucial property of the generators which enables the analysis of  $\text{pcf } A$  is this:

**THEOREM 9.8.** *If  $A$  is a set of regular cardinals such that  $|A| < \min A$  and if  $\kappa \in \text{pcf } A$  then  $\prod_{\nu \in B_\kappa} \nu / J_\kappa$  has true cofinality  $\kappa$ .*

In fact, the technology provided by the pcf theory goes beyond analyzing the ideals  $J_\kappa$ . Perhaps the most general result on reduced products is the following. (This is explicitly stated for  $\kappa > 2^{|A|}$  in Lemma 2.4 of [30], and can be proved for  $\kappa > |A|^+$  as in Theorems 7.3 and 7.9 of [5]):

**THEOREM 9.9 (Splitting Theorem).** *Let  $I$  be an ideal on  $A$ , let  $\lambda_a$ ,  $a \in A$ , be limit ordinals of cofinality greater than  $|A|^+$  and let  $\kappa$  be a regular cardinal,  $\kappa > |A|^+$ . If  $\prod_{a \in A} \lambda_a / I$  is  $\kappa$ -directed, then either  $\prod_{a \in A} \lambda_a / I$  is  $\kappa^+$ -directed, or it has true cofinality  $\kappa$ , or there exist  $A_1, A_2 \notin I$ ,  $A_1 \cup A_2 = A$ , such that  $\prod_{a \in A_1} \lambda_a / I$  is  $\kappa^+$ -directed and  $\prod_{a \in A_2} \lambda_a / I$  has true cofinality  $\kappa$ .*

#### REFERENCES

- [1] J. E. BAUMGARTNER and K. PRIKRY, *On a theorem of Silver*, **Discrete Mathematics**, vol. 14 (1976), pp. 17–21.
- [2] ———, *Singular cardinals and the generalized continuum hypothesis*, **American Mathematical Monthly**, vol. 84 (1977), pp. 108–113.
- [3] L. BUKOVSKÝ, *The continuum problem and the powers of alephs*, **Commentationes Mathematicae Universitatis Carolinae**, vol. 6 (1965), pp. 181–197.
- [4] ———, *Changing cofinality of  $\aleph_2$* , **Set theory and hierarchy theory** (W. Marek et al., editors), Lecture Notes in Mathematics, vol. 537, Springer-Verlag, Berlin, 1976, pp. 37–49.

- [5] M. BURKE and M. MAGIDOR, *Shelah's pcf theory and its applications*, *Annals of Pure and Applied Logic*, vol. 50 (1990), pp. 207–254.
- [6] G. CANTOR, *Über unendliche, lineare Punktmannigfaltigkeiten*, *Mathematische Annalen*, vol. 21 (1883), pp. 545–591.
- [7] P. J. COHEN, *The independence of the continuum hypothesis I*, *Proceedings of the National Academy of Sciences, U.S.A.*, vol. 50 (1963), pp. 1143–1148.
- [8] ———, *The independence of the continuum hypothesis II*, *Proceedings of the National Academy of Sciences, U.S.A.*, vol. 51 (1964), pp. 105–110.
- [9] J. CUMMINGS, *A model in which GCH holds at successors but fails at limits*, *Transactions of the American Mathematical Society*, vol. 329 (1992), pp. 1–39.
- [10] K. DEVLIN and R. JENSEN, *Marginalia to a theorem of Silver*, *Logic Conference, Kiel 1974, Proceedings* (G. H. Müller et al., editors), Lecture Notes in Mathematics, vol. 499, Springer-Verlag, Berlin, 1975, pp. 115–142.
- [11] A. J. DODD and R. B. JENSEN, *The core model*, *Annals of Mathematical Logic*, vol. 20 (1981), pp. 43–75.
- [12] ———, *The covering lemma for  $K$* , *Annals of Mathematical Logic*, vol. 22 (1982), pp. 1–30.
- [13] W. B. EASTON, *Powers of regular cardinals*, *Annals of Mathematical Logic*, vol. 1 (1970), pp. 139–178.
- [14] M. FOREMAN and W. H. WOODIN, *The generalized continuum hypothesis can fail everywhere*, *Annals of Mathematics*, vol. 133 (1991), pp. 1–35.
- [15] F. GALVIN and A. HAJNAL, *Inequalities for cardinal powers*, *Annals of Mathematics*, vol. 101 (1975), pp. 491–498.
- [16] M. GITIK, *Changing cofinalities and the nonstationary ideal*, *Israel Journal of Mathematics*, vol. 56 (1986), pp. 280–314.
- [17] ———, *The negation of the singular cardinal hypothesis from  $\mathfrak{o}(\kappa) = \kappa^{++}$* , *Annals of Pure and Applied Logic*, vol. 43 (1989), pp. 209–234.
- [18] ———, *The strength of the failure of the singular cardinal hypothesis*, *Annals of Pure and Applied Logic*, vol. 51 (1991), pp. 215–240.
- [19] ———, *On measurable cardinals violating the continuum hypothesis*, *Annals of Pure and Applied Logic*, vol. 63 (1993), pp. 227–240.
- [20] ———, *Blowing up the power of a singular cardinal*, *Annals of Pure and Applied Logic*, to appear.
- [21] M. GITIK and M. MAGIDOR, *The singular cardinals hypothesis revisited*, *Set theory and the continuum* (H. Judah et al., editors), Springer-Verlag, Berlin, 1992, pp. 243–279.
- [22] ———, *Extender based forcing*, *The Journal of Symbolic Logic*, vol. 59 (1994), pp. 445–460.
- [23] M. GITIK and W. J. MITCHELL, *Indiscernible sequences for extenders, and the singular cardinal hypothesis*, to appear.
- [24] M. GITIK and S. SHELAH, *On certain indestructibility of strong cardinals and a question of Hajnal*, *Archive for Mathematical Logic*, vol. 28 (1989), pp. 35–42.
- [25] K. GÖDEL, *Consistency-proof for the generalized continuum hypothesis*, *Proceedings of the National Academy of Sciences, U.S.A.*, vol. 25 (1939), pp. 220–224.
- [26] ———, *What is Cantor's continuum problem?*, *American Mathematical Monthly*, vol. 54 (1947), pp. 515–525.
- [27] S. H. HECHLER, *Powers of singular cardinals and a strong form of the negation of the generalized continuum hypothesis*, *Zeitschrift für Mathematische Logik*, vol. 19 (1973), pp. 83–84.
- [28] T. JECH, *Properties of the gimel function and a classification of singular cardinals*, *Fundamenta Mathematicae*, vol. 81 (1973), pp. 57–64.

- [29] ———, *On the cofinality of countable products of cardinal numbers, A tribute to Paul Erdős* (A. Baker et al., editors), Cambridge University Press, Cambridge, 1990, pp. 289–305.
- [30] ———, *Singular cardinal problem: Shelah's theorem on  $2^{\aleph_\omega}$* , *Bulletin of the London Mathematical Society*, vol. 24 (1992), pp. 127–139.
- [31] T. JECH and K. PRIKRY, *On ideals of sets and the power set operation*, *Bulletin of the American Mathematical Society*, vol. 82 (1976), pp. 593–595.
- [32] T. JECH and S. SHELAH, *On a conjecture of Tarski on products of cardinals*, *Proceedings of the American Mathematical Society*, vol. 112 (1991), pp. 1117–1124.
- [33] ———, *Possible pcf algebras*, *The Journal of Symbolic Logic*, to appear.
- [34] R. B. JENSEN, *The fine structure of the constructible hierarchy*, *Annals of Mathematical Logic*, vol. 4 (1972), pp. 229–308.
- [35] ———, *Inner models and large cardinals*, this BULLETIN, vol. 1 (1995), pp. xxx–xxxx.
- [36] A. KANAMORI, *Higher set theory*, Springer-Verlag, Berlin, 1994.
- [37] J. KÖNIG, *Zum Kontinuumproblem*, *Mathematische Annalen*, vol. 60 (1905), pp. 177–180.
- [38] K. KUNEN, *On the GCH at measurable cardinals*, *Logic Colloquium '69* (R. O. Gandy and C. E. M. Yates, editors), North-Holland, Amsterdam, 1971, pp. 107–110.
- [39] M. MAGIDOR, *Chang's conjecture and powers of singular cardinals*, *The Journal of Symbolic Logic*, vol. 42 (1977), pp. 272–276.
- [40] ———, *On the singular cardinals problem I*, *Israel Journal of Mathematics*, vol. 28 (1977), pp. 1–31.
- [41] ———, *On the singular cardinals problem II*, *Annals of Mathematics*, vol. 106 (1977), pp. 517–547.
- [42] ———, *Changing cofinality of cardinals*, *Fundamenta Mathematicae*, vol. 99 (1978), pp. 61–71.
- [43] A. R. D. MATHIAS, *Surrealist landscape with figures*, *Periodica Mathematica Hungarica*, vol. 10 (1979), pp. 109–175.
- [44] W. J. MITCHELL, *The core model for sequences of measures I*, *Mathematical Proceedings of the Cambridge Philosophical Society*, vol. 95 (1984), pp. 229–260.
- [45] ———, *On the singular cardinal hypothesis*, *Transactions of the American Mathematical Society*, vol. 329 (1992), pp. 507–530.
- [46] K. NAMBA, *Independence proof of  $(\omega, \omega_\alpha)$ -distributive law in complete Boolean algebras*, *Commentarii Mathematici Universitatis Sancti Pauli*, vol. 19 (1970), pp. 1–12.
- [47] K. PRIKRY, *Changing measurable into accessible cardinals*, *Dissertationes Mathematicae*, vol. 68 (1970), pp. 5–52.
- [48] L. B. RADIN, *Adding closed cofinal sequences to large cardinals*, *Annals of Mathematical Logic*, vol. 22 (1982), pp. 243–261.
- [49] F. ROWBOTTOM, *Some strong axioms of infinity incompatible with the axiom of constructibility*, *Annals of Mathematical Logic*, vol. 3 (1971), pp. 1–44.
- [50] D. SCOTT, *Measurable cardinals and constructible sets*, *Bulletin de l'Académie Polonaise des Sciences*, vol. 9 (1961), pp. 521–524.
- [51] S. SHELAH, *Jonsson algebras in successor cardinals*, *Israel Journal of Mathematics*, vol. 30 (1978), pp. 57–64, [Sh 68].
- [52] ———, *A note on cardinal exponentiation*, *The Journal of Symbolic Logic*, vol. 45 (1980), pp. 56–66, [Sh 71].
- [53] ———, *Proper forcing*, *Lecture Notes in Mathematics*, vol. 940, Springer-Verlag, Berlin, 1982.
- [54] ———, *The singular cardinals problem: independence results*, *Surveys in set theory* (A. R. D. Mathias, editor), Cambridge University Press, Cambridge, 1983, pp. 116–134, [Sh 137].

- [55] ———, *On power of singular cardinals*, *Notre Dame Journal of Formal Logic*, vol. 27 (1986), pp. 263–299, [Sh 111].
- [56] ———, *More on powers of singular cardinals*, *Israel Journal of Mathematics*, vol. 59 (1987), pp. 299–326, [Sh 256].
- [57] ———, *Successors of singulars, cofinalities of reduced products of cardinals and productivity of chain conditions*, *Israel Journal of Mathematics*, vol. 62 (1988), pp. 213–256, [Sh 282].
- [58] ———, *Products of regular cardinals and cardinal invariants of products of Boolean algebras*, *Israel Journal of Mathematics*, vol. 70 (1990), pp. 129–187, [Sh 345].
- [59] ———, *Cardinal arithmetic for skeptics*, *Bulletin of the American Mathematical Society*, vol. 26 (1992), pp. 197–210, [Sh 400a].
- [60] ———, *More on cardinal arithmetic*, *Archive for Mathematical Logic*, vol. 32 (1993), pp. 399–428, [Sh 410].
- [61] ———, *Cardinal arithmetic*, Oxford Logic Guides, Oxford University Press, Oxford, 1994.
- [62] ———, *Advances in cardinal arithmetic*, Proceedings of the Banff Conference in Alberta, to appear, [Sh 420].
- [63] ———, *Further cardinal arithmetic*, *Israel Journal of Mathematics*, to appear, [Sh 430].
- [64] ———, *The generalized continuum hypothesis revisited*, to appear, [Sh 460].
- [65] ———, *The pcf-theorem revisited*, A special volume dedicated to Paul Erdős, edited by R. Graham and J. Nešetřil, to appear, [Sh 506].
- [66] J. H. SILVER, *Some applications of model theory in set theory*, *Annals of Mathematical Logic*, vol. 3 (1971), pp. 45–110.
- [67] ———, *On the singular cardinals problem*, *Proceedings of the International Congress of Mathematicians*, 1974, pp. 265–268.
- [68] R. M. SOLOVAY, *Strongly compact cardinals and the GCH*, *Proceedings of the Tarski symposium*, American Mathematical Society, Providence, RI, 1974, pp. 365–372.
- [69] A. TARSKI, *Quelques théorèmes sur les alephs*, *Fundamenta Mathematicae*, vol. 7 (1925), pp. 1–14.

DEPARTMENT OF MATHEMATICS  
 THE PENNSYLVANIA STATE UNIVERSITY  
 UNIVERSITY PARK, PA 16802

*E-mail:* jech@math.psu.edu

## LINKED CITATIONS

- Page 1 of 3 -



You have printed the following article:

### **Singular Cardinals and the pcf Theory**

Thomas Jech

*The Bulletin of Symbolic Logic*, Vol. 1, No. 4. (Dec., 1995), pp. 408-424.

Stable URL:

<http://links.jstor.org/sici?sici=1079-8986%28199512%291%3A4%3C408%3ASCATPT%3E2.0.CO%3B2-5>

---

*This article references the following linked citations. If you are trying to access articles from an off-campus location, you may be required to first logon via your library web site to access JSTOR. Please visit your library's website or contact a librarian to learn about options for remote access to JSTOR.*

## References

### <sup>2</sup> **Singular Cardinals and the Generalized Continuum Hypothesis**

J. E. Baumgartner; K. Prikry

*The American Mathematical Monthly*, Vol. 84, No. 2. (Feb., 1977), pp. 108-113.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9890%28197702%2984%3A2%3C108%3ASCATGC%3E2.0.CO%3B2-H>

### <sup>7</sup> **The Independence of the Continuum Hypothesis**

Paul J. Cohen

*Proceedings of the National Academy of Sciences of the United States of America*, Vol. 50, No. 6. (Dec. 15, 1963), pp. 1143-1148.

Stable URL:

<http://links.jstor.org/sici?sici=0027-8424%2819631215%2950%3A6%3C1143%3ATIOTCH%3E2.0.CO%3B2-5>

### <sup>8</sup> **The Independence of the Continuum Hypothesis, II**

Paul J. Cohen

*Proceedings of the National Academy of Sciences of the United States of America*, Vol. 51, No. 1. (Jan. 15, 1964), pp. 105-110.

Stable URL:

<http://links.jstor.org/sici?sici=0027-8424%2819640115%2951%3A1%3C105%3ATIOTCH%3E2.0.CO%3B2-U>

**NOTE:** *The reference numbering from the original has been maintained in this citation list.*

## LINKED CITATIONS

- Page 2 of 3 -



<sup>9</sup> **A Model in Which GCH Holds at Successors but Fails at Limits**

James Cummings

*Transactions of the American Mathematical Society*, Vol. 329, No. 1. (Jan., 1992), pp. 1-39.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9947%28199201%29329%3A1%3C1%3AAMIWGH%3E2.0.CO%3B2-J>

<sup>14</sup> **The Generalized Continuum Hypothesis Can Fail Everywhere**

Matthew Foreman; W. Hugh Woodin

*The Annals of Mathematics*, 2nd Ser., Vol. 133, No. 1. (Jan., 1991), pp. 1-35.

Stable URL:

<http://links.jstor.org/sici?sici=0003-486X%28199101%292%3A133%3A1%3C1%3ATGCHCF%3E2.0.CO%3B2-D>

<sup>15</sup> **Inequalities for Cardinal Powers**

Fred Galvin; Andras Hajnal

*The Annals of Mathematics*, 2nd Ser., Vol. 101, No. 3. (May, 1975), pp. 491-498.

Stable URL:

<http://links.jstor.org/sici?sici=0003-486X%28197505%292%3A101%3A3%3C491%3AIFCP%3E2.0.CO%3B2-%23>

<sup>22</sup> **Extender Based Forcings**

Moti Gitik; Menachem Magidor

*The Journal of Symbolic Logic*, Vol. 59, No. 2. (Jun., 1994), pp. 445-460.

Stable URL:

<http://links.jstor.org/sici?sici=0022-4812%28199406%2959%3A2%3C445%3AEBF%3E2.0.CO%3B2-P>

<sup>32</sup> **On a Conjecture of Tarski on Products of Cardinals**

Thomas Jech; Saharon Shelah

*Proceedings of the American Mathematical Society*, Vol. 112, No. 4. (Aug., 1991), pp. 1117-1124.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9939%28199108%29112%3A4%3C1117%3AOCOTO%3E2.0.CO%3B2-X>

<sup>35</sup> **Inner Models and Large Cardinals**

Ronald Jensen

*The Bulletin of Symbolic Logic*, Vol. 1, No. 4. (Dec., 1995), pp. 393-407.

Stable URL:

<http://links.jstor.org/sici?sici=1079-8986%28199512%291%3A4%3C393%3AIMALC%3E2.0.CO%3B2-V>

## LINKED CITATIONS

- Page 3 of 3 -



<sup>39</sup> **Chang's Conjecture and Powers of Singular Cardinals**

Menachem Magidor

*The Journal of Symbolic Logic*, Vol. 42, No. 2. (Jun., 1977), pp. 272-276.

Stable URL:

<http://links.jstor.org/sici?sici=0022-4812%28197706%2942%3A2%3C272%3ACCAPOS%3E2.0.CO%3B2-A>

<sup>41</sup> **On the Singular Cardinals Problem II**

Menachem Magidor

*The Annals of Mathematics*, 2nd Ser., Vol. 106, No. 3. (Nov., 1977), pp. 517-547.

Stable URL:

<http://links.jstor.org/sici?sici=0003-486X%28197711%292%3A106%3A3%3C517%3AOTSCPI%3E2.0.CO%3B2-H>

<sup>45</sup> **On the Singular Cardinal Hypothesis**

W. J. Mitchell

*Transactions of the American Mathematical Society*, Vol. 329, No. 2. (Feb., 1992), pp. 507-530.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9947%28199202%29329%3A2%3C507%3AOTSCH%3E2.0.CO%3B2-T>

<sup>52</sup> **A Note on Cardinal Exponentiation**

Saharon Shelah

*The Journal of Symbolic Logic*, Vol. 45, No. 1. (Mar., 1980), pp. 56-66.

Stable URL:

<http://links.jstor.org/sici?sici=0022-4812%28198003%2945%3A1%3C56%3AANOCE%3E2.0.CO%3B2-P>