

INTERFACE AND MIXED BOUNDARY VALUE PROBLEMS ON n -DIMENSIONAL POLYHEDRAL DOMAINS

CONSTANTIN BĂCUȚĂ, ANNA L. MAZZUCATO*, VICTOR NISTOR,
AND LUDMIL T. ZIKATANOV

ABSTRACT. We prove a well-posedness result for mixed boundary value /interface problems of second-order, positive, strongly elliptic operators in weighted Sobolev spaces on bounded, non-convex, curvilinear polyhedral domain Ω in a manifold M . The typical weight we consider is the distance to the set of singular boundary points. Our model problem is $-\operatorname{div}(a\nabla u) = f$, in Ω , $u = 0$ on $\partial_D\Omega$, and $\partial_\nu u = 0$ on $\partial_N\Omega$, where $a > 0$ is constant on some polyhedral subdomains Ω_j defining a decomposition $\bar{\Omega} = \cup_j \bar{\Omega}_j$, and $\partial\Omega = \partial_D\Omega \cup \partial_N\Omega$ is a decomposition of the boundary of Ω into polyhedral subsets corresponding, respectively, to Dirichlet and Neumann boundary conditions. An important step in our proof is a *regularity* result, which holds for general strongly elliptic operators that are not necessarily positive. The well-posedness result applies to positive operators, provided the interfaces are smooth and there are no adjacent faces with Neumann boundary conditions.

CONTENTS

Introduction	1
1. The problem and statement of the main results	4
2. Polyhedral domains	10
3. Lie manifolds with boundary	16
4. Desingularization of polyhedra	20
5. Weighted Sobolev spaces	34
6. Proofs	39
References	45

INTRODUCTION

Let $\Omega \subset \mathbb{R}^n$ be an open, bounded set. Consider the boundary value problem

$$(1) \quad \begin{cases} \Delta u = f \\ u|_{\partial\Omega} = g, \end{cases}$$

Date: August 22, 2008.

2000 Mathematics Subject Classification. 35J25, 58J32, 52B70, 51B25.

Key words and phrases. Elliptic equations, polyhedral domains, mixed boundary conditions, interface, weighted spaces, well-posedness, Lie manifold.

* Corresponding author.

The authors were supported in part by NSF grants.

where Δ is the Laplace operator. For Ω smooth, this boundary value problem has a unique solution $u \in H^{s+2}(\Omega)$ depending continuously on $f \in H^s(\Omega)$ and $g \in H^{s+3/2}(\partial\Omega)$, $s > 1/2$. See the books of Evans [16], Lions and Magenes [36], or Taylor [54] for proofs of this basic and well known result.

It is also well known that this result does not extend to non-smooth domains Ω . For instance, Jerison and Kenig prove in [24] that if $g = 0$ and $\Omega \subset \mathbb{R}^3$ is an open, bounded set such that $\partial\Omega$ is Lipschitz, then Equation (1) has a unique solution in $W^{s,p}(\Omega)$ depending continuously on $f \in W^{s-2,p}(\Omega)$ if, and only if, $(1/p, s)$ belongs to a certain explicit hexagon, the Jerison-Kenig hexagon. They also prove a similar result if $\Omega \subset \mathbb{R}^2$. A consequence of this result, is that the smoothness of the solution u (measured by the order s of the Sobolev space $W^{s,p}(\Omega)$ containing it) will not exceed, in general, a certain bound that depends on the domain Ω and p , even if f is smooth.

In addition to the Jerison and Kenig paper mentioned above, a deep analysis of the difficulties that arise for $\partial\Omega$ Lipschitz is contained in the papers of Babuška [4], Băcuță, Bramble, and Xu [7], Babuška and Guo [21], Jerison and Kenig [22, 23], Kenig [27], Kenig and Toro [28], Mitrea and Taylor [43, 45, 46], Verchota [55], and others (see the references in the aforementioned papers). Results more specific to curvilinear polyhedral domains are contained in the papers of Costabel [10], Dauge [11, 12], Elschner [13, 14], Kondratiev [30, 31], Mazya and Rossmann [39], Rossmann [48] and others. Excellent references are also the monographs of Grisvard [18, 19] as well as the recent books [32, 33, 37, 38, 47].

In this paper, we consider the boundary value problem (1) when Ω is a *bounded curvilinear polyhedral domain* in \mathbb{R}^n , or, more generally, in a manifold M of dimension n and Poisson's equation $\Delta u = f$ is replaced by a positive, strongly elliptic scalar equation. We define curvilinear polyhedral domains inductively in Section 2. We allow polyhedral domains to be disconnected for technical reasons, that is, for the purpose of defining them inductively. Our results, however, are formulated for connected polyhedral domains. Many polyhedral domains are Lipschitz domains, but not all. This fact is discussed in detail by Vogel and Verchota in [56], where they also prove that the harmonic measure is absolutely continuous with respect to the Lebesgue measure on the boundary as well as the solvability of Equation (1) if $f = 0$ and $g \in L^{2-\epsilon}(\partial\Omega)$, thus generalizing several earlier, classical results.

Instead of working with the usual Sobolev spaces, as in the papers of Jerison and Kenig [24], Mitrea and Taylor [45], or Verchota and Vogel [56], we shall work in some weighted analogues of these papers. Let $\Omega^{(n-2)} \subset \partial\Omega$ be the set of singular (or non-smooth) boundary points of Ω , that is, the set of points $p \in \partial\Omega$ such $\partial\Omega$ is not smooth in a neighborhood of p . We shall denote by $\eta_{n-2}(x)$ the distance from a point $x \in \Omega$ to the set $\Omega^{(n-2)}$. We agree to take $\eta_{n-2} = 1$ if there are no such points, that is, if $\partial\Omega$ is smooth. We then consider the weighted Sobolev spaces

$$(2) \quad \mathcal{K}_a^\mu(\Omega) = \{u \in L_{\text{loc}}^2(\Omega), \eta_{n-2}^{|\alpha|-a} \partial^\alpha u \in L^2(\Omega), \text{ for all } |\alpha| \leq \mu\}, \quad \mu \in \mathbb{Z}_+,$$

which we endow with the induced Hilbert space norm. A similar definition, Definition 47, yields the weighted Sobolev spaces $\mathcal{K}_a^s(\partial\Omega)$, $s \in \mathbb{R}_+$. By including an extra weight h in the spaces $\mathcal{K}_a^\mu(\Omega)$ we obtain the spaces $h\mathcal{K}_a^\mu(\Omega)$ (h is required to be an admissible weight, see subsection 5.1). In this article, we denote the spaces $h\mathcal{K}_a^\mu(\Omega)$ as the *Babuška-Kondratiev spaces*, although several other authors contributed to their theory, including the essential work of Mazya and Plamenevskii. These spaces

are closely related to the weighted Sobolev spaces on non-compact manifolds considered, for example, in the works of Erkip and Schrohe [15], Grubb [20], Schrohe [49], as well as the sequence of papers of Schrohe and Schulze [50, 51] concerning related results on boundary value problems on non-compact manifolds and, more generally, the analysis on non-compact manifolds.

The main result of this article, Theorem 1.2 applies to operators with *piecewise smooth coefficients*, such as $\operatorname{div} a \nabla u = f$, where a is allowed to have only jumps. A simplified version of that result, when formulated for the Laplace operator Δ on \mathbb{R}^n with Dirichlet boundary conditions, reads as follows. In this theorem and throughout this paper, Ω will always denote an open set.

Theorem 0.1. *Let $\Omega \subset \mathbb{R}^n$ be a bounded, curvilinear polyhedral domain and $\mu \in \mathbb{Z}_+$. Then there exists $\eta > 0$ such that the boundary value problem (1) has a unique solution $u \in \mathcal{K}_{a+1}^{\mu+1}(\Omega)$ for any $f \in \mathcal{K}_{a-1}^{\mu-1}(\Omega)$, any $g \in \mathcal{K}_{a+1/2}^{\mu+1/2}(\partial\Omega)$, and any $|a| < \eta$. This solution depends continuously on f and g . If $a = \mu = 0$, this solution is the solution of the associated variational problem.*

The case $n = 2$ of the above theorem is Theorem 6.6.1 in the excellent monograph [32]. Results in higher dimensions related to the ones in our paper can be found, for instance, in [11, 32, 39, 47]. These works also use the framework of the $\mathcal{K}_a^\mu(\Omega)$ spaces. The Babuška–Kondratiev spaces $h\mathcal{K}_a^\mu(\Omega)$, with h an admissible weight (see Subsection 5.1 for the definition of admissible weights) are somewhat more general. We also take the dimension n of the ambient Euclidean space $\mathbb{R}^n \supset \Omega$ to be arbitrary. Furthermore, we impose mixed Dirichlet/Neumann boundary conditions and allow the boundary conditions to change along $n - 2$ -dimensional, piece-wise smooth hypersurfaces in each hyperface of Ω . To handle this situation, we include all points where the boundary conditions change in the singular set of Ω , giving rise to a polyhedral structure on Ω which is not entirely determined by geometry, but also takes into account the specifics of the boundary value problem. However, we consider only second order, strongly elliptic systems. This restriction allows us in particular to use coercive estimates. The method of layer potentials seems to give more precise results, but is less elementary (see e.g. [27, 44, 45]). We use manifolds in order to be able to prove estimates inductively.

The paper is organized as follows. In Section 1, we introduce the mixed boundary value/interface problem of Equation (6), and state the main results of the paper, Theorem 1.1 on the regularity of (6) in weighted spaces of arbitrarily high Sobolev index, and Theorem 1.2 on the solvability of (6) under additional conditions on the operator (positivity) and on the domain (smooth interfaces and no Neumann-Neumann hyperfaces). In Section 2, we give a notion of curvilinear, polyhedral domain in arbitrary dimension using induction, then we specialize to the familiar case of polygonal domains in \mathbb{R}^2 and polyhedral domains in \mathbb{R}^3 , and describe the desingularization $\Sigma(\Omega)$ of the domain Ω in this familiar setting. Before discussing the desingularization in higher dimension, we recall briefly needed notions from the theory of Lie manifolds with boundary in Section 3. Then, in Section 4 we show that $\Sigma(\Omega)$, also defined by induction on the dimension, naturally carries a structure of Lie manifold with boundary. We also discuss the construction of the canonical weight function r_Ω , which extends smoothly to $\Sigma(\Omega)$ and is comparable to the distance to the singular set. In turn, this Lie structure allows to identify the Babuška-Kondratiev spaces $\mathcal{K}_a^\mu(\Omega)$, $\mu \in \mathbb{Z}_+$, with standard Sobolev spaces on

$\Sigma(\Omega)$ with respect to a certain measure, and hence define the Babuška-Kondratiev spaces on the boundary $\mathcal{K}_a^s(\partial\Omega)$, $s \in \mathbb{R}$. Lastly Section 6 contains the proofs of the main results and most lemmas of the paper; in particular, it contains a proof of the weighted Hardy-Poincaré inequality used to establish positivity or strict coercivity for the problem of Equation (6).

We conclude this Introduction with a word on notation. By Ω we always mean an open set in \mathbb{R}^n . By a diffeomorphism, we mean a C^∞ invertible map with C^∞ inverse. C denotes a generic constant that may change from line to line. We also denote $\mathbb{Z}_+ = \{0, 1, 2, 3, \dots\}$.

Acknowledgements. . We thank Bernd Ammann, Ivo Babuška, Wolfgang Dahmen, Alexandru Ionescu, and Daniel Tătaru for useful discussions. We also thank Johnny Guzman for pointing the reference [31] to us. The second named author would like to thank Institute Henri Poincaré in Paris for its hospitality while parts of this work were being completed.

1. THE PROBLEM AND STATEMENT OF THE MAIN RESULTS

We begin by introducing the class of differential operators and the associated mixed Dirichlet-Neumann boundary value/interface problem that will be the object of study. For simplicity, we consider primarily the scalar case, although our results extend to systems as well. Then, we state the main results of this article, namely the well-posedness, Theorem 1.2, and regularity, Theorem 1.1, of the mixed boundary value/interface problem (see Equation (6)) in weighted Sobolev spaces for n -dimensional, curvilinear polyhedral domains $\Omega \subset \mathbb{R}^n$. Our analysis is general enough to extend to a bounded subdomain $\Omega \subset M$ of a compact Riemannian manifold M . Initially the reader may assume the polyhedron is straight, that is, informally every j -dimensional component of the boundary, $j = 1, \dots, n-1$ is a subset of an affine space. A complete definition of a curvilinear polyhedral domain is given in Section 2.

1.1. The differential operator P and the associated problem. We let $\Omega \subset \mathbb{R}^n$ be a bounded, curvilinear stratified polyhedral domain (see Definition 2.1). Ω need not be connected, nor convex. We assume that we are given a decomposition

$$(3) \quad \bar{\Omega} = \cup_{j=1}^N \bar{\Omega}_j,$$

into disjoint polyhedral subdomains. In particular, every face of Ω is also a face of one of the Ω_j . As discussed in Section 4, a face of codimension 1 is called a hyperface. For well-posedness results, we shall assume that

$$(4) \quad \Gamma = \cup_{j=1}^N \partial\Omega_j \setminus \partial\Omega,$$

is a finite collection of disjoint, smooth $(n-1)$ -hypersurfaces. We observe that, since each Ω_j is a polyhedron, each component of Γ intersects $\partial\Omega$ transversely. We refer to Γ as the *interface*.

We also assume that the boundary of Ω is partition into two disjoint polyhedral subsets

$$(5) \quad \partial\Omega = \partial_D\Omega \cup \partial_N\Omega,$$

with $\partial_N\Omega$ open in the topology of $\partial\Omega$. Each of $\partial_D\Omega$ and $\partial_N\Omega$ consists of a union of faces of Ω . For well-posedness results, we shall assume that $\partial_N\Omega$ does not contain adjacent faces of $\partial\Omega$.

We are interested in studying the following mixed boundary value/interface problem for a certain class of elliptic, scalar operators P described below:

$$(6) \quad \begin{cases} Pu = f & \text{on } \Omega, \\ u|_{\partial\Omega} = g_D & \text{on } \partial_D\Omega, \\ D_\nu^P u|_{\partial_N\Omega} = g_N & \text{on } \partial_N\Omega, \\ u^+ = u^-, D_\nu^{P^+} u = D_\nu^{P^-} u & \text{on } \Gamma, \end{cases}$$

Above, ν is the unit outer normal to $\partial\Omega$, which is defined almost everywhere, D_ν^P is the conormal derivative associated to the operator P (see (10)), and \pm refers to one-sided, non-tangential approaches to the interface Γ . The coefficients of P will have in general jump discontinuities along Γ .

We next introduce the class of differential operators that we consider. At first, the reader may assume $P = -\Delta$, the Laplace operator. We shall write $Re(z) := \frac{1}{2}(z + \bar{z})$, or simply $Re z$ for the real part of a complex number z .

Let $u \in H_{\text{loc}}^2(\Omega)$. We shall study the following scalar, differential operator in divergence form

$$(7) \quad Pu(x) = - \sum_{j,k=1}^n \partial_j [A_{jk}(x) \partial_k u(x)] + \sum_{j=1}^n B_j(x) \partial_j u(x) + C(x)u(x).$$

The coefficients A_{jk}, B_j, C are real valued with only jump discontinuities on the interface Γ , the operator P is required to be uniformly strongly elliptic and to satisfy another positivity condition. More precisely, the coefficients of P must satisfy:

$$(8a) \quad A_{jk}, B_j, C \in \oplus_{j=1}^N \mathcal{C}^\infty(\bar{\Omega}_j) \cap L^\infty(\bar{\Omega})$$

$$(8b) \quad Re \left(\sum_{j,k=1}^n [A_{jk}(x)] \xi_j \bar{\xi}_k \right) \geq \epsilon \sum_{j=1}^n |\xi_j|^2, \quad \xi_j \in \mathbb{C}, \text{ and}$$

$$(8c) \quad 2C(x) - \sum_{j=1}^n \partial_j B_j(x) \geq 0.$$

Our results extend to systems satisfying the *strong Legendre–Hadamard* condition, namely

$$(9) \quad Re \left(\sum_{j,k=1}^n \sum_{p,q=1}^m [A_{jk}(x)]_{pq} \xi_{jp} \bar{\xi}_{kq} \right) \geq \epsilon \sum_{j=1}^n \sum_{p=1}^m |\xi_{jp}|^2, \quad \xi_{jp} \in \mathbb{C},$$

and a condition on the lower-order terms equivalent to (8c). This condition is not satisfied however by the system of anisotropic elasticity in \mathbb{R}^3 , for which nevertheless the well-posedness result holds if the elasticity tensor is positive definite on symmetric matrices (see [40]).

In (8a), the “regularity condition on the coefficients of P ” means that the coefficients and their derivatives of all orders have well-defined limits on each non-tangential approach to Γ , but as equivalence classes in L^∞ they may have jump discontinuities along the interface. This condition can be relaxed, but it allows us to state a regularity result of arbitrary order in each subdomains for the solution to the problem (6). The conormal derivative associated to the operator P is formally

defined by:

$$(10) \quad D_P^\nu u(x) = \sum_{i,j=1}^n \nu_i A_{ij} \partial_j u(x),$$

where ν is the unit outer co-normal vector to the boundary of Ω . We give meaning to (10) in the sense of trace at the boundary. In particular, for u regular enough $D_P^\nu u$ is defined almost everywhere on the boundary as a non-tangential limit.

The problem (6) with $g_D = 0$ is interpreted in a weak (or variational) sense, using the bilinear form $B(u, v)$ defined by:

$$(11) \quad B(u, v) := \sum_{j,k=1}^n (A_{jk} \partial_k u, \partial_j v) + \sum_{j=1}^n (B_j \partial_j u, v) + (Cu, v),$$

which is well-defined for any $u, v \in H^1(\Omega)$. Then, (6) is weakly equivalent to

$$(12) \quad B(u, v) = (f, v)_{L^2(\Omega)} + (g_N, v)_{\partial_N \Omega},$$

where the second paranthesis denotes the pairing between a distribution and a (suitable) function. The jump or *transmission* conditions, $u^+ = u^-$, $D_P^\nu u^+ = D_P^\nu u^-$ at the interface Γ follow from the weak formulation and the H^1 -regularity of weak solutions, and hence justify passing from the strong formulation (6) to the weak one (12). Otherwise, in general, the difference $D_P^{\nu^+} u - D_P^{\nu^-} u$ may be non-zero and may be included as a distributional term in f .

Condition (8c) implies the Poincaré type inequality $Re B(u, u) > C(\eta_{n-2} u, \eta_{n-2} u)_{L^2}$, if $\partial_D \Omega \neq \emptyset$. In fact, it is enough to assume that the latter is satisfied instead of (8c). For applications, however, it is more convenient to have the concrete condition (8c).

Problems of the form (6) arise in many applications. An important example is given by (linear) elastostatics. In this case, $[Pu]^i = -\sum_{jkl=1}^3 \partial_j C^{ijkl} \partial_k u^l$, $i = 1, 2, 3$, where C is the fourth-order elasticity tensor, modelling the response of an elastic body under small deformations. Dirichlet or displacement boundary conditions correspond to clamping (parts of) the boundary, while Neumann or traction boundary conditions correspond to loading mechanically (parts of) the boundary. Interfaces arise due to the use of different materials. A careful analysis of mixed Dirichlet/Neumann boundary value problems for linear elastostatics in 3-dimensional curvilinear, polyhedral domains, was carried out by two of the authors in [40]. There, the concept of a “domain with polyhedral structure” is more general than in this paper and includes cracks. In [35], they studied mixed boundary value/interface problems and the implementation of the Finite Element Method on “domains with polygonal structure” with nonsmooth interfaces (see also [8]). The results of this paper can be extended to include domains with cracks, as in [40] and [35], but the topological machinery used there, including the notion of an “unfolded boundary” [11] in arbitrary dimensions is significantly more complex.

1.1.1. *Operators on manifolds.* We turn to consider the assumptions on P when the domain Ω is a curvilinear, polyhedral domain in a manifold M of the same dimension. Let then E be a vector bundles on M endowed with a hermitian metric. A coordinate free expression of the conditions in Equations (8a)–(8c) is obtained as follows. We assume that there exist a metric preserving connection $\nabla : \Gamma(E) \rightarrow$

$\Gamma(E \otimes T^*M)$, a smooth endomorphism $A \in \text{End}(E \otimes T^*M)$, and a first order differential operator $P_2 : \Gamma(E) \rightarrow \Gamma(E)$ with smooth coefficients such that

$$(13) \quad A + A^* \geq 2\epsilon I \text{ for some } \epsilon > 0.$$

Then we define $P_1 = \nabla^* A \nabla$ and $P = P_1 + P_2$. In particular, the operator P will satisfy the strong Lagrange–Hadamard condition in a neighborhood of Ω in M . Note that if $\Omega \subset \mathbb{R}^n$ and the vector bundle E is trivial, then the condition of (13) reduce to the conditions of (8), by taking ∇ to be the trivial connection. We can allow A to have jump discontinuities as well.

1.2. The main results. We are ready to state the principal results of this paper. *We continue to assume hypotheses (3)–(5) on the domain Ω and its decomposition into disjoint subdomains Ω_j separated by the interface Γ .*

We begin with a regularity results for solutions to the problem (6) in weighted Sobolev spaces $h\mathcal{K}_a^\mu$, $\mu \in \mathbb{Z}_+$, $a \in \mathbb{R}$, where

$$\mathcal{K}_a^\mu(\Omega) := \{u \in L^2_{\text{loc}}(\Omega), \eta_{n-2}^{|\alpha|-a} \partial^\alpha u \in L^2(\Omega), \text{ for all } |\alpha| \leq \mu\}, \quad \mu \in \mathbb{Z}_+,$$

and

$$h\mathcal{K}_a^\mu(\Omega) := \{hu, u \in \mathcal{K}_a^\mu(\Omega)\}.$$

(See Section 5 for a detailed discussion and main properties of these spaces.) Above, η_{n-2} is the distance to the singular set in Ω given in Definition 2.5, while h is a so-called admissible weight described in Section 5.1. Initially, the reader may assume that $h = r_\Omega^b$, $b \in \mathbb{R}$, where r_Ω is a function comparable to the distance function η_{n-2} close to the singular set, but with better regularity than η_{n-2} away from the singular set. (We refer again to Subsection 5.1 for more details). The weight h is important in applications of the theory developed here for numerical methods, where appropriate choices of h yield quasi-optimal rates of convergence for the Finite Element approximation to the weak solution of the problem (6) (see [6, 35]).

Theorem 1.1. *Let $\Omega \subset \mathbb{R}^n$ be a bounded, curvilinear polyhedral domain of dimension n . Assume that the operator P satisfies conditions (8a) and (8b). Let $\mu \in \mathbb{Z}_+$, $a \in \mathbb{R}$, and $u \in h\mathcal{K}_{a+1}^1(\Omega)$ be such that $Pu \in h\mathcal{K}_{a-1}^{\mu-1}(\Omega_j)$, for all j , $u|_{\partial_D \Omega} \in h\mathcal{K}_{a+1/2}^{\mu+1/2}(\partial_D \Omega)$, $D_\nu^P u|_{\partial_N \Omega} \in h\mathcal{K}_{a-1/2}^{\mu-1/2}(\partial_N \Omega)$. If h is an admissible weight, then $u \in h\mathcal{K}_{a+1}^{\mu+1}(\Omega_j)$, for all $j = 1, \dots, N$, and*

$$(14) \quad \|u\|_{h\mathcal{K}_{a+1}^{\mu+1}(\Omega_j)} \leq C \left(\sum_{k=1}^N \|Pu\|_{h\mathcal{K}_{a-1}^{\mu-1}(\Omega_k)} + \|u\|_{h\mathcal{K}_{a+1}^0(\Omega)} + \|u|_{\partial_D \Omega}\|_{h\mathcal{K}_{a+1/2}^{\mu+1/2}(\partial_D \Omega)} + \|u|_{\partial_N \Omega}\|_{h\mathcal{K}_{a-1/2}^{\mu-1/2}(\partial_N \Omega)} \right)$$

for a constant $C = C(\Omega, P, \mu, a, h) > 0$, independent of u .

The proof of the regularity theorem exploits Lie manifolds and their structure, and can be found in Section 6. Note that in this theorem we do not require the interfaces to be smooth and we allow for adjacent faces with Neumann boundary conditions.

Under additional conditions on the set Ω and its boundary ensuring strict co-ercivity of the bilinear form B of equation (11), we obtain a well-posedness result for problem (6). In [35], two of the authors obtained a well-posedness result in an augmented space on polygonal domains with “Neumann-Neumann vertices,” *i.e.*,

vertices for which both sides joining at the vertex are given Neumann boundary conditions, and for which the interface Γ is not smooth. Such result is based on specific spectral properties of operator pencils near the vertices and is not easily extendable to higher dimension.

Theorem 1.2. *Let $\Omega \subset \mathbb{R}^n$ be a bounded, connected curvilinear polyhedral domain of dimension n . Assume that $\partial\Omega_N$ does not contain any two adjacent hyperfaces, that $\partial_D\Omega$ is not empty, and that the interface Γ is smooth. In addition, assume that the operator P satisfies conditions (8). Let $\mathcal{W}_\mu(\Omega)$, $\mu \in \mathbb{Z}_+$, be the set of admissible weights h such that the map $\tilde{P}(u) := (Pu, u|_{\partial_D\Omega}, D_\nu^P u|_{\partial_N\Omega})$ establishes an isomorphism*

$$\begin{aligned} \tilde{P} : \{u \in \bigoplus_{j=1}^N h\mathcal{K}_1^{\mu+1}(\Omega_j) \cap h\mathcal{K}_1^1(\Omega), u^+ = u^-, D_\nu^P u^+ = D_\nu^P u^- \text{ on } \Gamma\} \\ \rightarrow \bigoplus_{j=1}^N h\mathcal{K}_{-1}^{\mu-1}(\Omega_j) \oplus h\mathcal{K}_{1/2}^{\mu+1/2}(\partial_D\Omega) \oplus h\mathcal{K}_{-1/2}^{\mu-1/2}(\partial_N\Omega). \end{aligned}$$

Then the set $\mathcal{W}_\mu(\Omega)$ is an open set containing 1.

Theorem 1.2 reduces to a well-known, classical result when Ω is a smooth bounded domain. (See however Remark 6.11 for a result on smooth domains that is not classical.) The same is true for the following result, Theorem 1.3, which works for general domains on manifolds. Note however that for manifolds it is more difficult to express the coercive property, so for more complete results we restrict to the case of operators of Laplace type.

Theorem 1.3. *Let $\Omega \subset M$ be a bounded, connected curvilinear polyhedral domain of dimension n . Assume that every connected component of Ω has a non-empty boundary and that the operator P satisfies condition (13). Assume additionally that no two hyperfaces of $\partial\Omega$ are endowed with Neumann boundary conditions and that the interface Γ is smooth. Let $c \in \mathbb{C}$ and $\mathcal{W}'_\mu(\Omega)$ be the set of admissible weights h such that the map $\tilde{P}_c(u) := (Pu + cu, u|_{\partial\Omega}, \partial_\nu u|_{\partial\Omega})$ establishes an isomorphism*

$$\begin{aligned} \tilde{P}_c : \{u \in \bigoplus_{j=1}^N h\mathcal{K}_1^{\mu+1}(\Omega_j) \cap h\mathcal{K}_1^1(\Omega), u^+ = u^-, D_\nu^P u^+ = D_\nu^P u^- \text{ on } \Gamma\} \\ \rightarrow \bigoplus_{j=1}^N h\mathcal{K}_{-1}^{\mu-1}(\Omega_j) \oplus h\mathcal{K}_{1/2}^{\mu+1/2}(\partial_D\Omega) \oplus h\mathcal{K}_{-1/2}^{\mu-1/2}(\partial_N\Omega). \end{aligned}$$

Then the set $\mathcal{W}'_\mu(\Omega)$ is an open set. This set contains 1 if the real part of c is large or if P is the Laplace operator associated to some smooth metric on M and $c = 0$.

For the rest of this section, Ω and P will be as in Theorem 1.2. We discuss some immediate consequences of Theorem 1.2. Analogous results can be obtained from Theorem 1.3, but we will not state them explicitly. The continuity of the inverse of \tilde{P} is made explicit in the following corollary.

Corollary 1.4. *There exists a constant $C = C(\Omega, P, \mu, a, h) > 0$, independent of f , g_D , and g_N , such that*

$$\begin{aligned} \|u\|_{h\mathcal{K}_1^1(\Omega)} + \|u\|_{h\mathcal{K}_1^{\mu+1}(\Omega_j)} &\leq C \left(\sum_{j=1}^N \|Pu\|_{h\mathcal{K}_1^{\mu-1}(\Omega_j)} \right. \\ &\quad \left. + \|u|_{\partial_D\Omega}\|_{h\mathcal{K}_1^{\mu+1/2}(\partial_D\Omega)} + \|D_\nu^P u|_{\partial_N\Omega}\|_{h\mathcal{K}_1^{\mu+1/2}(\partial_N\Omega)} \right), \end{aligned}$$

for any $u \in h\mathcal{K}_1^1(\Omega)$ and any j .

From the fact that η_{n-2} is equivalent to r_Ω by Proposition 4.9 and Corollary 4.11, we obtain the following corollary.

Corollary 1.5. *There exists $\eta > 0$ such that*

$$\begin{aligned} (P, D_\nu^P) : \{u \in \bigoplus_{j=1}^N \mathcal{K}_{a+1}^{\mu+1}(\Omega_j) \cap \mathcal{K}_1^1(\Omega), u|_{\partial_D\Omega} = 0, \\ u^+ = u^-, D_\nu^P u^+ = D_\nu^P u^- \text{ on } \Gamma\} \rightarrow \bigoplus_{j=1}^N h\mathcal{K}_{a-1}^{\mu-1}(\Omega_j) \oplus h\mathcal{K}_{a-1/2}^{\mu-1/2}(\partial_N\Omega) \end{aligned}$$

is an isomorphism for all $\mu \in \mathbb{Z}_+$ and all $|a| < \eta$.

Proof. From the results in Sections 5 and 5.1, $\mathcal{K}_{a+1}^{\mu+1} = r_\Omega^a \mathcal{K}_1^{\mu+1}$ and r_Ω^a is an admissible weight for any $a \in \mathbb{R}$. The result then follows from the fact that $\mathcal{W}_\mu(\Omega)$ is an open set containing the weight 1 by Theorem 1.2. \square

The following corollary gives a characterization of the set $\mathcal{W}(\Omega)$ in the spirit of [8]. There, similar arguments give that for $n = 2$ the constant η in the previous corollary is $\eta = \pi/\alpha_M$, where α_M is the largest angle of Ω . See also [31].

Corollary 1.6. *Let $h = r_\Omega^a$ be an admissible weight such that either $a \geq 0$ or $a \leq 0$. Assume that for all $\lambda \in [0, 1]$ the map*

$$(P, D_\nu^P) : \{u \in h^\lambda \mathcal{K}_1^1(\Omega), u|_{\partial_D\Omega} = 0, D_\nu^P u|_{\partial_N\Omega} = 0\} \rightarrow h^\lambda \mathcal{K}_1^{-1}(\Omega)$$

is Fredholm. Then $h \in \mathcal{W}_\mu(\Omega)$.

The corollary holds for more general weights $h = \prod_H x_H^{a_H}$, where x_H is the distance to an hyperface H at infinity (see Section 5.1), as long as all $a_H \geq 0$ or all $a_H \leq 0$.

Proof. We proceed as in [8]. The family $P_\lambda := h^{-\lambda} P h^\lambda$ is continuous for $\lambda \in [0, 1]$, consists of Fredholm operators by hypothesis, and is invertible for $\lambda = 0$ by Theorem 1.2. It follows that the family P_λ consists of Fredholm operators of index zero. To prove that these operators are isomorphisms, it is hence enough to prove that they are either injective or surjective. Assume first that $a \geq 0$ in the definition of h . Then $h^\lambda \mathcal{K}_1^1(\Omega) \subset \mathcal{K}_1^1(\Omega)$. Therefore P is injective on $h^\lambda \mathcal{K}_1^1(\Omega) \cap \{u|_{\partial_D\Omega} = 0, D_\nu^P u|_{\partial_N\Omega} = 0\}$. This, in turn, gives that P_λ is injective.

Assume that $a \leq 0$ and consider

$$(15) \quad P_\lambda : h^\lambda \mathcal{K}_1^1(\Omega) \cap \{u|_{\partial\Omega} = 0, D_\nu^P u|_{\partial_N\Omega} = 0\} \rightarrow h^\lambda \mathcal{K}_1^{-1}(\Omega).$$

We have $(P_\lambda)^* = (P^*)_{-\lambda}$. The same argument as above shows that P_λ^* is injective, and hence that it is an isomorphism, for all $0 \leq \lambda \leq 1$. Hence P_λ is an isomorphism for all $0 \leq \lambda \leq 1$. \square

2. POLYHEDRAL DOMAINS

In this section we introduce the class of domains to which the results of the previous sections apply. We then specialize to domains in 2 and 3 dimensions and provide ample examples. The reader may at first concentrate on this case. We describe how to desingularize the domain in arbitrary dimension later in the paper, using the theory of Lie manifold introduced in the next section.

Let Ω be a proper open set in \mathbb{R}^n or more generally in a smooth manifold M of dimension n . Our main focus is the analysis of partial differential equations on Ω , specifically the mixed boundary value/interface problem (6). For this reason, we give Ω a structure that is not entirely determined by geometry, rather it takes into account the boundary and interface conditions for the operator P in problem (6).

We assume that Ω is given a smooth stratification:

$$(16) \quad \Omega^{(0)} \subset \Omega^{(1)} \subset \dots \subset \Omega^{(n-2)} \subset \Omega^{(n-1)} := \partial\Omega \subset \Omega^{(n)} := \bar{\Omega}.$$

We recall that a *smooth stratification* $S_0 \subset S_1 \subset \dots \subset X$ of a topological space X is an increasing sequence of closed sets $S_j = S_j(X)$ such that each point of X has a neighborhood that meets only finitely many of the sets S_j , S_0 is a discrete subset, $S_{j+1} \setminus S_j$, $j \geq 0$, is a disjoint union of smooth manifolds of dimension $j + 1$, and $X = \cup S_j$. Some of the sets S_j may be empty for $0 \leq j \leq j_0 < \dim(X)$.

We will always assume that the stratification $\{\Omega^{(j)}\}$ satisfies the condition $\Omega^{(j)} \setminus \Omega^{(j-1)}$ has finitely many connected components. This assumption is automatically satisfied if Ω is bounded, and it is not crucial, but simplifies some of the later constructions.

We proceed by induction on the dimension to define a polyhedral structure on Ω . Our definition is very closely related to that of Whitney stratified spaces [58]. We first agree that a curvilinear polyhedral domain of dimension $n = 0$ is simply a finite set of points. Then, we assume that we have defined curvilinear polyhedral domains in dimension $\leq n - 1$, $n \geq 1$, and define a curvilinear polyhedral domain in a manifold M of dimension n next. We shall denote by B^l the open unit ball in \mathbb{R}^l and by $S^{l-1} := \partial B^l$ its boundary. In particular, we identify $B^0 = \{1\}$, $B^1 = (-1, 1)$, and $S^0 = \{-1, 1\}$.

Definition 2.1. Let M be a smooth manifold of dimension $n \geq 1$. Let $\Omega \subset M$ be an open subset endowed with the stratification (16). Then $\Omega \subset M$ is a *stratified, curvilinear polyhedral domain* if for every point $p \in \partial\Omega$, there exist a neighborhood V_p in M such that:

- (i) if $p \in \Omega^{(l)} \setminus \Omega^{(l-1)}$, $l = 1, \dots, n - 1$, there is a curvilinear polyhedral domain $\omega_p \subset S^{n-l-1}$, $\bar{\omega}_p \neq S^{n-l-1}$, and
 - (ii) a diffeomorphism $\phi_p : V_p \rightarrow B^{n-l} \times B^l$ such that $\phi_p(p) = 0$ and
- $$(17) \quad \phi_p(\Omega \cap V_p) = \{rx', 0 < r < 1, x' \in \omega_p\} \times B^l,$$

inducing a homeomorphism of stratified spaces.

Given any $p \in \partial\Omega$, let $0 \leq \ell(p) \leq n - 1$ be the smallest integer such that $p \in \Omega^{(\ell(p))}$, but $p \notin \Omega^{(\ell(p)-1)}$ (by convention we set $\Omega^{(l)} = \emptyset$ if $l < 0$). By construction, $\ell(p)$ is unique given p . Then, the domain $\omega_p \subset S^{n-\ell(p)-1}$ in the definition above will be called the *link of Ω at p* . Recall that we identify the "ball" $B^0 = \{1\}$ and the "sphere" $S^0 = \partial B^1 = \{-1, 1\}$. In particular if $\ell(p) = n - 1$, then Ω_P is a point.

The notion of a stratified polyhedron is well known in the literature (see for example the monograph [53]). However, Our definition is more general, and well suited for applications to partial differential equations. See the paper of Babuška and Guo [5], the paper of Mazya and Rossmann [39], and the papers and Verchota and Vogel [57, 56] for related definitions. We remark that according to the above definition, Ω does not need to be bounded, nor connected, nor convex. *For applications to the analysis of boundary value/interface problems, however, we will always assume Ω is connected.* The boundary $\partial\Omega$ need not be connected either, but it does have finitely many connected components. We also stress that polyhedral domains will always be open subsets.

The condition $\overline{\omega_p} \neq S^{n-l-1}$ can be relaxed to $\omega_p \neq S^{n-l-1}$, thus allowing for cracks and slits, but not punctured domains of the form $M \setminus \{p\}$. We will not pursue this generality in the paper, given also that submanifolds of codimension greater than 2 consists of irregular boundary points for elliptic equations and may lead to ill-posedness in boundary value problems. We refer to the articles [40, 35] for a detailed analysis of polyhedral domains with cracks in 2 and 3 dimensions.

We continue with some comments on the definition before providing several concrete examples in dimension $n = 1, 2, 3$. We denote by tB^l the ball of radius t in \mathbb{R}^l , $l \in \mathbb{N}$, centered at the origin. We also let tB^0 to be a point independent of t . Sometimes it is convenient to replace Condition (17) with the equivalent condition that there exist $t > 0$ such that

$$(18) \quad \phi_p(\Omega \cap V_p) = \{rx', 0 < r < t, x' \in \omega_p\} \times tB^l.$$

We shall interchange conditions (17) and (18) at will from now on. For a cone or an infinite wedge, $t = +\infty$, so cones and wedges are particular examples of polyhedral domain.

We have the following simple result that is an immediate consequence of the definitions.

Proposition 2.2. *Let $\psi : M \rightarrow M'$ be a diffeomorphism and let $\Omega \subset M$ be a curvilinear polyhedral domain. Then $\psi(\Omega)$ is also a curvilinear polyhedral domain.*

Next, we introduce the *singular set* of Ω , $\Omega_{\text{sing}} := \Omega^{(n-2)}$. A point $p \in \Omega^{(n-2)}$ will be called a *singular point* for Ω . We recall that a point $x \in \partial\Omega$ is called a *smooth boundary point* of Ω if the intersection of $\partial\Omega$ with a small neighborhood of p is a smooth manifold of dimension $n - 1$. In view of Definition 2.1, the point p is smooth if ϕ_p defines a diffeomorphism

$$(19) \quad \phi_p(\Omega \cap V_p) = (0, t) \times B^{n-1}.$$

This observation is consistent with ω_p being a point in this case, since it is a polyhedral domain of dimension 0.

Any point $p \in \partial\Omega$ that is not a smooth boundary point in this sense is a singular point. But the singular set may include other points as well, in particular the points where the boundary conditions change, *i.e.*, the points of the boundary of $\partial_D\Omega$ in $\partial\Omega$, and the points where the interface Γ meets $\partial\Omega$. It is known [25, 26] that the solution to the problem (6) near such points behaves in a similar way as in the neighborhood of non-smooth boundary points. We call the non-smooth points in $\partial\Omega$ the *true* or *geometric* singular points, while we call all the other singular points as *artificial* singular points.

The true singular points can be characterized by the condition that the domain ω_p of Definition 2.1 be an “irreducible” subset of the sphere S^{n-l-1} , in the sense of the following definition.

Definition 2.3. A subset $\omega \subset S^{k-1} := \partial B^k$, the unit sphere in \mathbb{R}^k will be called *irreducible* if $\mathbb{R}_+\omega := \{rx', r > 0, x' \in \omega\}$ cannot be written as $V + V'$ for a linear subspace $V \subset \mathbb{R}^k$ of dimension ≥ 1 and V' an arbitrary subset of \mathbb{R}^{n-k} . (The sum does not have to be a direct sum and, in fact, V' is not assumed to be an affine subspace.)

For example, $(0, \alpha) \subset S^1$ is irreducible if, and only if, $\alpha \neq \pi$. A subset $\omega \subset S^{k-1}$ strictly contained in an open half-space is irreducible, but the intersection of S^{k-1} , $k \geq 2$, with an open half-space is not irreducible.

If $p \in \Omega^{(0)}$, then we shall call p a *vertex* of Ω and we shall interpret the condition (17) as saying that ϕ_p defines a diffeomorphism such that

$$(20) \quad \phi_p(\Omega \cap V_p) = \{rx', 0 < r < t, x' \in \omega_p\}.$$

This interpretation is consistent with our convention that the set B^0 (the zero dimensional unit ball) consists of a single point. We shall call any open, connected component of $\Omega^{(1)} \setminus \Omega^{(0)}$ an (open) *edge* of Ω , necessarily a smooth curve in M . Similarly, any open, connected component of $\Omega^{(j)} \setminus \Omega^{(j-1)}$ shall be called a (open) *j-face* if $2 \leq j \leq n-1$. A $n-1$ -face will be called a *hyperface*. A *j-face* H is a smooth manifold of dimension j , but in general it is not a curvilinear polyhedral domain (except if $n=2$), because there might not exist a j -manifold containing the closure of H in $\partial\Omega$. This point will be addressed in terms of the desingularization $\Sigma(\Omega)$ of Ω constructed in Section 4.

Notations 2.4. From now on, Ω will denote a curvilinear polyhedral domain in a manifold M of dimension n with given stratification $\Omega^{(0)} \subset \Omega^{(1)} \subset \dots \subset \Omega^{(n)} := \bar{\Omega}$.

Some or all of the sets $\Omega^{(j)}$, $j = 0, \dots, n-2$, in the stratification of Ω may be empty. In fact, $\Omega^{(n-2)}$ is empty if, and only if, $\bar{\Omega}$ is a smooth manifold, possibly with boundary, a particular case of a curvilinear, stratified polyhedron. Finally we introduce the notion of distance to the singular set $\Omega^{(n-2)}$ of Ω (if not empty) on which the constructions of the Sobolev spaces $\mathcal{K}_a^\mu(\Omega)$ given in Section 5 is based. If $\Omega^{(n-2)} = \emptyset$, we let $\eta_{n-2} \equiv 1$.

Definition 2.5. Let Ω be a curvilinear, stratified polyhedral domain of dimension n . The distance function $\eta_{n-2}(x)$ from x to the singular set $\Omega^{(n-2)}$ is

$$(21) \quad \eta_{n-2}(x) := \inf_{\gamma} \ell(\gamma),$$

where $\ell(\gamma)$ is the length of the curve γ , and γ ranges through all smooth curves $\gamma : [0, 1] \rightarrow \bar{\Omega}$, $\gamma(0) = x$, $p := \gamma(1) \in \Omega^{(n-2)}$.

If Ω is not bounded, for example Ω is an infinite cone, then we modify the definition of the distance function as follows:

$$(22) \quad \eta_{n-2}(x) := \chi(\inf_{\gamma} \ell(\gamma)), \quad \text{where}$$

$$\chi \in C^\infty([0, +\infty)), \quad \chi(s) = \begin{cases} s, & 0 \leq s \leq 1 \\ \geq 1, & s \geq 1 \\ 2, & s \geq 3. \end{cases}$$

2.1. Curvilinear polyhedral domains in 1, 2, and 3 dimensions. In this subsection we give some examples of curvilinear polyhedral domains Ω in \mathbb{R}^2 , in S^2 , or in \mathbb{R}^3 . These examples are crucial in understanding Definition 2.1, which we specialize here for $n = 2, n = 3$. The desingularization $\Sigma(\Omega)$ and the function r_Ω will be introduced in the next subsection in these special cases.

We have already defined a polyhedron in dimension 0 as a finite collection of points. Accordingly, a subset $\Omega \subset \mathbb{R}$ or $\Omega \subset S^1$ is a *curvilinear polyhedral domain* if, and only if, it is a finite union of open intervals.

Let M be a smooth 2-manifold or \mathbb{R}^2 . Definition 2.1 can be more explicitly stated as follows.

Definition 2.6. A subset $\Omega \subset M$ together with smooth stratification $\Omega^{(0)} \subset \Omega^{(1)} \equiv \partial\Omega \subset \Omega^{(2)} \equiv \Omega$ will be called a *curvilinear, stratified polygonal domain* if, for every point of the boundary $p \in \partial\Omega$, there exists a neighborhood $V_p \subset M$ of p and a diffeomorphism $\phi_p : V_p \rightarrow B^2$, $\phi_p(p) = 0$, such that:

- (a) $\phi_p(V_p \cap \Omega) = \{ (r \cos \theta, r \sin \theta), 0 < r < 1, \theta \in \omega_p \}$, where ω_p is a union of open intervals of the unit circle such that $\overline{\omega_p} \neq S^1$;
- (b) if $p \in \Omega^{(1)} \setminus \Omega^{(0)}$, then ω_p is exactly an interval of length π .

Any point $p \in \Omega^{(0)}$ is a *vertex* of Ω , and p is a true vertex precisely when ω_p is not an interval of length π . The open, connected components of $\partial\Omega \setminus \Omega^{(0)}$ are the (open) *sides* of Ω . In view of condition (b) above, sides are smooth curves $\gamma_j : [0, 1] \rightarrow M, j = 1, \dots, N$, with no common interior points. Recall that by hypothesis, there are finitely many vertices and sides. The condition that $\overline{\omega_p} \neq S^1$ implies that either a side γ_j has a vertex in common with another side γ_k or γ_j is a closed smooth curve or an infinite smooth curve. In the special case $\Omega^{(1)} \setminus \Omega^{(0)} = \emptyset$, Ω has only isolated conical points (see Example 25 in the next subsection), while if $\Omega^{(0)} = \emptyset$, Ω has smooth boundary.

Notations 2.7. Any *curvilinear, stratified polygon* in \mathbb{R}^2 will be denoted by \mathbb{P} and its stratification by $\mathbb{P}^{(0)} \subset \mathbb{P}^{(1)} \subset \dots \subset \mathbb{P}^{(n-1)} = \partial\mathbb{P} \subset \mathbb{P}^{(n)} = \overline{\mathbb{P}}$.

Let now M be a smooth 3-manifold or \mathbb{R}^3 . Definition 2.1 can also be stated more explicitly.

Definition 2.8. A subset $\Omega \subset M$ together with a smooth stratification $\Omega^{(0)} \subset \Omega^{(1)} \subset \Omega^{(2)} \equiv \partial\Omega \subset \Omega^{(3)} \equiv \overline{\Omega}$ will be called a *curvilinear, stratified polyhedral domain* if, for every point of the boundary $p \in \partial\Omega$, there exists a neighborhood $V_p \subset M$ of p and a diffeomorphism $\phi_p : V_p \rightarrow B^l \times B^{3-l}, \phi_p(p) = 0$, such that:

- (a) $\phi_p(V_p \cap \Omega) = \{ (y, rx'), y \in B^j, 0 < r < t, x' \in \omega_p \}$, where $\omega_p \subset S^{2-l}$ is such that $\overline{\omega_p} \neq S^2, t \in (0, +\infty]$;
- (b) if $l = 0$ (i.e., if $p \in \Omega^{(0)}$), then $\omega_p \subset S^2$ is a stratified, curvilinear polygonal domain;
- (c) if $l = 1$ (i.e., if $p \in \Omega^{(1)} \setminus \Omega^{(0)}$), then ω_p is a finite, disjoint union of finitely many open intervals in S^1 of total length less than 2π .
- (d) if $l = 2$ then p is a smooth boundary point;
- (e) ϕ_p preserves the stratifications;

Each point $p \in \Omega^{(0)}$ is a *vertex* of Ω and p is a true vertex precisely when ω_p is an irreducible subset of S^2 (according to Definition 2.3). The open, connected components of $\Omega^{(1)} \setminus \Omega^{(0)}$ are the *edges* of Ω , smooth curves with no interior common

points by condition (c) above. The open, connected components of $\Omega^{(2)} \setminus \Omega^{(1)}$, smooth surfaces with no common interior points, are the *faces* of Ω . Recall that by hypothesis, there are only finitely many vertices, edges, and faces in Ω . The condition that $\overline{\omega_p}$ is not the whole sphere S^{2-l} ($l = 1, 0$) implies that either an edge γ_j has a vertex in common with another edge γ_k or γ_j is a closed smooth curve or an infinite smooth curve (such as in a wedge), and similarly for faces. Again, in the the case $\Omega^{(1)} = \Omega^{(0)}$, Ω has only isolated conical points, in the case $\Omega^{(0)} = \emptyset$, Ω has only edge singularities, and in the case $\Omega^{(1)} = \Omega^{(0)} = \emptyset$, Ω is smooth.

The following subsection contains several examples.

2.2. Definition of $\Sigma(\Omega)$ and of r_Ω if $n = 2$ or $n = 3$. We now introduce the desingularization $\Sigma(\Omega)$ for some of the typical examples of curvilinear polyhedral domains in $n = 2$ or $n = 3$. Associated to the singularization is the function r_Ω , which is comparable with the distance to the singular set η_{n-2} but is more regular. We also frame these definitions as examples. The general case (of which the examples considered here are particular cases) is in Section 4. The reader can skip this part at first.

The case $n = 2$ of a polygonal domain \mathbb{P} in \mathbb{R}^2 is particularly simple. We use the notation in Definition 2.6.

Example 2.9. The desingularization $\Sigma(\mathbb{P})$ of \mathbb{P} will replace each of the vertices A_j , $j = 1, \dots, k$, of \mathbb{P} with a segment of length $\alpha_j > 0$, where α_j is the magnitude of the angle at A_j (if A_j is an artificial vertex, then $\alpha_j = \pi$). We can realize $\Sigma(\mathbb{P})$ in three dimensions as follows. Let θ_j be the angle in a polar coordinates system (r_j, θ_j) centered at A_j . Let ϕ_j be a smooth function on \mathbb{P} that is equal to 1 on $\{r_j < \epsilon\}$ and vanishes outside $V_j := \{r_j < 2\epsilon\}$. By choosing $\epsilon > 0$ small enough, we can arrange that the sets V_j do not intersect. We define then

$$\Phi : \overline{\mathbb{P}} \setminus \{A_1, A_2, \dots, A_k\} \rightarrow \mathbb{P} \times \mathbb{R} \subset \mathbb{R}^3$$

by $\Phi(p) = (p, \sum \phi_j(p)\theta_j(p))$. Then $\Sigma(\mathbb{P})$ is (up to a diffeomorphism) the closure in \mathbb{R}^3 of $\Phi(\mathbb{P})$. The desingularization map is $\kappa(p, z) = p$. The structural Lie algebra of vector fields $\mathcal{V}(\mathbb{P})$ on $\Sigma(\mathbb{P})$ is given by (the lifts of) the smooth vector fields X on $\overline{\mathbb{P}} \setminus \{A_1, A_2, \dots, A_k\}$ that on $V_j = \{r_j < 2\epsilon\}$ can be written as

$$(23) \quad X = a_r(r_j, \theta_j)r_j\partial_{r_j} + a_\theta(r_j, \theta_j)\partial_{\theta_j},$$

with a_r and a_θ smooth functions of (r_j, θ_j) on $[0, 2\epsilon] \times [0, \alpha_j]$. We can take $r_\Omega(x) := \psi(x) \prod_{j=1}^k r_j(x)$, where ψ is a smooth, nowhere vanishing function on $\Sigma(\Omega)$. (Such a factor ψ can always be introduced, and the function r_Ω is determined only up to this factor. We shall omit this factor in the examples below.)

The example of a domain with a single edge or vertex is one of the most instructive.

Example 2.10. Let first Ω be the wedge

$$(24) \quad \mathbb{W} := \{(r \cos \theta, r \sin \theta, z), 0 < r, 0 < \theta < \alpha, z \in \mathbb{R}\},$$

where $0 < \alpha < 2\pi$, and $x = r \cos \theta$ and $y = r \sin \theta$ define the usual cylindrical coordinates (r, θ, z) , with $(r, \theta, z) \in [0, \infty) \times [0, 2\pi) \times \mathbb{R}$. Then the manifold of generalized cylindrical coordinates is, in this case, just the domain of the cylindrical coordinates on $\overline{\mathbb{W}}$:

$$\Sigma(\mathbb{W}) = [0, \infty) \times [0, \alpha] \times \mathbb{R}.$$

The desingularization map is $\kappa(r, \theta, z) = (r \cos \theta, r \sin \theta, z)$ and the structural Lie algebra of vector fields of $\Sigma(\mathbb{W})$ is

$$a_r(r, \theta, z)r\partial_r + a_\theta(r, \theta, z)\partial_\theta + a_z(r, \theta, z)r\partial_z,$$

where a_r , a_z , and a_θ are smooth functions on $\Sigma(\mathbb{W})$. Note that the vector fields in $\mathcal{V}(\mathbb{W})$ may not extend to the closure $\overline{\mathbb{W}}$. We can take $r_\Omega = r$, the distance to the Oz -axis.

At this stage, we can describe a domain with one conical point and its desingularization in any dimension.

Example 2.11. Let next Ω be a domain with one conical point, that is, Ω is a curvilinear, stratified polyhedron in \mathbb{R}^n such that $\Omega^{(j)} = \Omega^{(0)}$ for all $1 \leq j \leq n - 2$. We assume Ω is bounded for simplicity. Let $p \in \Omega^{(0)}$ be the single vertex of Ω . Then there exists a neighborhood V_p of p such that, up to a local change of coordinates,

$$(25) \quad V_p \cap \Omega = \{rx', 0 \leq r < \epsilon, x' \in \omega\},$$

for some smooth, connected domain $\omega \subset S^{n-1} := \partial B^n$. Then we can realize $\Sigma(\Omega)$ in \mathbb{R}^{2n} as follows. Assume $p = 0$, the origin, for simplicity. We define $\Phi(x) = (x, |x|^{-1}x)$ for $x \neq p$, where $|x|$ is the distance from x to the origin (*i.e.*, to p). Then $\Sigma(\Omega)$ is defined to be the closure of the range of Φ . The map κ is the projection onto the first n components. Then κ is one-to-one, except that $\kappa^{-1}(p) = \{p\} \times \omega$. We can take $r_\Omega(x) = |x|$. The Lie algebra of vector fields $\mathcal{V}(\Omega)$ consists of the vector fields on $\Sigma(\Omega)$ that are tangent to $\kappa^{-1}(p)$. This example is due to Melrose [42].

Example 2.12. Let $\Omega \subset \mathbb{R}^3$ be a convex polyhedral domain, such that all edges are straight segments. To construct $\Sigma(\Omega)$, we combine the ideas used in the previous examples. First, for each edge e we define (r_e, θ_e, z_e) to be a coordinate system aligned to that edge and such that $\theta_e \in (0, \alpha_e)$, as in Example 2.10. Let v_1, v_2, \dots, v_b be the set of vertices of Ω and e_1, \dots, e_a be the set of edges. Then, for x not on any edge of Ω , we define $\Phi(x) \in \mathbb{R}^{3+a+b}$ by

$$\Phi(x) = (x, \theta_{e_1}, \theta_{e_2}, \dots, \theta_{e_a}, |x - v_1|^{-1}(x - v_1), \dots, |x - v_b|^{-1}(x - v_b)).$$

The desingularization $\Sigma(\Omega) \subset \mathbb{R}^{3+a+b}$ is defined as the closure of the range of Φ . The resulting set will be a manifold with corners with several different types of hyperfaces. Namely, the manifold $\Sigma(\Omega)$ will have a hyperface for each face of Ω , a hyperface for each edge of Ω , and, finally, a hyperface for each vertex of Ω . The last two types of hyperfaces are the so-called *hyperfaces at infinity* of $\Sigma(\Omega)$. Let x_H be the distance to the hyperface H . We can take then $r_\Omega = \prod_H x_H$, where H ranges through the hyperfaces at infinity of $\Sigma(\Omega)$.

We can imagine $\Sigma(\Omega)$ as follows. Let $\epsilon > 0$. Remove the sets $\{x \in \Omega, |x - v_j| \leq \epsilon\}$ and $\{x \in \Omega, |x - e_k| \leq \epsilon^2\}$. Call the resulting set Ω_ϵ . Then, for ϵ small enough, the closure of Ω_ϵ is diffeomorphic to $\Sigma(\Omega)$.

The example above can be generalized to a curvilinear, stratified polyhedron, using local change of coordinates as in Example 2.9 in 2 dimensions. A detailed construction will be given in Section 4.

A nonstandard example of curvilinear polyhedral domain is given below.

Example 2.13. We start with a connected polygonal domain \mathbb{P} with connected boundary and we deform it until one, and exactly one of the vertices, say A , touches

the interior of another edge, say $[B, C]$. Let Ω be the resulting connected open set. Ω will be a curvilinear polyhedral domain. We define the set $\Sigma(\Omega)$ as for the polygonal domain \mathbb{P} , but by introducing polar coordinates in the whole neighborhood of the point A .

If we deform \mathbb{P} to Ω , $\Sigma(\mathbb{P})$ will deform continuously to a space $\Sigma'(\Omega)$, different from $\Sigma(\Omega)$. For certain purposes, the desingularization $\Sigma'(\Omega)$ is better suited than $\Sigma(\Omega)$.

3. LIE MANIFOLDS WITH BOUNDARY

Both the construction of the desingularization of a general, curvilinear, stratified polyhedron in n dimensions in Section 4, used in the definition of weighted Sobolev spaces on the boundary, and the proof of a weighted Hardy-Poincaré inequality in Subsection 6.2, crucial in establishing coercive estimates for the mixed boundary value/interface problem (6), rely on properties of manifolds with a Lie structure at infinity.

In order to make this paper as self-contained as possible, we now recall the definition of a Lie manifold from [3] and of a Lie manifold with boundary from [1]. We also recall a few other needed definitions and results from those papers.

3.1. Definition. We recall that a topological space \mathfrak{M} is, by definition, a *manifold with corners* if every point $p \in \mathfrak{M}$ has a coordinate neighborhood homeomorphic to $[0, 1)^k \times (-1, 1)^{n-k}$, $k = 1, \dots, n-1$, such that the transition functions are smooth (including at the boundary). Given $p \in \mathfrak{M}$, the least integer k with the above property is called the *depth* of p . Since the transition functions are smooth, it therefore makes sense to talk about smooth functions on \mathfrak{M} , these being the functions that correspond to smooth functions on $[0, 1)^k \times (-1, 1)^{n-k}$. We denote by $\mathcal{C}^\infty(\mathfrak{M})$ the set of smooth functions on a manifold with corners \mathfrak{M} .

Throughout this paper, \mathfrak{M} will denote a manifold with corners, not necessarily compact. We shall denote by \mathfrak{M}_0 the interior of \mathfrak{M} and by $\partial\mathfrak{M} = \mathfrak{M} \setminus \mathfrak{M}_0$ the boundary of \mathfrak{M} . The set \mathfrak{M}_0 consists of the set of points of depth zero of \mathfrak{M} . It is usually no loss of generality to assume that \mathfrak{M}_0 is connected. Let \mathfrak{M}_k denote the set of points of \mathfrak{M} of depth k and F_0 be a connected component of \mathfrak{M}_k . We shall call F_0 an *open face of codimension k* of \mathfrak{M} and $F := \overline{F_0}$ a *face of codimension k* of \mathfrak{M} . A face of codimension 1 will be called a *hyperface* of \mathfrak{M} , so that $\partial\mathfrak{M}$ is the union of all hyperfaces of \mathfrak{M} . In general, a face of \mathfrak{M} need not be a smooth manifold (with or without corners). A face $F \subset \mathfrak{M}$ which is a submanifold with corners of \mathfrak{M} will be called an *embedded face*.

Anticipating, a Lie manifold with boundary \mathfrak{M}_0 is the interior of a manifold with corners \mathfrak{M} together with a Lie algebra of vector fields \mathcal{V} on \mathfrak{M} satisfying certain conditions. To state these conditions, it will be convenient of first introduce a few other concepts.

Definition 3.1. Let \mathfrak{M} be a manifold with corners and \mathcal{V} be a space of vector fields on \mathfrak{M} . Let $U \subset \mathfrak{M}$ be an open set and Y_1, Y_2, \dots, Y_k be vector fields on $U \cap \mathfrak{M}_0$. We shall say that Y_1, Y_2, \dots, Y_k form a *local basis of \mathcal{V} on U* if the following three conditions are satisfied:

- (i) there exist vector fields $X_1, X_2, \dots, X_k \in \mathcal{V}$, $Y_j = X_j$ on $U \cap \mathfrak{M}_0$;
- (ii) \mathcal{V} is closed under products with smooth functions in $\mathcal{C}^\infty(\mathfrak{M})$ (i.e., $\mathcal{V} = \mathcal{C}^\infty(\mathfrak{M})\mathcal{V}$) and for any $X \in \mathcal{V}$, there exist smooth functions $\phi_1, \phi_2, \dots, \phi_k \in$

$C^\infty(\mathfrak{M}_0)$ such that

$$(26) \quad X = \phi_1 X_1 + \phi_2 X_2 + \dots + \phi_k X_k \quad \text{on } U \cap \mathfrak{M}_0;$$

and

- (iii) if $\phi_1, \phi_2, \dots, \phi_k \in C^\infty(\mathfrak{M})$ and $\phi_1 X_1 + \phi_2 X_2 + \dots + \phi_k X_k = 0$ on $U \cap \mathfrak{M}_0$, then $\phi_1 = \phi_2 = \dots = \phi_k = 0$ on U .

We now recall structural Lie algebras of vector fields from [3].

Definition 3.2. A subspace $\mathcal{V} \subseteq \Gamma(\mathfrak{M}, T\mathfrak{M})$ of the Lie algebra of all smooth vector fields on \mathfrak{M} is said to be a *structural Lie algebra of vector fields on \mathfrak{M}* provided that the following conditions are satisfied:

- (i) \mathcal{V} is closed under the Lie bracket of vector fields;
- (ii) every vector field $X \in \mathcal{V}$ is tangent to all hyperfaces of \mathfrak{M} ;
- (iii) $C^\infty(\mathfrak{M})\mathcal{V} = \mathcal{V}$; and
- (iv) for each point $p \in \mathfrak{M}$ there exist a neighborhood U_p of p in \mathfrak{M} and a local basis of \mathcal{V} on U_p .

The concept of Lie structure at infinity, defined next, is also taken from [3].

Definition 3.3. A *Lie structure at infinity* on a smooth manifold \mathfrak{M}_0 is a pair $(\mathfrak{M}, \mathcal{V})$, where \mathfrak{M} is a compact manifold, possibly with corners, and $\mathcal{V} \subset \Gamma(\mathfrak{M}, T\mathfrak{M})$ is a structural Lie algebra of vector fields on \mathfrak{M} with the following properties:

- (i) $\mathfrak{M}_0 = \mathfrak{M} \setminus \partial\mathfrak{M}$, the interior of \mathfrak{M} , and
- (ii) If $p \in \mathfrak{M}_0$, then any local basis of \mathcal{V} in a neighborhood of p is also a local basis of the tangent space to \mathfrak{M}_0 . (In particular, the constant k of Equation (26) equals n , the dimension of \mathfrak{M}_0 .)

A *manifold with a Lie structure at infinity* (or, simply, a *Lie manifold*) is a manifold \mathfrak{M}_0 together with a Lie structure at infinity $(\mathfrak{M}, \mathcal{V})$ on \mathfrak{M}_0 . We shall sometimes denote a Lie manifold as above by $(\mathfrak{M}_0, \mathfrak{M}, \mathcal{V})$, or, simply, by $(\mathfrak{M}, \mathcal{V})$, because \mathfrak{M}_0 is determined as the interior of \mathfrak{M} .

Let \mathcal{V}_b be the set of vector fields on \mathfrak{M} that are tangent to all faces of \mathfrak{M} . Then $(\mathfrak{M}, \mathcal{V}_b)$ is a Lie manifold [42, 41]. See [34, 3, 1] for more examples.

3.2. Riemannian metric. Let $(\mathfrak{M}, \mathcal{V})$ be a Lie manifold and g a Riemannian metric on $\mathfrak{M}_0 := \mathfrak{M} \setminus \partial\mathfrak{M}$. We shall say that g is *compatible* (with the Lie structure at infinity $(\mathfrak{M}, \mathcal{V})$) if, for any $p \in \mathfrak{M}$, there exist a neighborhood U_p of p in \mathfrak{M} and a local basis X_1, X_2, \dots, X_n of \mathcal{V} on U_p that is orthonormal with respect to g on U_p .

It was proved in [3] that (\mathfrak{M}_0, g_0) is necessarily of infinite volume and complete. Moreover, all the covariant derivatives of the Riemannian curvature tensor of g are bounded. Under additional mild assumptions, we also know that the injectivity radius is bounded from below by a positive constant, *i.e.*, (\mathfrak{M}_0, g_0) is of bounded geometry. (A *manifold with bounded geometry* is a Riemannian manifold with positive injectivity radius and with bounded covariant derivatives of the curvature tensor, see for example [9] or [52] and references therein). We assume from now on that $r_{\text{inj}}(\mathfrak{M}_0)$, the injectivity radius of (\mathfrak{M}_0, g_0) , is positive.

3.3. \mathcal{V} -differential operators. We are especially interested in the analysis of the differential operators generated using only derivatives in \mathcal{V} . Let $\text{Diff}_{\mathcal{V}}^*(\mathfrak{M})$ be the algebra of differential operators on \mathfrak{M} generated by multiplication with functions in $\mathcal{C}^\infty(\mathfrak{M})$ and by differentiation with vector fields $X \in \mathcal{V}$. The space of order m differential operators in $\text{Diff}_{\mathcal{V}}^*(\mathfrak{M})$ will be denoted $\text{Diff}_{\mathcal{V}}^{*m}(\mathfrak{M})$. A differential operator in $\text{Diff}_{\mathcal{V}}^*(\mathfrak{M})$ will be called a \mathcal{V} -differential operator. We define the set $\text{Diff}_{\mathcal{V}}^*(\mathfrak{M}; E, F)$ of \mathcal{V} -differential operators acting between sections of smooth vector bundles $E, F \rightarrow \mathfrak{M}$ in the usual way [1, 3].

A simple but useful property of the differential operator in $\text{Diff}_{\mathcal{V}}^*(\mathfrak{M})$ is that

$$(27) \quad x^s P x^{-s} \in \text{Diff}_{\mathcal{V}}^*(\mathfrak{M})$$

for any $P \in \text{Diff}_{\mathcal{V}}^*(\mathfrak{M})$ and any defining function x of some hyperface of \mathfrak{M} [41, 2]. This property is easily proved using the fact that X is tangent to the hyperface defined by x , for any $X \in \mathcal{V}$ (a proof of a slightly more general fact is included in Corollary 6.3).

3.4. Lie manifolds with boundary. A subset $\mathfrak{N} \subset \mathfrak{M}$ is called a *submanifold with corners* of \mathfrak{M} if \mathfrak{N} is a closed submanifold of \mathfrak{M} such that \mathfrak{N} is transverse to all faces of \mathfrak{M} and any face of \mathfrak{N} is a component of $\mathfrak{N} \cap F$ for some face F of \mathfrak{M} .

The following definition is a reformulation of a definition of [1].

Definition 3.4. Let $(\mathfrak{N}, \mathcal{W})$ and $(\mathfrak{M}, \mathcal{V})$ be Lie manifolds, where $\mathfrak{N} \subset \mathfrak{M}$ is a submanifold with corners and

$$\mathcal{W} = \{X|_{\mathfrak{N}}, X \in \mathcal{V}, X|_{\mathfrak{N}} \text{ tangent to } \mathfrak{N}\}.$$

We shall say that $(\mathfrak{N}, \mathcal{W})$ is a *tame* submanifold of $(\mathfrak{M}, \mathcal{V})$ if, for any $p \in \partial\mathfrak{N}$ and any $X \in T_p\mathfrak{M}$, there exist $Y \in \mathcal{V}$ and $Z \in T_p\mathfrak{N}$ such that $X = Y(p) + Z$.

Let $\mathfrak{N} \subset \mathfrak{M}$ be a submanifold with corners. We assume that \mathfrak{M} and \mathfrak{N} are endowed with the Lie structures $(\mathfrak{N}, \mathcal{W})$ and $(\mathfrak{M}, \mathcal{V})$. We shall say that \mathfrak{N} is a *regular* submanifold of $(\mathfrak{M}, \mathcal{V})$ if we can choose a tubular neighborhood V of $\mathfrak{N}_0 := \mathfrak{N} \setminus \partial\mathfrak{N} = \mathfrak{N} \cap \mathfrak{M}_0$ in \mathfrak{M}_0 , a compatible metric g_1 on \mathfrak{N}_0 , a product-type metric g_1 on V that reduces to g_1 on \mathfrak{N}_0 , and a compatible metric on \mathfrak{M}_0 that restricts to g_1 on V . Theorem 5.8 of [3] states that every tame submanifold is regular. The point of this result is that it is much easier to check that a submanifold is tame than to check that it is regular.

In the case when \mathfrak{N} is of codimension one in \mathfrak{M} , the condition that \mathfrak{N} be tame is equivalent to the fact that there exists a vector field $X \in \mathcal{V}$ that restricts to a normal vector of \mathfrak{N} in \mathfrak{M} . The neighborhood V will then be of the form $V \simeq (\partial\mathfrak{N}_0) \times (-\varepsilon_0, \varepsilon_0)$. Moreover, there will exist a compatible metric on \mathfrak{M}_0 that restricts to the product metric $g_1 + dt^2$ on V , where g_1 is a compatible metric on \mathfrak{N}_0 .

Let A be a subset of \mathfrak{M} . We denote by $\partial_{\mathfrak{M}}A$ the boundary of A computed within \mathfrak{M} , which should not be confused with $\partial A = A \setminus A_0$, where A_0 is the interior of A . Let $\mathbb{D} \subset \mathfrak{M}$ be an open subset. We say that \mathbb{D} is a *Lie domain* in \mathfrak{M} if, and only if, $\partial_{\mathfrak{M}}\mathbb{D} = \partial_{\mathfrak{M}}\mathbb{D}$ and $\partial_{\mathfrak{M}}\mathbb{D}$ is a regular submanifold of \mathfrak{M} . A typical example of a Lie domain $\mathbb{D} \subset \mathfrak{M}$ is obtained by considering a regular submanifold with corners $\mathfrak{N} \subset \mathfrak{M}$ of codimension one with the property that $\mathfrak{M} \setminus \mathfrak{N}$ consists of two connected components. Any of these two components will be a Lie domain.

Definition 3.5. A *Lie manifold with boundary* is a triple $(\mathfrak{D}_0, \mathfrak{D}, \mathcal{V}')$, where \mathfrak{D}_0 is a smooth manifold with boundary, \mathfrak{D} is a compact manifold with corners containing \mathfrak{D}_0 as an open subset, and \mathcal{V}' is a Lie algebra of vector fields on \mathfrak{D} with the property that there exists a Lie manifold $(\mathfrak{M}_0, \mathfrak{M}, \mathcal{V})$, a Lie domain \mathbb{D} in \mathfrak{M} and a diffeomorphism $\phi : \mathfrak{D} \rightarrow \overline{\mathbb{D}}$ such that $\phi(\mathfrak{D}_0) = \overline{\mathbb{D}} \cap \mathfrak{M}_0$ and $D\phi(\mathcal{V}|_{\mathbb{D}}) = \mathcal{V}'$.

We continue with some simple observations. First note that if $(\mathfrak{D}_0, \mathfrak{D}, \mathcal{V})$ is a Lie manifold with boundary, then \mathfrak{D}_0 is determined by $(\mathfrak{D}, \mathcal{V})$. Indeed, if we remove from \mathfrak{D} the hyperfaces H with the property that \mathcal{V} consists only of vectors tangent to H , then the resulting set is \mathfrak{D}_0 . Therefore, we can denote the Lie manifold with boundary $(\mathfrak{D}_0, \mathfrak{D}, \mathcal{V})$ simply by $(\mathfrak{D}, \mathcal{V})$.

Another observation is that $\partial\mathfrak{D}_0$, the boundary of \mathfrak{D}_0 has a canonical structure of Lie manifold $(\partial\mathfrak{D}_0, D = \partial_{\mathfrak{M}}\mathbb{D}, \mathcal{W})$, where $\mathcal{W} = \{X|_D, X \in \mathcal{V}, X|_D \text{ is tangent to } D\}$. The compactification D is the closure of $\partial\mathfrak{D}_0$ in \mathfrak{D} .

3.5. Sobolev spaces. The main reason for considering Lie manifolds (with or without boundary) in our setting is that they carry some naturally defined Sobolev spaces and these Sobolev spaces behave almost exactly like the Sobolev spaces on a compact manifold with a smooth boundary. Let us recall one of the equivalent definitions in [1].

Definition 3.6. Fix a Lie manifold $(\mathfrak{M}, \mathcal{V})$. The spaces $L^2(\mathfrak{M}_0) = L^2(\mathfrak{M}_0)$ are defined using the natural volume form on \mathfrak{M}_0 given by an arbitrary compatible metric g on \mathfrak{M}_0 (*i.e.*, compatible with the Lie structure at infinity). All such volume forms are known to define the same space $L^2(\mathfrak{M})$, but with possibly different norms. Let $k \in \mathbb{Z}_+$. Choose a finite set of vector fields $\mathcal{X} \subset \mathcal{V}$ such that $\mathcal{C}^\infty(\mathfrak{M})\mathcal{X} = \mathcal{V}$. The system \mathcal{X} gives rise to the norm

$$(28) \quad \|u\|_{\mathcal{X}, \Omega}^2 := \sum \|X_1 X_2 \dots X_l u\|_{L^2(\Omega)}^2, \quad 1 \leq p < \infty,$$

the sum being over all possible choices of $0 \leq l \leq k$ and all possible choices of vector fields $X_1, X_2, \dots, X_l \in \mathcal{X}$, not necessarily distinct. We then set

$$H^k(\mathfrak{M}_0) = H^k(\mathfrak{M}) := \{u \in L^2(\mathfrak{M}), \|u\|_{\mathcal{X}, \mathfrak{M}} < \infty\}.$$

The spaces $H^s(\mathfrak{M}_0) = H^s(\mathfrak{M})$ are defined by duality (with pivot $L^2(\mathfrak{M}_0)$) when $-s \in \mathbb{Z}_+$, and then by interpolation, as above.

Let $(\mathfrak{D}_0, \mathfrak{D}, \mathcal{V})$ be a Lie manifold with boundary. We shall assume that \mathfrak{D} is the closure of a Lie domain \mathbb{D} of the Lie manifold \mathfrak{M} . The Sobolev spaces $H^k(\mathfrak{D}_0)$ are defined as the set of *restrictions* to \mathfrak{D}_0 of distributions $u \in H^k(\mathfrak{M}_0)$, using the notation of Definition 3.5, $k \in \mathbb{Z}$. In particular, we obtain the following description of $H^k(\mathfrak{D}_0)$ from [1].

Lemma 3.7. *We have, for $k \geq 0$,*

$$H^k(\mathfrak{D}_0) = \{u \in L^2(\mathfrak{M}), \|u\|_{\mathcal{X}} < \infty\},$$

and

$$H^{-k}(\mathfrak{D}_0) = H_0^k(\mathfrak{D}_0)^*,$$

where $H_0^k(\mathfrak{D}_0)$ is the closure of $\mathcal{C}_c^\infty(\mathfrak{D}_0)$ in $H^k(\mathfrak{M}_0)$.

Proof. The first part is obvious (see [1], for example). For the second part, let us notice that the dual of a subspace $Y \subset X$ is the quotient of the dual of X by the annihilator of Y . We then use this result for $Y = H_0^k(\mathfrak{D}_0)$. \square

The hyperfaces of \mathfrak{D} that do not intersect the boundary $\partial\mathfrak{D}_0$ of the manifold with boundary \mathfrak{D}_0 will be called *hyperfaces at infinity*. Let x_H be a defining function of the hyperface H of \mathfrak{D} . Any function of the form $h = \prod x_H^{a_H}$, where H ranges through the set of hyperfaces at infinity of \mathfrak{D} and $a_H \in \mathbb{R}$, will be called an *admissible weight*. If h is an admissible weight, we set

$$hH^\mu(\mathfrak{D}_0) = \{hu; u \in H^\mu(\mathfrak{D}_0)\}$$

with the induced norm.

Later in the paper, we will identify the weighted Sobolev spaces $\mathcal{K}_a^s(\Omega)$ with suitable spaces $hH^s(\mathfrak{D}_0)$ in Proposition 5.8 and utilize the spaces $hH^s(\partial\mathfrak{D}_0)$ to define the spaces $\mathcal{K}_a^s(\partial\Omega)$ on the boundary in Definition 5.9, for Ω a curvilinear, stratified polyhedral domain in dimension n . The following proposition, which summarizes the relevant results from Theorem 3.4 and 3.7 from [1], will then imply Theorem 5.10.

Proposition 3.8. *The restriction to the boundary extends to a continuous, surjective map $hH^\mu(\mathfrak{D}_0) \rightarrow hH^{\mu-1/2}(\partial\mathfrak{D}_0)$, for any $\mu \geq 1$ and any admissible weight h . The kernel of this map, for $\mu = 1$, consists of the closure of $\mathcal{C}_c^\infty(\mathfrak{D}_0)$ in $hH^1(\mathfrak{D}_0)$.*

For \mathbb{D} , \mathfrak{D} , \mathfrak{D}_0 as in the proposition above, $hH^s(\mathbb{D})$, $hH^s(\mathfrak{D})$, and $hH^s(\mathfrak{D}_0)$ will all denote the same space.

4. DESINGULARIZATION OF POLYHEDRA

In this section, we introduce a desingularization procedure that we shall use for studying curvilinear polyhedral domains. The desingularization will carry a natural structure of Lie manifold with boundary. This construction will allow us to study curvilinear polyhedral domains using Lie manifolds with boundary.

As before, $\Omega \subset M$ denotes a curvilinear, stratified polyhedral domain in an n -dimensional manifold M . We shall construct by induction on n a *canonical* manifold with corners $\Sigma(\Omega)$ and a differentiable map $\kappa : \Sigma(\Omega) \rightarrow \overline{\Omega}$ that is a diffeomorphism from the interior of $\Sigma(\Omega)$ to Ω . In particular, the map κ allows us to identify Ω with a subset of $\Sigma(\Omega)$. We shall also construct a canonical Lie algebra of vector fields $\mathcal{V}(\Omega)$ on $\Sigma(\Omega)$. The manifold $\Sigma(\Omega)$ will be called the *desingularization of Ω* , the map κ will be called the *desingularization map*, and the Lie algebra of vector fields will be called the *structural Lie algebra of vector fields of $\Sigma(\Omega)$* . We shall also introduce in this section a smooth weight function r_Ω equivalent to η_{n-2} .

The space $\Sigma(\Omega)$ that we construct is not optimal if the links ω_p are not connected. A better desingularization would be obtained if one considers a diffeomorphism ϕ_{pC} for each connected component C of $V_p \cap \Omega$ that maps C to a conic set of the form $(0, 1)\omega_{pC} \times B^\lambda$, with λ largest possible. The difference between these two constructions is seen by looking at the Example 2.13.

Notations 4.1. *From now on V_p and $\phi_p : V_p \rightarrow tB^{n-l} \times tB^l$, $l = \ell(p)$, will denote a neighborhood of $p \in \partial\Omega$ in $M \supset \Omega$ and ϕ_p will be a diffeomorphism satisfying the conditions of Definition (2.1). In addition, $\omega_p \subset S^{n-l-1}$ will be the curvilinear, stratified polyhedron such that*

$$\phi_p(V_p \cap \Omega) = \{(rx', x''), r \in (0, t), x' \in \omega_p \text{ and } x'' \in tB^l\},$$

i.e., ω_p is the link of Ω at p . This notation will remain fixed throughout the paper.

Recall that $0 \leq \ell(p) \leq n - 1$ is defined to be the smallest integer such that $p \in \Omega^{(\ell(p))}$, but $p \notin \Omega^{(\ell(p)-1)}$. If $\ell(p) = 0$, then B^l is reduced to a point, and we just drop x'' from the notation above. We will assume that ϕ_p extends to the closure of V_p , if necessary.

4.1. The desingularization $\Sigma(\Omega)$. We now define the canonical desingularization of a curvilinear polyhedral domain $\Omega \subset M$, M an n -dimensional smooth manifold. For $n = 0$, Ω consists of finitely many points. Then we define $\Sigma(\Omega) = \Omega$ and $\kappa = id$. To define $\Sigma(\Omega)$ for general Ω , we shall proceed by induction.

We need first to make the important observation that the set ω_p , $p \in \partial\Omega$, of Definition 2.1 is determined up to a linear isomorphism of \mathbb{R}^{n-l-1} . Indeed, let $S_p \subset \partial\Omega$ be maximal connected manifold of dimension $l = \ell(p)$ passing through p that is, the connected component of $\Omega^{(l)} \setminus \Omega^{(l-1)}$ containing p . Let $(T_p S_p)^\perp = T_p M / T_p S_p$. The differential $D\phi_p : T_p M \rightarrow \mathbb{R}^n = T_0 \mathbb{R}^n$ of the map ϕ_p at p has then the property that $D\phi_p(T_p S_p) = T_0 \mathbb{R}^l$, $D\phi_p((T_p S_p)^\perp) = T_0 \mathbb{R}^n / T_0 \mathbb{R}^l = \mathbb{R}^{n-l}$. We will define a canonical set $\mathcal{N}_p \subset (T_p S_p)^\perp$ such that

$$D\phi_p(\mathcal{N}_p) \simeq \mathbb{R}_+ \omega_p.$$

Since the definition of \mathcal{N}_p , which we give next, is independent of any choices used in the definition of a polyhedral domain, it follows that ω_p is unique, up to a linear isomorphism of \mathbb{R}^{n-l-1} . It remains to define the set \mathcal{N}_p with the desired independence property. It is enough to define the *complement* of \mathcal{N}_p . This complement is the projection onto $(T_p S_p)^\perp = T_p M / T_p S_p$ of the set $\gamma'(0) \in T_p M$, where γ ranges through the set of smooth curves $\gamma : [0, 1] \rightarrow \bar{\Omega}$, $\gamma(t) \in \Omega$ for $t > 0$, and $\gamma(0) = p$.

We let then $\sigma_p := \mathcal{N}_p / \mathbb{R}_+$, the set of rays in \mathcal{N}_p , for $p \in \partial\Omega$. Any choice of a metric on $T_p M / T_p S_p \supset \mathcal{N}_p$ will identify σ_p with a subset of the unit sphere of $T_p M / T_p S_p$, which depends however on the metric. In particular, $D\phi_p : \sigma_p \rightarrow \omega_p$ is a diffeomorphism. If p is not in the singular set $\Omega^{(n-2)}$ of Ω , then σ_p consists exactly of one point. The map κ is the projection onto the second component and is one-to-one above Ω and above $\Omega^{(n-1)} \setminus \Omega^{(n-2)} \subset \partial\Omega$.

We now proceed with the induction step. Assume $\Sigma(\omega)$ and $\kappa : \Sigma(\omega) \rightarrow \bar{\omega}$ have been constructed for all curvilinear, stratified polyhedral domains ω of dimension at most $n - 1$. Let Ω be an arbitrary curvilinear, stratified polyhedral domain of dimension n . We define

$$(29) \quad \Sigma(\Omega, M) := \cup_p \{p\} \times \Sigma(\sigma_p, S^{n-\ell(p)-1}), \quad p \in \bar{\Omega}.$$

In particular, if Ω is a bounded domain with smooth boundary, then $\Sigma(\Omega) = \{0\} \times \Omega \simeq \Omega$. This definition is consistent as ω_p is a curvilinear polyhedral domain of dimension at most $n - 1$. Since M in $\Sigma(\Omega, M)$ is most of the time fixed, we will sometimes omit it from the notation. For example, we shall write $\Sigma(\sigma_p) = \Sigma(\sigma_p, S^{k-1})$.

Below, an open embedding will mean a diffeomorphism onto an open subset of the codomain.

Proposition 4.2. *Let $\Omega \subset M$ and $\Omega' \subset M'$ be curvilinear, stratified polyhedral domains and $\Phi : M \rightarrow M'$ be an open embedding such that $\Phi(\Omega)$ is a union of connected components of $\Omega' \cap \Phi(M)$. Then Φ defines a canonical map $\Sigma(\Phi) : \Sigma(\Omega, M) \rightarrow \Sigma(\Omega', M')$ such that*

$$\Sigma(\Phi \circ \Phi') = \Sigma(\Phi) \circ \Sigma(\Phi'),$$

for all open embeddings ϕ and ϕ' for which $\Sigma(\Phi \circ \Phi')$, $\Sigma(\Phi) \circ \Sigma(\Phi')$ are well-defined.

Proof. The proof is by induction. There is nothing to prove for $n = 0$. Let $p \in \overline{\Omega}$. We have that $\Phi(\overline{\Omega}) \subset \overline{\Omega'}$, and hence $\Phi(p) \in \overline{\Omega'}$, as well. Let V'_p be an open neighborhood of $\Phi(p)$ in M' such that there exists a diffeomorphism $\phi'_p : V'_p \rightarrow B^{n-l} \times B^l$ satisfying the condition (17) of the definition of a polyhedral domain (i.e., $\phi'_p(\Omega' \cap V'_p)$ is $\mathbb{R}_+\omega'_p \times B^l$, for some curvilinear polyhedral domain $\omega'_p \subset S^{n-l-1}$). By decreasing V'_p , if necessary, we can assume that $V'_p \subset \Phi(M)$. Then $V'_p \cap \Phi(\Omega)$ is a union of connected components of $V'_p \cap \Omega'$. Therefore ω'_p is a union of connected components of $\Phi(\omega_p)$, where $\omega_p \subset S^{n-l-1}$ is associated to $p \in \overline{\Omega}$ in the same way as ω'_p was associated to $\Phi(p) \in \overline{\Omega'}$. The induction hypothesis then gives rise to a canonical, injective map $\Sigma(\omega_p, S^{n-l-1}) \rightarrow \Sigma(\omega'_p, S^{n-l-1})$. The map $\Sigma(\phi)$ is obtained by combining these different maps.

The functoriality (i.e., the relation $\Sigma(\phi \circ \phi') = \Sigma(\phi) \circ \Sigma(\phi')$) is proved similarly by induction. \square

Here is a corollary of the above proof.

Corollary 4.3. *If $\Omega = \Omega' \cup \Omega''$ is the disjoint union of two open sets, then the inclusions $\Sigma(\Omega', M) \subset \Sigma(\Omega, M)$ and $\Sigma(\Omega'', M) \subset \Sigma(\Omega, M)$ defined in Proposition 4.2 realize $\Sigma(\Omega, M) = \Sigma(\Omega', M) \cup \Sigma(\Omega'', M)$, where the union is a disjoint union.*

Proof. We use the same argument as in the proof of Proposition 4.2. \square

The desingularization has a simple behavior with respect to products.

Lemma 4.4. *We have a canonical identification*

$$\Sigma(M' \times \Omega, M' \times M) = M' \times \Sigma(\Omega, M),$$

for any smooth manifolds M and M' and any curvilinear polyhedral domain $\Omega \subset M$.

Proof. Since M' is smooth, we can choose the structural local diffeomorphism $\phi_{(p,q)}$ in $M' \times \Omega$ to be given by $\phi_p \times \psi_q$, where ψ_p is a local coordinate chart defined in a neighborhood of $p \in M'$ and ϕ_q is the local diffeomorphism of a neighborhood of q in Ω . Indeed, then

$$(30) \quad \Sigma(M' \times \Omega) := \cup_{p,q} \{(p, q)\} \times \Sigma(\sigma_{(p,q)}) = \cup_{p,q} \{(p, q)\} \times \Sigma(\sigma_q) = M' \times \Sigma(\Omega),$$

where $q \in \overline{\Omega}$ and $p \in M'$. Consequently, there is a canonical bijection $\sigma_{(p,q)} \simeq \sigma_q$ for any $q \in \overline{\Omega}$ and any $p \in M'$ (so (p, q) is in the closure of $M' \times \Omega$ in $M' \times M$). \square

It remains to define the topology and differentiable structure on $\Sigma(\Omega)$. These definitions will again be canonical if we require that the map of the above lemma, as well as the maps κ and $\Sigma(\phi)$, be differentiable, for any open embedding ϕ .

Let $V_p \subset M$ and ϕ_p be as in Equation (18). By Proposition 4.2, we may assume that ϕ_p is the identity, so that $p = 0$, $V_p = B^{n-l} \times B^l$, and $V_p \cap \Omega = I\omega_p \times B^l$, with $I = (0, 1)$. Let $(\Sigma(\omega_p), \kappa'_p)$ be the canonical desingularization of ω_p in S^{n-l-1} . We shall need the following lemma.

Lemma 4.5. *We have a canonical identification*

$$\Sigma(V_p \cap \Omega, M) = [0, 1) \times \Sigma(\omega_p, S^{n-l-1}) \times B^l$$

and the desingularization map

$$(31) \quad \kappa_p : [0, 1) \times \Sigma(\omega_p) \times B^l \rightarrow \overline{V_p \cap \Omega} \subset B^{n-l} \times B^l$$

is given by $\kappa_p(r, x', y) = (r\kappa'_p(x'), y)$.

Proof. We shall replace p with 0 below, since we have assumed $p = 0$. This will simplify the notation. The closure of $V_0 \cap \Omega$ in $V_0 = V_p$ is the disjoint union $\{0\} \times B^l \cup \overline{I\omega_0} \times B^l$. Accordingly we decompose $\Sigma(V_0 \cap \Omega, M)$ into two disjoint sets, corresponding to this splitting of the closure of $V_0 \cap \Omega$. Recall that by definition $\Sigma(V_0 \cap \Omega, M)$ is the union $\cup_{p \in \overline{V_0 \cap \Omega}} \{p\} \times \Sigma(\sigma_p, S^{n-l-1})$. Using also Lemma 4.4, we then obtain

$$\begin{aligned} \Sigma(V_0 \cap \Omega, M) &= \Sigma(V_0 \cap \Omega, M \setminus (\{0\} \times B^l)) \cup \bigcup_{q \in B^l} \{(0, q)\} \times \Sigma(\omega_0) \\ &= \Sigma((0, 1) \times \omega_0 \times B^l, (0, 1) \times S^{n-l-1} \times B^l) \cup \bigcup_{q \in B^l} \{(0, q)\} \times \Sigma(\omega_0) \\ &= (0, 1) \times \Sigma(\omega_0) \times B^l \cup \{0\} \times \Sigma(\omega_0) \times B^l \\ &= [0, 1) \times \Sigma(\omega_0) \times B^l. \end{aligned}$$

The formula for κ_0 follows from the definition. □

Since $\Sigma(\Omega, M)$ is the union of all the sets $\Sigma(V_p \cap \Omega, M)$, with V_p in the covering above, we can define the topology and smooth structure on $\Sigma(\Omega, M)$ as follows.

Definition 4.6. Let $\phi_p : V_p \rightarrow B^{n-l} \times B^l$ and ω_p be as in Definition 2.1. The topology and smooth structure on $\Sigma(\Omega, M)$ are such that the induced structure on $\Sigma(V_p \cap \Omega, M)$ is the same as the one obtained from the canonical identification $\Sigma(V_p \cap \Omega, M) = [0, 1) \times \Sigma(\omega_p) \times B^l$ of Lemma 4.5.

We need to prove that the transition functions are smooth. This follows from the fact that the maps ϕ_p are diffeomorphisms.

We have therefore completed the definition of the desingularization $\Sigma(\Omega, M)$ and of its smooth structure.

4.2. The distance to singular boundary points. We continue with a study of the geometric and, especially, metric properties of $\Sigma(\Omega, M)$. Since M will be fixed from now on, we shall write $\Sigma(\Omega) = \Sigma(\Omega, M)$. We first argue that $\Sigma(\Omega)$ has embedded faces and hence that every hyperface of $\Sigma(\Omega)$ has a defining function.

Let F_0 be an open hyperface of a manifold with corners \mathfrak{M} . Then F_0 is a manifold of dimension $n - 1$. Its closure F , in general, will not necessarily be a manifold, because it may have self-intersections. (A typical example is the boundary of a curvilinear polygonal domain with only one vertex, the “tear drop domain.”) By induction, however, it follows that $F \cap V_p$ will be a manifold, for any p . In particular, we obtain that all (closed) faces of $\Sigma(\Omega)$ are embedded submanifolds of $\Sigma(\Omega)$. Let H be a hyperface of $\Sigma(\Omega)$, since H is an embedded submanifold of codimension 1, there will exist a function $x_H > 0$ on Ω , $H = \{x_H = 0\}$, and $dx_H \neq 0$ on H . A function x_H with this property is called a *defining function of H* , see Melrose’s book [42].

One of the main reasons for introducing the desingularization space $\Sigma(\Omega)$ is the following result.

Proposition 4.7. *Let Ω be a bounded, curvilinear, stratified polyhedral domain and g_1 and g_2 be two smooth Riemannian metrics on M . Fix k and assume $\Omega^{(k)} \neq \emptyset$. Let $f_j(x)$ be the modified distance from $x \in \overline{\Omega}$ to the set $\Omega^{(k)}$ in the metric g_j ,*

computed within $\bar{\Omega}$. Then the quotient f_2/f_1 extends to a continuous function on $\Sigma(\Omega)$.

Proof. It is enough to prove the given property in the neighborhood of every point $p \in \bar{\Omega}$. So let us fix $p \in \bar{\Omega}$. By replacing V_p with a smaller neighborhood of p , if necessary, we can also assume that $g_2(\xi) \leq Cg_1(\xi)$, which implies that $f_2 \leq Cf_1$, and hence that f_2/f_1 is bounded.

We shall prove the statement by induction on n . In the case $n = 1$, the only possibility is that $k = 0$, or otherwise $\Omega^{(k)} = \emptyset$. Then $f(x)$ is the distance to the vertices of Ω . Recall that Ω is a disjoint union of open intervals in this case, so that we can reduce to consider a single interval. If say $\Omega = [a, b]$, then close to a , $f_j(x) = a_j(x - a)$, with a_j smooth near a and $a_j(a) \neq 0$. The same situation holds at b . This proves our result in the case $n = 1$. We now proceed with the induction step.

The function f_1/f_2 is clearly continuous on the open set Ω . Fix $p \in \partial\Omega$. We shall construct an open neighborhood U_p of p in $\bar{\Omega}$ such that f_1/f_2 extends to a continuous function on $\kappa^{-1}(U_p)$. Let V_p be as in the definition of polyhedral domains (Definition 2.1). We shall identify $V_p \cap \Omega$ with $I\omega_p \times B^l$ using the diffeomorphism ϕ_p of Equation (18). If $l > k$, that is, $p \in \Omega^{(l)} \setminus \Omega^{(k)}$, then both f_1 and f_2 extend to continuous, non-vanishing functions on $V_p \cap \bar{\Omega}$, which can be lifted to continuous, non-vanishing functions on $\kappa^{-1}(V_p \cap \Omega)$. We shall assume hence that $k \geq l$.

On a smaller neighborhood $V' \subset V_p$ if necessary, we can arrange that the function f_1 gives the distance to $V_p^{(k)}$, that is, that the point of $\Omega^{(k)}$ closest to $x \in V' \cap \Omega$ is, in fact, in V_p . By decreasing V' even further, we can further arrange that the same holds for f_2 . Then we shall take $U_p := V'$.

To prove that f_2/f_1 extends to a continuous function on $\kappa^{-1}(U_p)$, it is enough to do that in the case $\Omega = V_p \cap \Omega$, because the quotient f_2/f_1 does not change on $U_p \cap \Omega$ if we replace Ω with $V_p \cap \Omega$, as explained in the paragraph above. We can also assume that g_2 is the standard Euclidean metric, but then we have to prove that f_1/f_2 extends to a *nowhere vanishing* continuous function on $\Sigma(\Omega)$. (Using also Proposition 4.2, we have reduced to the case $\Omega = I\omega_p \times B^l \subset \mathbb{R}^n$, $I = (0, t)$.)

The scaling property of the Euclidean metric and our assumption that $k \geq l$ imply that

$$f_2(rx', x'') = rf_2(x', x''),$$

for any $r \in [0, 1]$. Let g_0 be a constant metric on \mathbb{R}^n that coincides with g_1 at the origin.

Let f_0 be associated to g_0 in the same way as f_j is associated to g_j , for $j = 1, 2$, i.e., $f_0(x) = \text{dist}(x, \Omega^{(k)})$ using the metric g_0 . We then have similarly $f_0(rx', x'') = rf_0(x', x'')$, so that the quotient $f_0(rx', x'')/f_1(rx', x'')$ does not depend on r . We can therefore fix $r = 1$. Consequently, we can work with the lower dimensional polyhedral domain $\omega := \omega_p \times B^l$ instead of $\Omega = I\omega_p \times B^l$, and prove that f_0/f_1 extends by continuity to $\Sigma(\omega)$. It remains to see that we can use induction to prove the existence of this extension. Since ω is by construction a stratified polyhedron, we denote by $\omega^{(k)} = \omega^{(k)} \times B^l$ $k < n$, its associated stratification, where we set $\omega_p^{(k-l-1)} = \emptyset$ if $k-l-1 < 0$ as before. Let f'_1 be the distance function to $\omega^{(k-1)}$ on ω (i.e., computed *within* $\bar{\omega}$, with respect to the metric induced by g_1 , as in the statement of Proposition 4.7). We let $f'_1 = 1$ if $\omega_p^{(k-l-1)} = \emptyset$.

We define f'_0 similarly. The inductive hypothesis guarantees that f'_0/f'_1 extends to a continuous function on $\Sigma(\omega) = \Sigma(\omega_p) \times B^l$. On the other hand, it is easy to see that both f_1/f'_1 and f'_1/f_1 extend to continuous functions on $\bar{\omega}$ if we set them to be equal to 1 on $\omega^{(k-1)}$. The same is true of f_0/f'_0 and f'_0/f_0 . Putting all these estimates together, it follows that

$$f_0/f_1 = (f_0/f'_0)(f'_0/f'_1)(f'_1/f_1)$$

extends to a continuous, nowhere vanishing function on $\Sigma(\omega)$.

Let us tackle now the case g_2 arbitrary. Let f_0 be defined as before. We then have that $f_2(rx', x'') = rf_0(x', x'') + r^2h(rx', x'')$, with h a continuous function on $\Sigma(V_p \times \Omega)$ that vanishes on $\Omega^{(k)}$. Then

$$\frac{f_2}{f_1} = \frac{f_0}{f_1} + r \frac{h(rx', x'')}{f_1(x', x'')}.$$

The function f_0/f_1 was already shown to extend by continuity to $\Sigma(\Omega)$. The same argument as above shows that h/f_1 extends by continuity to a nowhere vanishing function on

$$[\epsilon, 1) \times \Sigma(\omega_p) \times B^l \subset (0, 1) \times \Sigma(\omega_p) \times B^l =: \Sigma(\Omega).$$

The continuity of f_2/f_1 then follows from the boundedness of f_2/f_1 .

The resulting function does not vanish at $r = 0$, because it is equal to f_0/f_1 there. It was already proved that it does not vanish for $\epsilon > 0$. The proof is complete. \square

We shall need also the following corollary of the above proof.

Corollary 4.8. *Identify $V_p \cap \Omega = \Omega$ with $I\omega_p \times B^l$, $I = (0, a)$, $l = \ell(p)$, using the diffeomorphism ϕ_p given in Definition 2.1. Let g be a smooth metric on V_p , and let $f(x)$ be the distance from x to $\Omega^{(k)}$, $k \geq l$, $f'(x', x'')$ be the distance from $(x', x'') \in \omega := \omega_p \times B^l$ to $\omega^{(k-1)}$ (within $\bar{\omega}$, as in Proposition 4.7) if $\omega^{(k-1)} \neq \emptyset$, and $f'(x', x'') = 1$ otherwise. Assume ω_p is connected. Then*

$$f(rx', x'')/rf'(x', x'')$$

extends to a continuous, nowhere vanishing function on $\Sigma(\Omega) = [0, a) \times \Sigma(\omega_p) \times B^l$.

Proof. Assume first that $\omega^{(k-1)} \neq \emptyset$, where $\omega^{(k)}$ is defined as in Proposition 4.7. Let f_0 and f'_0 be defined in the same way f and f' were defined, but replacing g with a constant metric g_0 . Then the proof of Proposition 4.7 gives that $f_0(rx', x'')/rf'_0(x', x'')$ is independent of r . Hence $f_0(rx', x'')/rf'_0(x', x'')$ extends to a continuous, nowhere vanishing function on $\Sigma(\Omega)$, as it was shown in the proof of Proposition 4.7. Then

$$\frac{f(rx', x'')}{rf'(x', x'')} = \frac{f(rx', x'')}{f_0(rx', x'')} \times \frac{f_0(rx', x'')}{rf'_0(rx', x'')} \times \frac{f'_0(x', x'')}{f'(x', x'')}.$$

We have just argued that the middle quotient in the above product extends to a continuous function on $\Sigma(\Omega)$. The other two quotients also extend to continuous functions on $\Sigma(\Omega)$, by Proposition 4.7 applied to Ω and ω .

Let us assume now that $\omega^{(k-1)} = \emptyset$. Then the same proof applies, given that $f'_0/f' = 1$ clearly extends to a continuous function on $\Sigma(\Omega)$. \square

4.3. The weight function r_Ω . Recall that $\eta_{n-2}(x)$, given in Definition 2.5, denotes the distance from $x \in \bar{\Omega}$ to the singular set $\Omega^{(n-2)}$.

The main goal of this subsection is to define on any curvilinear polyhedral domain Ω a function

$$r_\Omega : \bar{\Omega} \rightarrow [0, \infty)$$

that lifts to a *smooth* function on $\Sigma(\Omega)$ and is *equivalent* to η_{n-2} . (Additional properties of r_Ω will be established later on.) This will lead to a definition of the Sobolev spaces $\mathcal{K}_a^m(\Omega)$ as weighted Sobolev spaces on Lie manifolds with boundary, Proposition 5.8. We again proceed by induction on n .

We define $r_\Omega = 1$ if $n \leq 1$ (recall $\Omega^{(n-2)} = \emptyset$ if $n < 2$) or if $\Omega^{(n-2)} = \emptyset$, that is, Ω is a smooth manifold, possibly with boundary.

Assume now that a function r_ω was defined for all curvilinear polyhedral domains ω of dimension $< n$ and let us define it for a given bounded n -dimensional curvilinear polyhedral domain Ω .

We localize first to a neighborhood of a generic point $p \in \partial\Omega$ and then use a partition of unity argument. We recall that by definition there exists a neighborhood V_p of p in M , a diffeomorphism $\phi_p : V_p \rightarrow B^{n-l} \times B^l$, for some $0 \leq l = \ell(p) \leq n-1$, and a polyhedral domain $\omega_p \subset S^{n-l-1}$ such that $\phi_p(V_p \cap \Omega) = I \omega_p \times B^l$, $I = (0, \epsilon)$, see Condition (17)). Therefore, we can assume that ϕ_p is the identity map and replace in what follows V_p with $\phi_p^{-1}(\frac{1}{2}B^{n-l} \times \frac{1}{2}B^l)$. Since r_Ω is already defined equal to 1 if $p \in \Omega^{(n-1)} \setminus \Omega^{(n-2)}$, we restrict $n-l-1 \geq 1$ above. Let r_{ω_p} the function associated to the curvilinear polyhedral domain ω_p . Then we define

$$(32) \quad r_{V_p}(rx', x'') := rr_{\omega_p}(x'), \quad (rx', x'') \in \Omega \subset V_p,$$

if $x' \in \omega_p$, $x'' \in B^l$, and $1 \leq l = \ell(p) \leq n-2$. Following our usual procedures, we set $r_{V_p}(rx') = rr_{\omega_p}(x')$ if $l = 0$.

We consider next a locally finite covering of $\bar{\Omega}$ with open sets U_α of one of the three following forms

- (i) $U_\alpha \subset \bar{U}_\alpha \subset \Omega$ with ∂U_α smooth;
- (ii) $U_\alpha = V_p$ with $\ell(p) = n-1$ (i.e., p is not in the singular set of $\bar{\Omega}$); or
- (iii) such that for any $x \in U_\alpha \cap \Omega$, the point of $\Omega^{(n-2)}$ closest to x is in V_p with $\ell(p) \leq n-2$, and

$$(33) \quad p \in U_\alpha \subset \bar{U}_\alpha \subset V_p.$$

A condition similar to (iii) above was already used in the proof of Proposition 4.7. The conditions (i) and (ii) above correspond exactly to the case when $(\partial U_\alpha \cap \partial\Omega)$ is smooth (this includes the case when $(\partial U_\alpha \cap \partial\Omega)$ is empty).

We then set

$$(34) \quad r_\alpha = \begin{cases} 1 & \text{if } (\partial U_\alpha \cap \partial\Omega) \text{ is smooth} \\ r_{V_p} & \text{if } U_\alpha \text{ is as in (33)}. \end{cases}$$

and define

$$(35) \quad r_\Omega = \sum_\alpha \varphi_\alpha r_\alpha,$$

where φ_α is a smooth partition of unity subordinated to U_α . If Ω is not bounded, we define instead:

$$(36) \quad r_\Omega = \chi\left(\sum_\alpha \varphi_\alpha r_\alpha\right),$$

where χ is defined as in (22). We notice that the definition of r_Ω is not canonical, because it depends on a choice of a covering $\{U_\alpha\}$ of $\bar{\Omega}$ as above and a choice of a subordinated partition of unity.

Proposition 4.9. *Let Ω be a curvilinear, stratified polyhedral domain of dimension $n \geq 2$. Then r_Ω defined in Equation (35) (or (36)) is continuous on $\bar{\Omega}$ and $r_\Omega \circ \kappa$ is smooth on $\Sigma(\Omega)$. Moreover, η_{n-2}/r_Ω extends to a continuous, nowhere vanishing function on $\Sigma(\Omega)$ and r_α/r_Ω extends to a smooth function on $\Sigma(V_p \cap \Omega)$.*

Proof. Let $\eta_{-1} := 1$ for the inductive step. We shall prove the statement on η_{n-2}/r_Ω by induction on $n \geq 1$. Since $r_\Omega = 1$ for polyhedral domains of dimension $n = 1$, the result is obviously true for $n = 1$. We now proceed with the inductive step.

We shall use the above results, in particular, Proposition 4.7, for $k = n - 2 \geq 0$. Thus $f = \eta_{n-2}$ in the notation of Proposition 4.7. Let $f_\alpha(x)$ be the distance from $x \in V_p$ to $V_p \cap \Omega^{(n-2)}$, if $U_\alpha \subset V_p$ is as in Equation (33) (so $\ell(p) \leq n - 2$ in this case). Thus $f_\alpha = f$ on $U_\alpha \cap \Omega$, by the construction of U_α . We identify once again $V_p \cap \Omega$ with $(0, \epsilon) \omega_p \times B^l$, $l = \ell(p)$, using the diffeomorphism ϕ_p , and set again $\omega := \omega_p \times B^l$. Also, for any $x \in \omega$, let $f'_\alpha(x)$ be the distance from x to the singular set $\omega^{(n-l-2)}$ of ω if it is not empty, $f'_\alpha(x) = 1$ otherwise. Let r_α be as in the definition of r_Ω , Equation (35). Then

$$\frac{f_\alpha(rx', x'')}{r_\alpha(rx', x'')} = \frac{f_\alpha(rx', x'')}{rf'_\alpha(x', x'')} \frac{f_\alpha(x', x'')}{r_{\omega_p}(x', x'')}, \quad \text{for } (rx', x'') \in V_p \cap \Omega.$$

The quotient $f_\alpha(rx', x'')/rf'_\alpha(x', x'')$ extends to a continuous, nowhere vanishing function on $\Sigma(V_p \cap \Omega)$, by Corollary 4.8. By the induction hypothesis, the quotient $f_\alpha(x', x'')/r_{\omega_p}(x', x'')$ also extends to a continuous, nowhere vanishing function on $\Sigma(\omega) = \Sigma(\omega_p) \times B^l$. Since this quotient is independent of r , it also extends to a continuous, nowhere vanishing function on $\Sigma(V_p \cap \Omega)$. Hence f_α/r_α extends to a continuous, nowhere vanishing function on $\Sigma(V_p)$. Therefore

$$r/f = \sum_\alpha \varphi_\alpha r_\alpha / f = \sum_\alpha \varphi_\alpha r_\alpha / f_\alpha$$

extends to a continuous function on $\Sigma(\Omega)$.

The quotient r/f is immediately seen to be non-zero everywhere, from the definition. Hence f/r also extends to a continuous function on $\Sigma(\Omega)$.

We have already noticed that r_α/f extends to a continuous, nowhere vanishing function on $\Sigma(V_p)$. Hence $r_\alpha/r_\Omega = (r_\alpha/f)(f/r_\Omega)$ extends to a continuous, nowhere vanishing function on $\Sigma(V_p \cap \Omega)$. Since both r_α and r_Ω are smooth on $\Sigma(V_p \cap \Omega)$ and the set of zeroes of r_Ω is the union of transversal manifolds on which r_Ω has simple zeroes, it follows that r_α/r_Ω extends to a smooth function on $\Sigma(V_p)$. Since $\bar{U}_\alpha \subset V_p$ is compact, it follows from a compactness argument that r_α and r are equivalent on U_α . The proof is complete. \square

We can now prove the following result, which will be used in the proof of Theorem 6.4.

Proposition 4.10. *Let Ω be a bounded, curvilinear, stratified polyhedral domain. Suppose r_Ω, r'_Ω are two functions on $\bar{\Omega}$ defined by formula (35) (or (36)) with possibly different choices of open covering $\{U_\alpha\}$, subordinate partition $\{\varphi_\alpha\}$, and*

diffeomorphisms ϕ_p . Then r'_Ω/r_Ω extends to a smooth, nowhere vanishing function on $\Sigma(\Omega)$.

Proof. We know from Proposition 4.9, that f/r'_Ω and f/r_Ω extend to continuous, nowhere vanishing functions on $\Sigma(\Omega)$. Hence r'_Ω/r_Ω extends to a continuous, nowhere vanishing function on $\Sigma(\Omega)$. Since both r'_Ω and r_Ω are smooth functions on $\Sigma(\Omega)$ and the set of zeroes of r_Ω is a union of transverse manifolds, each a set of simple zeroes of r_Ω , it follows that the quotient r'_Ω/r_Ω is smooth on $\Sigma(\Omega)$. \square

We obtain the following corollary. Let $H \subset \Sigma(\Omega)$ be a hyperface (*i.e.*, face of maximal dimension) of $\Sigma(\Omega)$. Recall that a *defining function* of H is a smooth function $x_H \geq 0$ defined on $\Sigma(\Omega)$, such that $H = \{x = 0\}$ and $dx_H \neq 0$ on H . All the faces of $\Sigma(\Omega)$ are closed subsets of $\Sigma(\Omega)$, by definition. We have already noticed that any face of $\Sigma(\Omega)$ has a defining function. We then have the following corollary.

Corollary 4.11. *Let $\eta = \prod_H x_H$, where H ranges through the set of hyperfaces of $\Sigma(\Omega)$ that do not intersect $\partial\Omega \setminus \Omega^{(n-2)}$. Then η/r_Ω extends to a smooth, nowhere vanishing function on $\Sigma(\Omega)$.*

Proof. This is a local statement that can be checked by induction, as in the previous proofs. \square

In particular, since the function r_Ω is anyway determined only up to a factor of $h \in \mathcal{C}^\infty(\Sigma(\Omega))$, $h \neq 0$, we obtain that we could take $r_\Omega = \prod_H x_H$, where H ranges through the set of hyperfaces of $\Sigma(\Omega)$ that do not intersect $\partial\Omega \setminus \Omega^{(n-2)}$. The function r_Ω , for various versions of the set Ω , will play an important role in the inductive definition of the structural Lie algebra of vector fields $\mathcal{V}(\Omega)$ on $\Sigma(\Omega)$, which we address next.

4.4. The structural Lie algebra of vector fields. We now proceed to define by induction a canonical Lie algebra of vector fields $\mathcal{V}(\Omega)$ on $\Sigma(\Omega)$, for Ω a curvilinear, stratified polyhedral domain of dimension $n \geq 1$. In view of Corollary 4.3, we can assume that Ω is connected. We denote by

$$\mathcal{X}(M) := \Gamma(M; TM)$$

the space of vector fields on a manifold M .

We let

$$(37) \quad \mathcal{V}(\Omega) = \mathcal{X}(\bar{\Omega}) = \mathcal{X}(\Sigma(\Omega)) \quad \text{if } n = 1.$$

In other words, there is no restriction on the vector fields $X \in \mathcal{V}(\Omega)$, if Ω has dimension one.

Assume now that the Lie algebra of vector fields $\mathcal{V}(\omega)$ has been defined on $\Sigma(\omega)$ for all curvilinear polyhedral domains ω of dimension $1 \leq k \leq n-1$ and let us define $\mathcal{V}(\Omega)$ for a curvilinear polyhedral domain of dimension n . We fix $p \in \partial\Omega$ and let V_p and ϕ_p be as in Definition 2.1, as usual. We identify $V_p \cap \Omega$ with $(0, 1)\omega_p \times B^l$ using ϕ_p . Assume $1 \leq \ell(p) \leq n-2$, so that in particular ω_p is a curvilinear polyhedral domain of dimension ≥ 1 . We notice that

$$T([0, 1) \times \Sigma(\omega_p) \times B^l) = T([0, 1)) \times T\Sigma(\omega_p) \times TB^l$$

and hence

$$\mathcal{X}([0, 1) \times \Sigma(\omega_p) \times B^l) = \mathcal{X}([0, 1)) \times \mathcal{X}(\Sigma(\omega_p)) \times \mathcal{X}(B^l).$$

Then we define

$$(38) \quad \mathcal{V}(V_p \cap \Omega) = \{X = (X_1, X_2, X_3) \in \mathcal{X}([0, 1]) \times \mathcal{X}(\Sigma(\omega_p)) \times \mathcal{X}(B^l)\}$$

where X_1 , X_2 , and X_3 are required to satisfy the following two conditions:

$$(39) \quad Y_1 := r_\Omega^{-1}X_1 \text{ and } Y_3 := r_\Omega^{-1}X_3 \text{ are smooth}$$

and

$$(40) \quad X_2(t, x', x'') \in \mathcal{V}(\{t\} \times \omega_p \times \{x''\}) = \mathcal{V}(\omega_p), \text{ for any fixed } t, y.$$

In Condition (39) above, smooth means, smooth *including at* $r = 0$. If $\ell(p) = 0$, then we just drop the component X_3 , but keep the same conditions on X_1 and X_2 . By Proposition 4.10, the definition of $\mathcal{V}(V_p \cap \Omega)$ is independent of the choice of r_Ω . All vector fields are assumed to be smooth.

Finally, we define $\mathcal{V}(\Omega)$ to consist of the vector fields $X \in \mathcal{X}(\Sigma(\Omega))$ such that $X|_{V_p \cap \Omega} \in \mathcal{V}(V_p \cap \Omega)$ for all $p \in \Omega^{(n-2)}$. In particular, only the smoothness condition is imposed on our vector fields at the smooth points of $\partial\Omega$. Note that the vector fields in $\mathcal{V}(\Omega)$ may not extend to the closure $\bar{\Omega}$, in general. This was seen in Example 2.10.

4.5. Lie manifolds with boundary. We now proceed to show that the pair $(\Sigma(\Omega), \mathcal{V}(\Omega))$ defines a Lie manifold with boundary, introduced in [1], and the construction of which was recalled in Definition 3.5.

We first establish some lemmata.

Lemma 4.12. *Let $X \in \mathcal{X}(\Sigma(\Omega))$ be such that $X = 0$ in a neighborhood of the boundary of $\Sigma(\Omega)$. Then $X \in \mathcal{V}(\Omega)$.*

Proof. The result follows immediately by induction from the definition of $\mathcal{V}(\Omega)$. \square

We also get the following simple fact.

Lemma 4.13. *If $f : \Sigma(\Omega) \rightarrow \mathbb{C}$ is a smooth function and $X \in \mathcal{V}(\Omega)$, then $X(f)$ is a smooth function on $\Sigma(\Omega)$ and $fX \in \mathcal{V}$.*

Proof. The vector field X is smooth on $\Sigma(\Omega)$, hence $X(f)$ is smooth on $\Sigma(\Omega)$. The second statement is local, so it is enough to check it on Ω and on each V_p , on which it is as a direct consequence of the definition and induction. \square

Lemma 4.14. *For any $X \in \mathcal{V}(\Omega)$ and any continuous function $f : \bar{\Omega} \rightarrow \mathbb{C}$ such that $f \circ \kappa$ is smooth on $\Sigma(\Omega)$, we have*

$$X(f) = \tilde{f} r_\Omega,$$

where \tilde{f} is a smooth function on $\Sigma(\Omega)$. In particular, $X(r_\Omega) = f_X r_\Omega$, where f_X is a smooth function on $\Sigma(\Omega)$.

Proof. This is a local statement that can be checked by induction in any neighborhood V_p , using the definition, as follows. Let us use the notation of Equation (38), and (39) and write

$$X = (X_1, 0, 0) + (0, X_2, 0) + (0, 0, X_3).$$

We shall write, with abuse of notation, $X_1 = (X_1, 0, 0)$. Define X_2 and X_3 similarly. It is enough to check that $X_j f(rx', x'')$ is of the indicated form, for $j = 1, 2, 3$. We have $X_1 = r_\Omega Y_1$ and $X_3 = r_\Omega Y_3$, where Y_1 and Y_3 are smooth (in appropriate

spaces), by Equation (39). This observation proves our lemma if $X = X_1$ or $X = X_3$. If $X = X_2$, then we have

$$(41) \quad (Xf)(r, x', x'') = X_2(f(rx', x'')) = r_{\omega_p} f_1(r, x', x''),$$

with f_1 a smooth function on $\Sigma(V_p \cap \Omega) = [0, \epsilon) \times \Sigma(\omega_p) \times \mathbb{R}^l$, by the induction hypothesis. Moreover, given that by assumption (40) $\kappa_* X$ is a vector field tangent to the sphere S^{n-l-1} , we see that $Xf(0, x', x'') = 0$. Therefore $Xf = rr_{\omega_p} \tilde{f}$, for some smooth function \tilde{f} on $\Sigma(V_p \cap \Omega)$. Let us denote $r_\alpha = rr_{\omega_p}$, as in Equation (34) and in Proposition 4.9. Proposition 4.9 gives that r_α/r_Ω is smooth on its domain of definition. Hence $Xf = r_\alpha f_1 = r_\Omega (r_\alpha/r_\Omega) f_1 = r_\Omega \tilde{f}$, with \tilde{f} smooth on each $\Sigma(V_p \cap \Omega)$. Hence \tilde{f} is smooth on $\Sigma(\Omega)$. \square

We next characterize which vector fields on Ω are restrictions of vector fields on $\mathcal{V}(\Omega)$. We begin by showing that the restriction property is local.

Lemma 4.15. *Let Y be a vector field on Ω with the property that every point $p \in \bar{\Omega}$ has a neighborhood U_p in M such that $Y = X_U$ on $U \cap \Omega$, for some $X_U \in \mathcal{V}(\Omega)$. Then there exists $X \in \mathcal{V}(\Omega)$ such that Y is the restriction of X to Ω .*

Proof. Let us cover $\bar{\Omega}$ with a locally finite family of sets U_p , $p \in B \subset \bar{\Omega}$. Let ψ_p , $p \in B$, be a subordinated partition of unity.

We claim that $X = \sum_{p \in B} \psi_p X_{U_p} \in \mathcal{V}(\Omega)$ (by Lemma 4.13) satisfies $X(x) = Y(x)$, $x \in \Omega$. Indeed, $X(x) = \sum_{p \in B} \psi_p(x) X_{U_p}(x) = (\sum \psi_p(x)) Y(x) = Y(x)$. \square

We can now prove the following lemma.

Lemma 4.16. *Let Y be a smooth vector field on $\bar{\Omega}$. Then $r_\Omega Y$ is the restriction to $\Omega \subset \Sigma(\Omega)$ of a vector field X in $\mathcal{V}(\Omega)$.*

Proof. By Lemma 4.15, it is enough to check this statement on a neighborhood V_p of some $p \in \partial\Omega$. We shall proceed by induction. Since the desingularization and the definition of r_Ω are covariant with respect to diffeomorphism (that respect the stratification of Ω), we can assume that $V_p = B^{n-l} \times B^l$ and that $V_p \cap \Omega \simeq (0, 1) \omega_p \times B^l$. Assume first that $Y = \partial_j$ is a constant vector field on V_p . Let $\alpha_t(x', x'') = (tx', x'')$. Then $D\alpha_t(\partial_j) = t\partial_j$. Therefore,

$$(42) \quad D\alpha_t(X) = X,$$

where $X = r_\Omega \partial_j$, where r_Ω can be taken, on V_p , to be given by rr_{ω_p} . Let us decompose $\partial_j = (Y_1, Y_2, Y_3)$ on V_p using the notation of Equation (38). Then Y_3 is constant. In fact, either $Y_3 = \partial_j$ or $Y_3 = 0$. In any instance, if we write $X = (X_1, X_2, X_3)$, then $X_3 = r_\Omega Y_3$ satisfies the condition of Equation (39). The relation (42) gives that $Y_1(r, x', x'') = a_1(x') \partial_r$ and $Y_2(r, x', x'') = r^{-1} Z(x')$, with a_1 a smooth function and Z a smooth vector field on $\bar{\omega}_p$. Clearly $X_1 = r_\Omega Y_1$ will satisfy the conditions of Equation (39). The induction hypothesis then gives that $X_2(r, x', x'') = r_\Omega Y_2(r, x', x'') = r_{\omega_p}(x') Z(x')$ is the restriction to $V_p \cap \Omega$ of a smooth vector field in $\mathcal{V}(V_p \cap \Omega)$. (This vector field depends only on the second factor in $\Sigma(V_p \cap \Omega) = [0, 1) \times \omega_p \times B^l$.) \square

We now identify a canonical metric on the vector fields \mathcal{V} . Recall that the concept of local basis of a space of vector fields was defined in Definition 3.1.

Proposition 4.17. *Let us fix a metric h on $M \supset \Omega$. Let $q \in \Sigma(\Omega)$ be arbitrary. Then there exists a neighborhood U of q in $\Sigma(\Omega)$ and $X_1, X_2, \dots, X_n \in \mathcal{V}(\Omega)$ that form a local basis of $\mathcal{V}(\Omega)$ on U and satisfy*

$$h(X_j, X_k) = r_\Omega^2 \delta_{jk}.$$

In other words, the vectors X_1, X_2, \dots, X_n form an orthonormal system on $\Omega \cap U$ for the metric $r_\Omega^{-2}h$. A local basis X_1, X_2, \dots, X_n with this property will be called a *local orthonormal basis of $\mathcal{V}(\Omega)$ over U* .

Proof. If $q \in \Omega \subset \Sigma(\Omega)$, the result follows from Lemma 4.12. Let $p = \kappa(q)$. We shall hence assume that $p \in \partial\Omega$. This is again a local statement in $p \in \partial\Omega$. We can therefore proceed by induction. If the dimension n of Ω is 1, then there is nothing to prove because $r_\Omega = 1$ in this case.

Once again, we let $\phi_p : V_p \rightarrow B^{n-l} \times B^l$ and ω_p be as in Definition 2.1. We can assume that ϕ_p is the identity map. If we can prove the result for the function $r = r_\Omega$, then we can prove it for the function $r' = f'r$, where $f', 1/f' \in C^\infty(\Sigma(\Omega))$, simply by replacing X_j with $f'X_j$. By Proposition 4.9, we can therefore assume that $r_\Omega = rr_{\omega_p}$ on $V_p \cap \Omega$. Let $q = (0, x', x'') \in [0, 1) \times \Sigma(\omega_p) \times B^l$.

Let h_0 be the standard metric on V_p . For the induction hypothesis, we shall need that the metric h_0 is given by

$$(43) \quad h_0(r, x', x'') = (dr)^2 + r^2(dx')^2 + (dx'')^2$$

on $\Omega \cap V_p = (0, 1)\omega_p \times B^l$. Here $(dx')^2$ denotes the metric on ω_p induced by the Euclidean metric on the sphere S^{n-l-1} . In other words, if $X = (X_1, X_2, X_3)$ is a vector field on $V_p \cap \Omega$, written using the product decomposition explained above (or as in the Equation (38)), then

$$h_0(X) = \|X_1\|^2 + r^2\|X_2\|^2 + \|X_3\|^2$$

where the norms come from the standard metrics, respectively, on $T[0, 1)$, on $TS^{n-l-1} \supset T\omega_p$, and on $T\mathbb{R}^l$.

Let us assume first that $h = h_0$, the standard metric on \mathbb{R}^n . By the induction hypothesis, we can construct $Y_2, \dots, Y_{n-l} \in \mathcal{V}(\omega_p)$ forming a local orthonormal basis of \mathcal{V} over some small neighborhood U' of x' in $\Sigma(\omega_p)$ (i.e., $\{Y_2, \dots, Y_{n-l}\} \subset \mathcal{V}(\omega_p)$ is orthonormal with respect to the metric $r_{\omega_p}^{-2}(dx')^2$). Here $(dx')^2$ denotes the metric on ω_p induced by the Euclidean metric on the sphere S^{n-l-1} , as above. Let $Y_1 = r_\Omega \partial_r$ and $Y_j = r_\Omega \partial_j$, $j = n-l+1, \dots, n$, where ∂_j forms the standard basis of \mathbb{R}^{l-1} . Then we claim that we can take $U = [0, 1) \times U' \times B^l$ and

$$(44) \quad \{X_1, X_2, \dots, X_n\} = \{Y_1\} \cup \{Y_2, \dots, Y_{n-l}\} \cup \{Y_{n-l+1}, \dots, Y_n\}.$$

(If $n-l=1$, then the second set in the above union is empty. If $l=0$, then the third set in the above union is empty.) Indeed, $\{X_1, \dots, X_n\}$ is a local basis by construction and by the local definition of $\mathcal{V}(\Omega)$ in Equation (38). Let us check that this is an orthonormal local basis. To this end, we shall use the form of the standard metric h_0 given in Equation (43), to obtain

$$h_0(X_1) = r_\Omega^2 \|\partial_r\|^2 = r_\Omega^2, \quad h_0(X_{n-l+1}) = \dots = h_0(X_n) = r_\Omega^2$$

$$\text{and } h_0(X_2) = \dots = h_0(X_{n-l}) = r^2 \|X_2\|^2 = r^2 r_{\omega_p}^2 = r_\Omega^2.$$

It is also clear that $\{X_1, X_2, \dots, X_n\}$ is an orthogonal system. This completes the induction step if $h = h_0$, the standard metric on \mathbb{R}^n .

If h is not the standard metric on V_l , we can nevertheless chose a matrix valued function T defined on a neighborhood of q in U such that $h(T\xi, T\eta) = h_0(\xi, \eta)$. We then let $X_j = TY_j$ and replace U with this smaller neighborhood. \square

This lemma gives the following corollary.

Corollary 4.18. *Let $X, Y \in \mathcal{V}(\Omega)$ and h be a fixed metric on M . Then the function $r_\Omega^{-2}h(X, Y)$, defined first on Ω , extends to a smooth function on $\Sigma(\Omega)$.*

Proof. This is a local statement in the neighborhood of each point $q \in \Sigma(\Omega)$. Let X_1, X_2, \dots, X_n be a local basis of \mathcal{V} on a neighborhood U of q in $\Sigma(\Omega)$ satisfying the conditions of Proposition 4.17 (i.e., orthogonal with respect to $r_\Omega^{-2}h$). Let $X = \sum \phi_j X_j$ and $Y = \sum \psi_j X_j$ on $U \cap \Omega$, where ϕ_j, ψ_j are smooth functions on $\Sigma(\Omega)$. Then $r_\Omega^{-2}h(X, Y) = \sum \phi_j \bar{\psi}_j$ is smooth on U . \square

Lemma 4.19. *Let $p \in \partial\Omega$ and X_1, X_2, \dots, X_n be vector fields on $\bar{\Omega}$ that define a local basis of TM on \bar{U} , for some neighborhood U of p . Then $r_\Omega X_1, r_\Omega X_2, \dots, r_\Omega X_n$ is a local basis of $\mathcal{V}(\Omega)$ on U , that is, for any $Y \in \mathcal{V}(\Omega)$, there exist unique smooth function $\phi_1, \phi_2, \dots, \phi_n$ on $\Sigma(\Omega)$ satisfying*

$$(45) \quad Y = \phi_1 r_\Omega X_1 + \phi_2 r_\Omega X_2 + \dots + \phi_n r_\Omega X_n \quad \text{on } U \cap \Omega \subset \Sigma(\Omega).$$

Conversely, if a vector field Y on Ω satisfies Condition (45) for any p and any local basis X_1, \dots, X_n of TM at p , then Y is the restriction to Ω of a vector field in $\mathcal{V}(\Omega)$.

Proof. The converse part is easier, so we prove it first. Let Y be a vector field on Ω that satisfies Condition (45) for any p and any local basis X_1, \dots, X_n of TM at p . Fix an arbitrary $p \in \Omega$. Lemmata 4.13 and 4.16 give that $\phi_j r_\Omega X_j$ is the restriction to Ω of a vector field in $\mathcal{V}(\Omega)$. Hence on each $U \cap \Omega$, Y is the restriction of a vector field $Y_U \in \mathcal{V}(\Omega)$. Lemma 4.15 then gives the converse part of our lemma.

We now prove the direct part of the lemma. We can assume that the vector fields X_1, \dots, X_n form an orthonormal system on U with respect to some fixed metric h on M . We know from Lemma 4.16 that $r_\Omega X_j \in \mathcal{V}(\Omega)$.

Let then $Y \in \mathcal{V}(\Omega)$ and note that $\phi_j = r_\Omega^{-1}h(Y, X_j) = r_\Omega^{-2}h(Y, r_\Omega X_j) \in C^\infty(\Sigma(\Omega))$, by Corollary 4.18. Then $Y = \sum_{j=1}^n \phi_j r_\Omega X_j$ on $U \cap \Omega$. The local uniqueness of the functions ϕ_j follows from the fact that $r_\Omega X_1, r_\Omega X_2, \dots, r_\Omega X_n$ also form a local basis of $T\Omega$ on $U \cap \Omega$. \square

We are now ready to prove the following characterizations of $\mathcal{V}(\Omega)$. We notice that the restriction map $\mathcal{V}(\Omega) \ni X \rightarrow X|_\Omega$ is injective, so we may identify $\mathcal{V}(\Omega)$ with a subspace of the space $\Gamma(\Omega, TM)$ of vector fields on Ω .

Proposition 4.20. *Let $\Omega \subset M$ be a curvilinear, stratified polyhedral domain of dimension n and let X be a smooth vector field on Ω . Fix an arbitrary metric h on M . Then $X \in \mathcal{V}(\Omega)$ if, and only if, $r_\Omega^{-1}h(X, Y)$ extends to a smooth function on $\Sigma(\Omega)$ for any smooth vector field Y on $\bar{\Omega}$.*

Proof. In one direction the result follows from Lemma 4.16 and Corollary 4.18. Indeed, let $X \in \mathcal{V}(\Omega)$ and Y be a smooth vector field on $\bar{\Omega}$. Then $r_\Omega Y \in \mathcal{V}(\Omega)$ by Lemma 4.16 and hence $r_\Omega^{-1}h(X, Y) = r_\Omega^{-2}h(X, r_\Omega Y)$ extends to a smooth function on $\Sigma(\Omega)$ by Corollary 4.18. (We have already used this argument in the proof of the previous lemma.)

Conversely, assume that $r_\Omega^{-1}h(X, Y)$ extends to a smooth function on $\Sigma(\Omega)$ for any smooth vector field on $\overline{\Omega}$. The statement that $X \in \mathcal{V}(\Omega)$ is a local statement, by Lemma 4.15. So let $p \in \overline{\Omega}$ and let U be an arbitrary neighborhood of p . Choose smooth vector fields defined in a neighborhood of $\overline{\Omega}$ in M such that X_1, X_2, \dots, X_n is a local orthonormal basis on U (orthonormal with respect to h). Let

$$\phi_j = r_\Omega^{-1}h(Y, X_j),$$

by assumption $\phi_j \in \mathcal{C}^\infty(\Sigma(\Omega))$. Then $Y = \sum_{j=1}^n \phi_j X_j$ on $U \cap \Omega$ and $\sum_{j=1}^n \phi_j X_j \in \mathcal{V}(\Omega)$. Lemma 4.15 then shows that $X \in \mathcal{V}(\Omega)$. \square

We now prove the main characterization of the structural Lie algebra of vector fields $\mathcal{V}(\Omega)$.

Theorem 4.21. *Let $\Omega \subset M$ be a bounded curvilinear, stratified polyhedral domain of dimension n . Then $\mathcal{V}(\Omega)$ is generated as a vector space by the vector fields of the form $\phi r_\Omega X$, where $\phi \in \mathcal{C}^\infty(\Sigma(\Omega))$ and X is a smooth vector field on $\overline{\Omega}$.*

Proof. We know that $\phi r_\Omega X \in \mathcal{V}(\Omega)$ whenever X is a smooth vector field on $\overline{\Omega}$, by Lemmata 4.13 and 4.16. This remark shows that the linear span of vectors of the form $\phi r_\Omega X$, where $\phi \in \mathcal{C}^\infty(\Sigma(\Omega))$ and X is a smooth vector field in a neighborhood of Σ , is contained in $\mathcal{V}(\Omega)$.

Conversely, let $Y \in \mathcal{V}(\Omega)$. Then Lemma 4.19 shows that we can find, in the neighborhood U_p of any point $p \in \overline{\Omega}$ vector fields $X_{1p}, X_{2p}, \dots, X_{np}$ and smooth functions ϕ_{jp} such that $Y = \sum \phi_{jp} r_\Omega X_{jp}$ on U_p . The result then follows using a finite partition of unity on $\Sigma(\Omega)$ subordinated to the covering U_p . \square

If we drop the condition that Ω be bounded, we obtain the following result, which was established in the first half of the above proof.

Proposition 4.22. *Let $\Omega \subset M$ be a curvilinear polyhedral domain of dimension n . Then $\mathcal{V}(\Omega)$ consists of the set of vector fields that locally can be written as linear combinations of vector fields of the form $\phi r_\Omega X$, where $\phi \in \mathcal{C}^\infty(\Sigma(\Omega))$ and X is a smooth vector field on $\overline{\Omega}$.*

We are finally in the position to endow $\Sigma(\Omega)$ with a structure of Lie manifold, which we will exploit in the following sections to study the mixed boundary value/interface problem (6). We set $\partial'\Sigma(\Omega)$ to be the union of all hyperfaces (*i.e.*, faces of maximal dimension) H of $\Sigma(\Omega)$ such that $\kappa(H) \subset \overline{\Omega}$ lies in the singular set $\Omega^{(n-2)}$, and let $\partial''\Sigma(\Omega) = \partial\Sigma(\Omega) \setminus \partial'\Sigma(\Omega)$. The next theorem is the principal result of this subsection.

Theorem 4.23. *Let Ω be a bounded curvilinear, stratified polyhedral domain and let*

$$\mathfrak{D}_0 := \Sigma(\Omega) \setminus \partial''\Sigma(\Omega) = \Omega \cup \partial'\Sigma(\Omega) = \kappa^{-1}(\overline{\Omega} \setminus \Omega^{(n-2)}).$$

Then $(\mathfrak{D}_0, \Sigma(\Omega), \mathcal{V}(\Omega))$ is a Lie manifold with boundary $\partial'\Sigma(\Omega)$. The projection map $\kappa : \mathfrak{D}_0 \rightarrow \overline{\Omega} \setminus \Omega^{(n-2)}$ is such that $\kappa^{-1}(p)$ consists of exactly one point.

Before proceeding with the proof, we observe that the above theorem justifies the following definition (cf. the corresponding definition for Lie manifolds after Definition (3.7)).

Definition 4.24. Any hyperface H of $\Sigma(\Omega)$ is called a *hyperface at infinity* if $\kappa(H) \subset \Omega^{(n-2)}$.

Proof. The last statement (on the number of elements in $\kappa^{-1}(p)$, $p \in \bar{\Omega} \setminus \Omega^{(n-2)}$) follows from the definition. Therefore, to prove the proposition, we need, using the notation of Definition 3.5, to construct a compactification \mathfrak{D} of \mathfrak{D}_0 that identifies with the closure of a Lie domain in a Lie manifold \mathfrak{M} .

We shall choose then $\mathfrak{D} = \Sigma(\Omega)$. Then we shall let \mathfrak{M} be the “double” of $\Sigma(\Omega)$, also denoted ${}^d\Sigma(\Omega)$. More precisely, \mathfrak{M} is obtained from the disjoint union of two copies of $\Sigma(\Omega)$ by identifying the hyperfaces that are not at infinity. We let \mathcal{V} to be the set of vector fields on \mathfrak{M} such that the restriction to either copy of $\Sigma(\Omega)$ is in $\mathcal{V}(\Omega)$.

Let \mathbb{D} be obtained from the closure of Ω in \mathfrak{M} by removing the closure of $\partial'\Sigma(\Omega)$. Then \mathbb{D} is an open subset of \mathfrak{M} whose closure is $\Sigma(\Omega)$. Moreover, $\partial_{\mathfrak{M}}\mathbb{D}$ (the boundary of \mathbb{D} regarded as a subset of \mathfrak{M}) is the closure of $\partial'\Sigma(\Omega)$. To prove our theorem, we shall check that \mathfrak{M} is a manifold with corners, that $(\mathfrak{M}, \mathcal{V})$ is a Lie manifold, and that $\partial\mathbb{D}$ is a regular submanifold of \mathfrak{M} . Each of these properties is local, so it can be checked in the neighborhood of a point of \mathfrak{M} .

Fix $V_p = (0, \epsilon) \times \omega_p \times B^l$. Then the union of the two copies of $\Sigma(V_p)$ is the double ${}^d\Sigma(V_p)$ of $\Sigma(V_p)$. Denote by ${}^d\omega_p$ the double of ω_p . Then

$${}^d\Sigma(V_p) = [0, \epsilon) \times {}^d\omega_p \times B^l.$$

An inductive argument then shows that ${}^d\Sigma(\Omega)$ is a manifold with corners and that $\partial\mathbb{D}$ is a regular submanifold of \mathfrak{M} .

Let us check that \mathcal{V} satisfies the conditions defining a Lie manifold structure on \mathfrak{M} . It follows from Theorem 4.21 that \mathcal{V} is a $\mathcal{C}^\infty(\mathfrak{M})$ -module (this checks condition (iii) of Definition 3.2). Theorem 4.21 and Lemmata 4.14, 4.13 show that \mathcal{V} is closed under Lie brackets (this checks condition (i) of Definition 3.2). Condition (ii) of that definition follows from the definition of $\mathcal{V}(\Omega)$. Condition (iv) of Definition 3.2 as well as Condition (ii) of Definition 3.3 were proved in Lemma 4.19. This shows that $(\mathfrak{M}, \mathcal{V})$ is a Lie manifold. \square

An immediate consequence of the above Proposition is that the boundary $\partial\mathfrak{D}_0 = \partial'\Sigma(\Omega)$ of $\mathfrak{D}_0 = \Sigma(\Omega) \setminus \partial''\Sigma(\Omega)$ will acquire the structure of a Lie manifold, as explained after the definition of a Lie manifold with boundary, Definition 3.5. Let D be the closure of $\partial\mathfrak{D}_0$ in \mathfrak{D} . Then the Lie structure at infinity is $(\partial\mathfrak{D}_0, D, \mathcal{W})$, where

$$(46) \quad \mathcal{W} = \{X|_D, X \in \mathcal{V}, X|_D \text{ is tangent to } D\}.$$

As always, $X \in \mathcal{W}$ is completely determined by its restriction to \mathfrak{D}_0 .

5. WEIGHTED SOBOLEV SPACES

One of the goal of this work, as mentioned already, is the study of mixed boundary value/interface problems for second-order elliptic operators on n -dimensional curvilinear, stratified polyhedral domains Ω in the framework of certain weighted Sobolev spaces. This framework is adapted to the singular geometry of polyhedral domains and allows to obtain optimal regularity, which does not hold in general in the standard (unweighted) spaces.

We begin by giving a rigorous definition of the weighted spaces. Let f be a continuous function on Ω , $f > 0$ on the interior of Ω . We define the μ th Sobolev space with weight f and index a by

$$(47) \quad \mathcal{K}_{a,f}^\mu(\Omega) = \{u \in L_{\text{loc}}^2(\Omega), f^{|\alpha|-a} \partial^\alpha u \in L^2(\Omega), \text{ for all } |\alpha| \leq \mu\}, \quad \mu \in \mathbb{Z}_+.$$

The norm on $\mathcal{K}_{a,f}^\mu(\Omega)$ is given by

$$(48) \quad \|u\|_{\mathcal{K}_{a,f}^\mu(\Omega)}^2 := \sum_{|\alpha| \leq \mu} \|f^{|\alpha|-a} \partial^\alpha u\|_{L^2(\Omega)}^2.$$

Definition 5.1. Let f, g be two continuous, non-negative functions on Ω . We shall say that f and g are *equivalent* (written $f \sim g$) if there exists a constant $C > 0$ such that

$$C^{-1}f(x) \leq g(x) \leq Cf(x),$$

for all $x \in \Omega$.

Clearly, if $f \sim g$, then the norms $\|u\|_{\mathcal{K}_{a,f}^\mu(\Omega)}$ and $\|u\|_{\mathcal{K}_{a,g}^\mu(\Omega)}$ are equivalent, and hence we have $\mathcal{K}_{a,f}^\mu(\Omega) = \mathcal{K}_{a,g}^\mu(\Omega)$ as Banach spaces.

Definition 5.2. We let $\mathcal{K}_a^\mu(\Omega) = \mathcal{K}_{a,f}^\mu(\Omega)$ and $\|u\|_{\mathcal{K}_a^\mu(\Omega)} = \|u\|_{\mathcal{K}_{a,f}^\mu(\Omega)}$, where $f = \eta_{n-2}$ is the distance to $\Omega^{(n-2)}$.

For example, $\mathcal{K}_0^0(\Omega) = L^2(\Omega)$. For Ω a polygon in the plane, $\eta_{n-2}(x) = \eta_0(x)$ is the distance from x to the vertices of Ω and the resulting spaces $\mathcal{K}_a^\mu(\Omega)$ are the spaces considered in Kondratiev's paper [30]. Above in Definition 5.2, we can and will replace η_{n-2} with the equivalent function r_Ω by Proposition 4.9.

If $\mu \in \mathbb{N} = \mathbb{Z}_+ \setminus \{0\}$, we define $\mathcal{K}_a^{-\mu}(\Omega)$ to be the dual of

$$(49) \quad \mathring{\mathcal{K}}_a^\mu(\Omega) := \mathcal{K}_a^\mu(\Omega) \cap \{\partial_\nu^j u|_{\partial\Omega} = 0, j = 0, 1, \dots, \mu - 1\}$$

with pivot $\mathcal{K}_0^0(\Omega)$. Later in this Section, we will identify $\mathcal{K}_a^m(\Omega)$ with a suitable space $hH^\mu(\Sigma(\Omega))$ using the Lie structure on $\Sigma(\Omega)$. Then, Theorem 3.4 of [1] (see also Lemma 3.7 and Proposition 3.8) gives that $\mathcal{C}_c^\infty(\Omega)$ is dense in $\mathring{\mathcal{K}}_a^\mu(\Omega)$ and consequently $\mathcal{K}_a^{-\mu}(\Omega)$ is the completion of the space of smooth functions u on Ω satisfying

$$(50) \quad \|u\|_{\mathcal{K}_a^{-\mu}(\Omega)} = \sup_{0 \neq v \in \mathcal{C}_c^\infty(\Omega)} \frac{|(u, v)|}{\|v\|_{\mathcal{K}_a^\mu(\Omega)}} < +\infty.$$

In order to make the identification $\mathcal{K}_a^m(\Omega) \approx hH^\mu(\Sigma(\Omega))$, we introduce next a class of "admissible weights" h .

5.1. The set of weights. If $h > 0$ on Ω , we denote

$$(51) \quad h\mathcal{K}_a^\mu(\Omega) := \{hu, u \in \mathcal{K}_a^\mu(\Omega)\},$$

with induced norm, that is $\|hu\|_{h\mathcal{K}_a^\mu(\Omega)} = \|u\|_{\mathcal{K}_a^\mu(\Omega)}$.

A main example of an admissible weight is η_{n-2}^a , for $a \in \mathbb{R}$. This example is sufficient to recover the weighted Sobolev spaces considered in [39] for instance. The definition of a general admissible weight involves the desingularization $\Sigma(\Omega)$. Let H be a hyperface at infinity of $\Sigma(\Omega)$, according to Definition 4.24, and let x_H be its defining function.

Definition 5.3. A function $h = \prod_H x_H^{a_H}$, where H ranges through the set of hyperfaces at infinity of $\Sigma(\Omega)$ and $a_H \in \mathbb{R}$, will be called an *admissible weight*. We denote the set of admissible weights by $\mathcal{W}(\Omega)$ and endow it with the topology defined by the exponents a_H .

As discussed after Corollary 4.11, we can always assume $r_\Omega := \prod_H x_H$. In particular, r_Ω^a , $a \in \mathbb{R}$, is the most important example of an admissible weight. It is more suitable to use this weight, which is intimately related to the structure of Lie manifold on $\Sigma(\Omega)$ ($\setminus \partial^n \Sigma(\Omega)$) described in Theorem 4.23, instead of η_{n-2}^a . We also have that

$$(52) \quad r_\Omega^t \mathcal{K}_a^\mu(\Omega) = \mathcal{K}_{a+t}^\mu(\Omega),$$

so in a statement about the spaces $h\mathcal{K}_a^\mu(\Omega)$, where h is an admissible weight, we can usually assume that $a = 0$, without loss of generality. All these spaces are therefore Babuška–Kondratiev spaces in the sense of the following definition.

Definition 5.4. Let h be an admissible weight on Ω . The *Babuška–Kondratiev* space of order $\mu \in \mathbb{Z}$ and weight h on Ω is the space $h\mathcal{K}_0^\mu(\Omega)$.

5.2. Sobolev spaces and Lie manifolds. We now identify the weighted Sobolev space $\mathcal{K}_a^\mu(\Omega)$ with $hH^\mu(\Sigma(\Omega))$, for a suitable admissible weight h ; more precisely, $h = r_\Omega^{a-n/2}$. The following description of $\mathcal{V}(\Omega)$ for Ω a curvilinear polyhedral domain in \mathbb{R}^n will be useful.

Corollary 5.5. Let $\Omega \subset \mathbb{R}^n$ be a bounded curvilinear, polyhedral domain. Then

$$\mathcal{V}(\Omega) = \{\phi_1 r_\Omega \partial_1 + \phi_2 r_\Omega \partial_2 + \dots + \phi_n r_\Omega \partial_n, \text{ where } \phi_j \in \mathcal{C}^\infty(\Sigma(\Omega))\}.$$

We shall denote by

$$(53) \quad \text{Diff}_\Omega^k := \text{Diff}_{\mathcal{V}(\Omega)}^k(\Sigma(\Omega))$$

the space of differential operators with coefficients in $\mathcal{C}^\infty(\Sigma(\Omega))$ of order $\leq k$ on $\Sigma(\Omega)$ generated by $\mathcal{V}(\Omega)$. The algebra of differential operators $\text{Diff}_\Omega^\infty$ is an example of the algebra of differential operators considered in 3.3. From the last corollary, we obtain directly the following lemma.

Lemma 5.6. Let X_1, X_2, \dots, X_k be smooth vector fields on M . Then

$$P := r_\Omega^k X_1 X_2 \dots X_k \in \text{Diff}_\Omega^k,$$

and ϕP , with P as above and $\phi \in \mathcal{C}^\infty(\Sigma(\Omega))$ generate Diff_Ω^k linearly.

Proof. For $k = 1$, this follows from Lemma 4.16. Next, we have

$$r_\Omega^{k+1} X_0 X_1 \dots X_k = r_\Omega X_0 r_\Omega^k X_1 \dots X_k - k X_0 (r_\Omega) r_\Omega^k X_1 \dots X_k.$$

The fact that $P \in \text{Diff}_\Omega^k$ then follows by induction, since $X_0(r_\Omega) \in \mathcal{C}^\infty(\Sigma(\Omega))$, by Lemma 4.14.

Conversely, we can similarly check by induction (using the same identity above) that the product $r_\Omega X_1 r_\Omega X_2 \dots r_\Omega X_k$ can be written as a linearly combination of differential operators of the form ϕP , with $\phi \in \mathcal{C}^\infty(\Sigma(\Omega))$ and P as above. Since $r_\Omega X$ generate $\mathcal{V}(\Omega)$ as a $\mathcal{C}^\infty(\Sigma(\Omega))$ -module (see the second part of Theorem 4.21), the result follows. \square

We next provide a different description of the weighted Sobolev spaces $\mathcal{K}_a^\mu(\Omega)$, $\mu \in \mathbb{Z}_+$. For a multiindex α , we denote

$$(54) \quad (r_\Omega \partial)^\alpha := (r_\Omega \partial_1)^{\alpha_1} (r_\Omega \partial_2)^{\alpha_2} \dots (r_\Omega \partial_n)^{\alpha_n}.$$

Theorem 5.7. *Let $\Omega \subset \mathbb{R}^n$ be a bounded curvilinear, stratified polyhedral domain and*

$$\|u\|_{\mu,a}^2 := \sum_{|\alpha| \leq \mu} \|r_\Omega^{-a} (r_\Omega \partial)^\alpha u\|_{L^2(\Omega)}^2.$$

Then $\|u\|_{\mu,a}$ is equivalent to $\|u\|_{\mathcal{K}_a^\mu(\Omega)}$ of Definition 5.2. In particular,

$$\mathcal{K}_a^\mu(\Omega) = \{u, \|u\|_{\mu,a} < \infty\}.$$

Proof. We have that

$$\begin{aligned} u &\in \mathcal{K}_a^\mu(\Omega) \\ &\Leftrightarrow r_\Omega^{|\alpha|-a} \partial^\alpha u \in L^2(\Omega) \quad \text{for all } |\alpha| \leq \mu \quad \text{by Proposition 4.9} \\ &\Leftrightarrow r_\Omega^{-a} (r_\Omega \partial)^\alpha u \in L^2(\Omega) \quad \text{for all } |\alpha| \leq \mu \quad \text{by Proposition 5.6.} \end{aligned}$$

Above the corresponding square integrability conditions define the topology on the indicated spaces. Therefore \Leftrightarrow also means that the topologies are the same. \square

We are in position to identify the spaces \mathcal{K}_a^μ with Sobolev spaces on Lie manifolds. If Ω bounded curvilinear, stratified polyhedral domain, we let

$$\mathfrak{D}_0 := \Sigma(\Omega) \setminus \partial''\Sigma(\Omega) = \Omega \cup \partial'\Sigma(\Omega) = \kappa^{-1}(\bar{\Omega} \setminus \Omega^{(n-2)}),$$

as in Theorem 4.23. Since $(\mathfrak{D}_0, \mathfrak{D} := \Sigma(\Omega), \mathcal{V}(\Omega))$ is a Lie manifold with boundary by the same theorem, the definitions of Sobolev spaces on Lie manifolds (with or without boundary) of Subsection 3.5 provide us with natural spaces $H^s(\Sigma(\Omega)) = H^s(\mathfrak{D}) = H^s(\mathfrak{D}_0)$ and $H^s(\partial'\Sigma(\Omega)) = H^s(\partial\mathfrak{D}_0)$. For the last equality we used that the boundary of \mathfrak{D}_0 is $\partial'\Sigma(\Omega)$.

Proposition 5.8. *Let Ω be an n -dimensional, bounded curvilinear, stratified polyhedral domain and let h be an admissible weight on Ω . We have an equality*

$$h\mathcal{K}_a^\mu(\Omega) = hr_\Omega^{a-n/2} H^\mu(\Sigma(\Omega)), \quad \mu \in \mathbb{Z}.$$

Proof. This is again a local statement. We can therefore assume that $\Omega \subset \mathbb{R}^n$. Furthermore, it is enough to prove the statement in the case $h = 1$, since the weight h does not enter into the condition on derivatives in the definition 51 of weighted spaces. Equation (52) and Proposition 4.9 show that we can also assume $a = 0$. Recall from Lemma 3.7 that the spaces $H^k(\Sigma(\Omega))$ are defined using $L^2(\Sigma(\Omega))$. In turn, $L^2(\Sigma(\Omega))$ is defined using the volume element of a compatible metric. A typical compatible metric is $r_\Omega^{-2} g_e$, where g_e is the Euclidean metric. Therefore the volume element on $\Sigma(\Omega)$ is $r_\Omega^{-n} dx$, where dx is the Euclidean volume element. In particular, $v \in L^2(\Omega) \Leftrightarrow v \in r_\Omega^{-n/2} L^2(\Sigma(\Omega))$.

We notice next that $r_\Omega^{-t} (r_\Omega \partial)^\alpha r_\Omega^t - (r_\Omega \partial)^\alpha$ is a linear combination with $\mathcal{C}^\infty(\Sigma(\Omega))$ -coefficients of monomials $(r_\Omega \partial)^\beta$, with $|\beta| < |\alpha|$, by the second part of Lemma 4.14.

From this observation we obtain

$$\begin{aligned} u &\in \mathcal{K}_0^\mu(\Omega) \Leftrightarrow (r_\Omega \partial)^\alpha u \in L^2(\Omega) \quad \text{for all } |\alpha| \leq \mu \quad \text{by Theorem 5.7} \\ &\Leftrightarrow (r_\Omega \partial)^\alpha u \in r_\Omega^{-n/2} L^2(\Sigma(\Omega)) \quad \text{for all } |\alpha| \leq \mu \\ &\Leftrightarrow (r_\Omega \partial)^\alpha r_\Omega^{n/2} u \in L^2(\Sigma(\Omega)) \quad \text{for all } |\alpha| \leq \mu \Leftrightarrow u \in r_\Omega^{-n/2} H^\mu(\Sigma(\Omega)). \end{aligned}$$

This proves that $\mathcal{K}_a^\mu(\Omega) = r_\Omega^{a-n/2} H^\mu(\Sigma(\Omega))$ for $\mu \in \mathbb{Z}_+$. For $\mu \in \mathbb{Z}_-$, we observe that, for $(\mathfrak{D}, \mathfrak{D}_0, \mathcal{V})$ a Lie manifold with boundary in a manifold with corner \mathfrak{M} , the set of restrictions of distributions $u \in H^{-\mu}(\mathfrak{M})$ to \mathfrak{D}_0 is the dual of the closure of $\mathcal{C}_c^\infty(\mathfrak{D}_0)$ in $H^{-\mu}(\mathfrak{M})$. Hence

$$\mathcal{K}_0^{-\mu}(\Omega) := \mathring{\mathcal{K}}_a^\mu(\Omega)^* = (r_\Omega^{-n/2} \mathring{H}^\mu(\Sigma(\Omega)))^* = r_\Omega^{-n/2} H^{-\mu}(\Sigma(\Omega)).$$

The proof is concluded. \square

The identification given in Proposition 5.8 above allows to define weighted spaces on the boundary $h\mathcal{K}_a^m(\partial\Omega)$. We recall that the closure of a hyperface of a curvilinear, stratified polyhedral domain Ω need not be contained in any smooth $n-1$ manifold. Consequently, we utilize the desingularization $\Sigma(\Omega)$. In the special case that $\Omega \subset \mathbb{R}^n$ is a (bounded) convex, stratified polyhedron that in addition has straight faces (*i.e.*, each connected component $D_j^{(l)}$ of $\Omega^{(l)} \setminus \Omega^{(l-1)}$, $l = 1, \dots, n-1$ is contained in an affine space $V_j^{(l)}$ of dimension l), for example an n -simplex, we can more simply define the spaces on the boundary as follows:

(55)

$$\mathcal{K}_a^\mu(D_j^{(n-1)}) = \{u \in L_{\text{loc}}^2(D_j^{(n-1)}), r_\Omega^{k-a} X_1 X_2 \dots X_k u \in L^2(D_j^{(n-1)}), 0 \leq k \leq l\},$$

for all choices of vector fields X_j in a basis of the linear space containing $D_j^{(n-1)}$. Then for any admissible weight h ,

$$(56) \quad h\mathcal{K}_a^\mu(\partial\Omega) = \{hu, u \in L_{\text{loc}}^2(\partial\Omega), u|_{D_j^{(n-1)}} \in \mathcal{K}_a^\mu(D_j^{(n-1)}), \text{ for all } j\},.$$

In the general case of a curvilinear, stratified polyhedral domain, we exploit the structure of Lie manifold on $\Sigma(\Omega)$, following the notation of Proposition 5.8.

Definition 5.9. Let Ω be a bounded, curvilinear, stratified polyhedral domain. Then we define

$$h\mathcal{K}_a^\mu(\partial\Omega) := hr_\Omega^{a-(n-1)/2} H^\mu(\partial'\Sigma(\Omega)),$$

for any admissible weight h .

Note that on each hyperface, the natural weight is the distance to the boundary, not the distance to the set of singular boundary points of that face. The spaces $\mathcal{K}_{-a}^{-\mu}(\partial\Omega)$ are defined to be the duals of $\mathcal{K}_a^\mu(\partial\Omega)$ with pivot $L^2(\partial\Omega)$. For reasons that will be explained later, we do not have to restrict to functions with vanishing trace when studying weighted Sobolev spaces on the boundary. In particular, the usual difficulties that appear in the treatment of Sobolev spaces of fractional order on smooth, bounded domains [36], do not arise when studying the weighted Sobolev spaces on $\partial\Omega$, and we can define the spaces $\mathcal{K}_a^s(\partial\Omega)$, with $s \notin \mathbb{Z}$, by complex interpolation. A similar attempt at defining $\mathcal{K}_a^s(\Omega)$, with $s \in \mathbb{Z} + 1/2$, would lead to the usual difficulties encountered in the case of smooth domain, [36].

We next prove a trace theorem, generalizing the corresponding well-known result for smooth domains. Let $\mathcal{C}_c^\infty(\Omega)$ be the space of compactly supported functions on the open set Ω .

Theorem 5.10. *The restriction $\mathcal{C}_c^\infty(\bar{\Omega} \setminus \Omega^{(n-2)}) \ni u \rightarrow u|_{\partial\Omega} \in \mathcal{C}_c^\infty(\partial\Omega \setminus \Omega^{(n-2)})$ extends to a continuous, surjective map*

$$Tr : \mathcal{K}_a^\mu(\Omega) \rightarrow \mathcal{K}_{a-1/2}^{\mu-1/2}(\partial\Omega), \quad \mu \geq 1.$$

Moreover, $\mathcal{C}_c^\infty(\Omega)$ is dense in the kernel of this map if $\mu = 1$.

The result is a consequence of similar results for Lie manifolds contained in Theorems 3.4 and 3.7 of [1] recalled here in Proposition 3.8.

Proof. The map $H^\mu(\Sigma(\Omega)) \rightarrow H^{\mu-1/2}(\partial'\Sigma(\Omega))$, where we follow the notation of Proposition 5.8, is well defined, continuous, and surjective by Proposition 3.8. Proposition 5.8 then shows that the map

$$h\mathcal{K}_a^\mu(\Omega) = hr_\Omega^{a-n/2}H^\mu(\Omega) \rightarrow hr_\Omega^{a-n/2}H^{\mu-1/2}(\partial\Omega) = h\mathcal{K}_{a-1/2}^{\mu-1/2}(\Omega)$$

is also well defined, continuous, and surjective.

The density of $\mathcal{C}_c^\infty(\Omega)$ in the subspace of elements in $h\mathcal{K}_a^1(\Omega)$ with trace zero also follows from Proposition 3.8 and Proposition 5.8. \square

6. PROOFS

In this section, we establish the main results of the paper, Theorems 1.1, 1.2, 1.3, using material from previous sections. We first discuss some results on the behavior of differential operators on the spaces $h\mathcal{K}_a^m(\Omega)$.

6.1. Differential operators. We recall that the algebra $\text{Diff}_\Omega^\infty$ is the natural algebra of differential operators on Ω associated to the Lie algebra of vector fields $\mathcal{V}(\Omega)$, namely, it is generated as an algebra by $X \in \mathcal{V}(\Omega)$ and $\phi \in \mathcal{C}^\infty(\Sigma(\Omega))$. (This algebra was used also in Equation (53) and in Subsection 3.3.)

Proposition 6.1. *Let P be a differential operator of order m on a manifold M with smooth coefficients. Let $\Omega \subset M$ be a curvilinear, stratified polyhedral domain. Then P maps $h\mathcal{K}_a^\mu(\Omega)$ to $h\mathcal{K}_{a-m}^{\mu-m}(\Omega)$ continuously, for any admissible weight h and any $\mu \in \mathbb{Z}$. Moreover, the resulting family $h^{-\lambda}Ph^\lambda : \mathcal{K}_a^\mu(\Omega) \rightarrow \mathcal{K}_{a-m}^{\mu-m}(\Omega)$ of bounded operators depends continuously on λ .*

Before proceeding with the proof, we discuss a corollary, which will be relevant in showing that Theorems 1.2 and 1.3 hold. following the notation of those theorems, below $\mathcal{W}_\mu(\Omega)$ represents the set of admissible weights h such that the map

$$\tilde{P}(u) := (Pu, u|_{\partial_D\Omega}, D_\nu^P u|_{\partial_N\Omega}) \text{ is an isomorphism } \left\{ \bigoplus_{j=1}^N h\mathcal{K}_1^{\mu+1}(\Omega_j) \cap \mathcal{K}_1^1(\Omega); u^+ = u^-, D_\nu^P u^+ = D_\nu^P u^- \text{ on } \Gamma \right\} \simeq \bigoplus_{j=1}^N h\mathcal{K}_{-1}^{\mu-1}(\Omega_j) \oplus h\mathcal{K}_{1/2}^{\mu+1/2}(\partial_D\Omega) \oplus h\mathcal{K}_{-1/2}^{\mu-1/2}(\partial_N\Omega).$$

Proposition 6.2. *The set $\mathcal{W}_\mu(\Omega)$ is open.*

Proof. This follows directly from Proposition 6.1. Indeed, the family $P : \bigoplus_{j=1}^N h\mathcal{K}_1^{\mu+1}(\Omega_j) \cap h\mathcal{K}_1^1(\Omega) \rightarrow h\mathcal{K}_{-1}^{\mu-1}(\Omega)$ is unitarily equivalent to $h^{-1}Ph$: $\bigoplus_{j=1}^N \mathcal{K}_1^{\mu+1}(\Omega_j) \cap \mathcal{K}_1^1(\Omega) \rightarrow \mathcal{K}_{-1}^{\mu-1}(\Omega)$. The result then follows since the set of invertible operators is open. \square

For the proof of Proposition 6.1, we observe that if $\Omega \subset \mathbb{R}^n$, the principal symbol of $(r_\Omega\partial)^\alpha$ is ξ^α . This result follows from the definition of the principal symbol in [3, 1] and from Corollary 5.5. (The reader can just assume $\sigma((r_\Omega\partial)^\alpha) = \xi^\alpha$ by definition.)

Corollary 6.3. *Let P be a differential operator of order m on M with smooth coefficients. Then*

- (i) $r_\Omega^m P \in \text{Diff}_\Omega$;
- (ii) P is (strongly) elliptic if, and only if, $r_\Omega^m P$ is (strongly) elliptic in Diff_Ω^m ;
- (iii) $h^\lambda P h^{-\lambda}$ depends continuously on λ ;
- (iv) P maps $h\mathcal{K}_a^\mu(\Omega) \rightarrow h\mathcal{K}_{a-m}^{\mu-m}(\Omega)$ continuously;

Proof. The relation $r_\Omega^m P \in \text{Diff}_\Omega^m$ was proved as part of Lemma 5.6. Strong ellipticity is a local property, so we can assume $\Omega \subset \mathbb{R}^n$. The proof of Lemma 5.6 shows that P and $r_\Omega^m P$ have the same principal symbol. Therefore they are elliptic (or strongly elliptic) at the same time.

For any $X \in \mathcal{V}$ and any defining function x of some hyperface at infinity of $\Sigma(\Omega)$, we have that $x^\lambda X x^{-\lambda} = X - \lambda x^{-1} X(x)$. Since $x^{-1} X(x)$ is a smooth function (as X is tangent to the face defined by x), we see that $x^\lambda X x^{-\lambda} \in \text{Diff}_\Omega^1$ and depends continuously on λ , establishing (iii). It also shows, in particular, that Diff_Ω^k is conjugation invariant with respect to defining functions of hyperfaces at infinity (Equation (27)). We can therefore assume that $h = 1$.

Since $(\Sigma(\Omega), \mathcal{V}(\Omega))$ is a Lie manifold with boundary (Theorem 4.23) any $P \in \text{Diff}_\Omega^k$ maps $H^\mu(\Sigma(\Omega)) \rightarrow H^{\mu-k}(\Sigma(\Omega))$ continuously. (This simple property, proved in [1], is an immediate consequence of the definitions.) The continuity of $P : \mathcal{K}_a^\mu(\Omega) \rightarrow \mathcal{K}_{a-m}^{\mu-m}(\Omega)$ then follows using also the fact that multiplication by r_Ω^a defines an isometry $\mathcal{K}_a^{\mu-m}(\Omega) \simeq \mathcal{K}_{a-m}^{\mu-m}(\Omega)$. \square

6.2. A weighted Hardy–Poincaré’s inequality. The stepping stones in the proof of our main result on the solvability of the mixed boundary value/interface problem (6), Theorem 1.2, consist of

- (i) a Hardy–Poincaré type inequality (Theorem 6.4);
- (ii) the regularity result for polyhedra (Theorem 1.1).

We address the Hardy–Poincaré inequality first and turn to the proof of the regularity result, which is more general and of independent interest in the next subsection. Let $dx = dx_1 dx_2 \dots dx_n$ denote the standard volume element in \mathbb{R}^n . We continue to denote by Ω a curvilinear, stratified polyhedral domain satisfying hypotheses (3)–(5).

Theorem 6.4. *Let Ω be a connected, curvilinear, stratified polyhedral domain $\Omega \subset M$. Assume that $\partial_D \Omega \neq \emptyset$ and $\partial_N \Omega$ does not contain any two adjacent hyperfaces. Then there exists a constant $\kappa_\Omega > 0$, depending only on the polyhedral structure of Ω , such that*

$$(57) \quad \|u\|_{\mathcal{K}_1^0(\Omega)}^2 := \int_\Omega \frac{|u(x)|^2}{\eta_{n-2}(x)^2} dx \leq \kappa_\Omega \int_\Omega |\nabla u(x)|^2 dx,$$

for any function $u \in H_{\text{loc}}^1(\Omega)$ such that $u|_{\partial_D \Omega} = 0$.

Above, if u/r is not square integrable, the statement of the theorem is understood to mean that ∇u is not square integrable either. By Propositions 4.9 and 4.10, we can replace the distance to the singular set η_{n-2} with the more regular weight r_Ω .

The proof proceeds by induction on the dimension n . We discuss first the case $n = 2, 3$.

The case $n = 2$. In view of the local nature of the definition of a curvilinear, stratified polygonal domain, Definition 2.6, it will be sufficient to have the Hardy–Poincaré inequality in a sector. By abuse of notation, we shall write $u(r, \theta) := u(r \cos \theta, r \sin \theta)$ for a function $u(x_1, x_2)$ expressed in polar coordinates. The proof

of the following elementary lemma can be found in e.g. [47][Subsection 2.3.1]. See also [29].

Lemma 6.5. *Let $\mathcal{C} = \mathcal{C}_R(\alpha, \beta) := \{(r \cos \theta, r \sin \theta) \in \mathbb{R}^2, 0 < r < R, \beta < \theta < \alpha\}$, $0 < \alpha - \beta < 2\pi$. Then*

$$\int_{\mathcal{C}} \frac{|u|^2}{r^2} dx \leq \frac{\pi^2}{\alpha^2} \int_{\mathcal{C}} |\nabla u|^2 dx$$

for any $u \in H_{\text{loc}}^1(\mathcal{C})$ satisfying $u(r, \theta) = 0$ if $\theta = \beta$ or $\theta = \alpha$. The same result holds if \mathcal{C} is the disjoint union of domains $\mathcal{C}_R(\alpha, \beta)$, for different values of R, α , and β .

From the Lemma above, we obtain the case $n = 2$ in Theorem 6.4, the first step in our induction proof. A detailed proof can be found e.g. in the papers [8, 40].

Lemma 6.6. *Let Ω be a connected, curvilinear, stratified polygonal domain in a two dimensional manifold M . Assume that $\partial_D \Omega \neq \emptyset$ and $\partial_N \Omega$ does not contain any two adjacent sides of Ω . Fix an arbitrary metric g on M and let $\eta_0(z)$ be the distance from z to the vertices of Ω . Then there exists a constant $\kappa_\Omega > 0$ such that*

$$\int_{\Omega} \frac{|u(z)|^2}{\eta_0(z)^2} dz \leq \kappa_\Omega \int_{\Omega} |\nabla u(z)|^2 dz$$

for any $u \in H_{\text{loc}}^1(\Omega)$ satisfying $u = 0$ on $\partial_D \Omega$.

The case $n = 3$ The proof Theorem 6.4 for $n = 3$ combines the methods used in the previous two Lemmata and the inequality for the case $n = 2$. We give a self-contained proof again, especially because the induction step in the general case is very similar.

The general and new case $n > 3$ will be completed using Proposition 4.10.

Proof. Let us fix, for any $p \in \partial\Omega$, a neighborhood V_p of p in M and a diffeomorphism $\phi_p : V_p \rightarrow U = B^{3-l} \times B^l$ as in Definition 2.8, where $l = \ell(p)$. We denote $\mathcal{C} := \phi_p(V_p \cap \Omega)$. We shall use the notation ω_p introduced in that definition. By decreasing V_p , if necessary, we may assume that ϕ_p extends to a diffeomorphism defined in a neighborhood of \bar{V}_p in \mathbb{R}^3 .

Since $\eta_{n-2} = \eta_1$ is the distance to the singular set $\Omega^{(1)}$ of Ω , we need only discuss two cases:

- (a) $l = \ell(p) = 0$, i.e., p is a true or artificial vertex;
- (b) $l = \ell(p) = 1$, i.e., p belongs to a true or artificial edge.

If $l = 0$, we denote by $\psi_0(x')$ the distance from a point $x' \in \omega_p \subset S^2$ to the vertices of ω_p and let $r_p(z) = \rho\psi_0(x')$, if $\phi_p(z) = \rho x'$, where $0 < \rho$ and $x' \in \omega_p$. If $l = 1$, we let $r_p(z) = r$ if $\phi_p(z) = (r \cos \theta, r \sin \theta, z)$, where $0 < r, 0 < \theta < \alpha$, and $z \in \mathbb{R}$. (These definitions agree with the general definition of r_Ω given in (35) with $r_p = r_\alpha$ given in (34).) As before, the function $\eta_1(x)/r_p(x)$ is bounded for any p , provided that we choose the neighborhoods V_p small enough, uniformly in p . Below, we shall write $u(x)$ instead of $u(\phi_p^{-1}(x))$, by abuse of notation.

If $l = 1$, $\mathcal{C} = \mathcal{C}' \times (-1, 1)$, so that we exploit the Hardy-Poincaré inequality in a sector of Lemma 6.5. In fact

$$(58) \quad \int_{V_p \cap \Omega} \frac{|u(z)|^2}{\eta_1(z)^2} dz = C \int_{\Omega \cap V_p} \frac{|u(x)|^2}{r^2} \left| \frac{\partial z}{\partial x} \right| dx \leq C \int_{\mathcal{C}} \frac{|u(x)|^2}{r^2} dx.$$

so that we obtain

$$\begin{aligned}
(59) \quad \int_{\mathcal{C}} \frac{|u(x)|^2}{r^2} dx &= \int_{-1}^1 \left(\int_{\mathcal{C}'} \frac{|u(x)|^2}{r^2} dx_1 dx_2 \right) dx_3 \\
&\leq \int_{-1}^1 \left(\int_{\mathcal{C}'} |\nabla_{(x_1, x_2)} u(x)|^2 dx_1 dx_2 \right) dx_3 \leq \int_{-1}^1 \left(\int_{\mathcal{C}'} |\nabla u(x)|^2 dx_1 dx_2 \right) dx_3 \\
&\leq C \int_{V_p \cap \Omega} |\nabla u(z)|^2 dz.
\end{aligned}$$

We perform a similar calculation on $V_p \cap \Omega$ when $l = 0$, using spherical coordinates instead. Recall that $\mathcal{C} = \phi_p(V_p \cap \Omega) = \{\rho x', 0 < \rho < 1, x' \in \omega_p\}$, hence following (58) and using that $C\eta_1(x) \geq \rho\psi_0(x)$ The inequality

$$(60) \quad \int_{V_p \cap \Omega} \frac{|u(z)|^2}{\eta_1(z)^2} dz \leq C \int_{\mathcal{C}} \frac{|u(x)|^2}{\rho^2 \psi_0(x')^2} dx, \quad x = \rho x', |x'| = 1,$$

Next, we observe that $\nabla u(\rho x') = \rho^{-1} \nabla' u(\rho x') + \partial_\rho u(\rho x')$, with ∇' the gradient of a function defined on ω_p , so that $|\nabla' u(\rho x')|^2 \leq \rho^2 |\nabla u(\rho x')|^2$, which gives

$$\begin{aligned}
(61) \quad \int_{\mathcal{C}} \frac{|u(x)|^2}{\rho^2 \psi_0(x')^2} dx &= \int_0^1 \left(\int_{\omega_p} \frac{|u(\rho x')|^2}{\psi_0^2} dx' \right) d\rho \\
&\leq C \int_0^1 \left(\int_{\omega_p} |\nabla' u(\rho x')|^2 dx' \right) d\rho \leq C \int_0^1 \left(\int_{\omega_p} \rho^2 |\nabla u(\rho x')|^2 dx' \right) d\rho \\
&\leq C \int_{V_p \cap \Omega} |\nabla u(z)|^2 dz.
\end{aligned}$$

We can rewrite the above inequalities simply as

$$(62) \quad \int_{V_p \cap \Omega} \frac{|u(z)|^2}{\eta_1(z)^2} dz \leq C_p \int_{V_p \cap \Omega} |\nabla u(z)|^2 dz \leq C_p \int_{\Omega} |\nabla u(z)|^2 dz.$$

where the constant C_p depends on the point $p \in \Omega^{(1)}$ but not on u .

To conclude the proof, as before we cover the singular set $\Omega^{(1)}$ with finitely many sets $V_p = V_{p_k}$. Let $C_0 > \eta_0^{-2}$ outside the union of the sets V_{p_k} . Let $\kappa_\Omega = C_0 C_\Omega + \sum C_{p_k}$. We add all inequalities (62) for $p = p_k$ and combine it with the standard Poincaré inequality (to cover $\bar{\Omega} \setminus \Omega^{(1)}$) to give

$$(63) \quad \int_{\Omega} f(z) |u(z)|^2 dz \leq \kappa_\Omega \int_{\Omega} |\nabla u(z)|^2 dz,$$

where $f(z) \geq \eta_1(z)^{-2}$ on V_{p_j} and $f(z) \geq C_0 \geq \eta_1(z)^{-2}$ outside the union of the sets V_{p_k} . The proof of Theorem 6.4 is now complete for $n = 3$. \square

The general case $n > 3$. To conclude the proof of theorem 6.4, we need only establish the induction step. such step follows very closely the proof of the case $n = 3$ and we do not include a proof therefore for sake of brevity. The only delicate point is showing that the ratio $\eta_{n-2}(x)/r_\alpha(x)$ is bounded on $\bar{\Omega}$, where η_{n-2} is the distance to the singular set $\Omega^{(n-2)}$ of Ω and r_α is as in Equation 34. This fact was established in Proposition 4.9.

We conclude with an immediate corollary of Theorem 6.4, which will be used in the proof of Theorem 1.2

Corollary 6.7. *There exists a constant $\kappa'_\Omega > 0$, depending only on Ω , such that*

$$\frac{1}{\kappa'_\Omega} \|u\|_{\mathcal{K}_1^1(\Omega)}^2 \leq \int_{\Omega} |\nabla u(x)|^2 dx,$$

for any function $u \in H_{\text{loc}}^1(\Omega)$ such that $u|_{\partial_D \Omega} = 0$, if $\partial_D \Omega \neq \emptyset$ and $\partial_N \Omega$ does not contain any two adjacent hyperfaces.

6.3. Proofs of the main results. In this subsection, we finally tackle the proofs of the main results of the paper. We first show how the proof of the regularity property for the mixed boundary value/interface problem (6), Theorem 1.1 can be obtained from the results of [1] and the theory developed in Section 4. The following result was proved in [1].

Theorem 6.8. *Let $(\mathfrak{M}, \mathcal{V})$ be a Lie manifold with boundary and $P_0 \in \text{Diff}^m(\mathfrak{M})$ be a second order, strongly elliptic operator. Let h be an admissible weight and $u \in hH^1(\mathfrak{M})$ be such that $Pu \in hH^{\mu-1}(\mathfrak{M})$ and $u|_{\partial \mathfrak{M}} \in hH^{\mu+1/2}(\partial \mathfrak{M})$, $\mu \in \mathbb{Z}_+$. Then $u \in hH^{\mu+1}(\mathfrak{M})$ and*

$$(64) \quad \|u\|_{hH^{\mu+1}(\mathfrak{M})} \leq C(\|P_0 u\|_{hH^{\mu-1}(\mathfrak{M})} + \|u\|_{hH^0(\mathfrak{M})} + \|u|_{\partial \mathfrak{M}}\|_{hH^{\mu+1/2}(\partial \mathfrak{M})}).$$

For mixed boundary value/interface problems we need the following extension of this theorem, which is proved exactly in the same way.

Theorem 6.9. *Let $(\mathfrak{M}, \mathcal{V})$ be a Lie manifold with boundary and $P_0 \in \text{Diff}^m(\mathfrak{M})$ be a second order, strongly elliptic operator with jumps discontinuities on sub Lie manifolds of \mathfrak{M} that partition it into subsets \mathfrak{M}_j . Assume that $\partial \mathfrak{M} = \partial_D \mathfrak{M} \cup \partial_N \mathfrak{M}$ is a disjoint decomposition into open, disjoint subsets. Let h be an admissible weight and $u \in hH^1(\mathfrak{M})$ be such that $Pu \in hH^{\mu-1}(\mathfrak{M}_j)$ and $u|_{\partial \mathfrak{M}} \in hH^{\mu+1/2}(\partial \mathfrak{M})$, $\mu \in \mathbb{Z}_+$. Then $u \in hH^{\mu+1}(\mathfrak{M}_j)$ and*

$$\begin{aligned} \|u\|_{hH^{\mu+1}(\mathfrak{M}_j)} \leq C & \left(\sum_k \|P_0 u\|_{hH^{\mu-1}(\mathfrak{M}_k)} + \|u\|_{hH^0(\mathfrak{M})} \right. \\ & \left. + \|u|_{\partial \Omega}\|_{hH^{\mu+1/2}(\partial_D \mathfrak{M})} + \|D_\nu^P u|_{\partial \Omega}\|_{hH^{\mu-1/2}(\partial_N \mathfrak{M})} \right). \end{aligned}$$

Theorem 1.1 then follows by applying the above theorem to $P_0 := r_\Omega^2 P$, which is strongly elliptic by Corollary 6.3(ii), and using the identifications of Proposition 5.8 and Definition 5.9.

We now prove Theorem 1.2 assuming the results stated in the previous subsection. The proof of Theorem 1.3 is completely similar.

Remark 6.10. In the statement of Theorems 1.2 and 1.3, the spaces $\mathcal{K}_1^{\mu+1}(\Omega_j)$ are defined intrinsically, without reference to Ω . However, the interface Γ is assumed smooth for well-posedness in this paper (more general conditions on Γ were for example considered in [35]) and the points where Γ intersects $\partial \Omega$, necessarily transversely, are included in the singular sets $\Omega^{(n-2)}$ of Ω ; consequently, r_Ω is equivalent to r_{Ω_j} in each Ω_j .

Proof. We first notice that Theorem 5.10 allows us to reduce the proof to the case $g_D = 0$.

We continue to denote with $\mathcal{W}_\mu(\Omega)$ the set of weights such that the operator \tilde{P} , defined by $\tilde{P}(u) := (Pu, u|_{\partial_D\Omega}, D_\nu^P u|_{\partial_N\Omega})$, is an isomorphism

$$(65) \quad \tilde{P} : \left\{ \bigoplus_{j=1}^N h\mathcal{K}_1^{\mu+1}(\Omega_j) \cap \mathcal{K}_1^1(\Omega); u|_{\partial_D\Omega} = 0, u^+ = u^-, D_\nu^P u^+ = D_\nu^P u^- \text{ on } \Gamma \right\} \rightarrow \bigoplus_{j=1}^N h\mathcal{K}_{-1}^{\mu-1}(\Omega_j) \oplus h\mathcal{K}_{-1/2}^{\mu-1/2}(\partial_N\Omega),$$

which is an open set by Proposition 6.2. Therefore, it is enough to show that $1 \in \mathcal{W}_\mu(\Omega)$ to complete the proof.

For solvability we consider the case $\mu = 0$. For $\mu = 0$, the problem (6) is interpreted in the weak sense (11), using that $\mathcal{K}_1^1(\Omega) \subset H^1(\Omega)$. More precisely, we let

$$(66) \quad \mathcal{H} := \{u \in \mathcal{K}_1^1(\Omega), u = 0 \text{ on } \partial_D\Omega\},$$

and we define the weak solution u of Equation (11) with $g_D = 0$ as the unique $u \in \mathcal{K}_1^1(\Omega)$ satisfying $u = 0$ on $\partial_D\Omega$ in trace sense and

$$(67) \quad B_P(u, v) = \Phi(v) \quad \text{for all } v \in \mathcal{H}.$$

where the element $\Phi \in \mathcal{H}^*$ is defined by $\Phi(u) = \int_\Omega f u \, dx + \int_{\partial_N\Omega} g_N u \, dS(x)$, this last integral being the pairing $\mathcal{K}_{1/2}^{1/2}(\partial\Omega) - \mathcal{K}_{-1/2}^{-1/2}(\partial\Omega)$. Here, we have employed the trace property, Theorem 5.10. We will establish the existence and uniqueness of u by using the Lax-Milgram Lemma and coercive estimates for P in weighted Sobolev spaces, which in turn follow from the (uniform) strong ellipticity of P and the Hardy-Poincaré inequality of Theorem 6.4. This result gives the first step of the proof, that is, $1 \in \mathcal{W}_0(\Omega)$. We refer to [17] for the version of the Lax-Milgram lemma needed in this proof, where P contains lower-order terms.

Indeed, the sesquilinear form B is continuous on $\mathcal{H} \times \mathcal{H}$ by Proposition 6.1. Furthermore, assumptions 8 on the coefficients A_{jk} , B_j , and C of the operator P , together with Corollary 6.7 imply the following inequality for the real part of $B(u, v)$

$$(68) \quad \begin{aligned} \operatorname{Re}(Pu, u) &= \int_\Omega \left(\operatorname{Re} \sum_{j,k=1}^n A_{jk} \partial_k u \overline{\partial_j u} \right) dx \\ &+ (2C - \sum_j \partial_j B_j)u, u / 2 \geq \epsilon \sum_{j=1}^n \|\partial_j u\|^2 = \epsilon \sum_{p=1}^n \|\nabla u_p\|_{L^2(\Omega)}^2 \\ &\geq \epsilon \|u\|_{\mathcal{K}_1^1(\Omega)}^2 =: \epsilon \|u\|_{\mathcal{K}_1^1(\Omega)}^2, \end{aligned}$$

which shows that B is strictly coercive on \mathcal{H} .

The assumptions of the Lax-Milgram lemma are therefore satisfied, Hence $P : \mathcal{H} \rightarrow \mathcal{H}^*$ is an isomorphism (*i.e.*, P is continuous with continuous inverse). proving that $1 \in \mathcal{W}_0\Omega$.

We next consider the case $\mu \geq 1$ and prove that $\mathcal{W}_0(\Omega) \subset \mathcal{W}_\mu(\Omega)$ for any $\mu \in \mathbb{Z}_+$, so that, in particular, $1 \in \mathcal{W}_\mu(\Omega)$. We pick $h \in \mathcal{W}_0(\Omega)$ and observe that by the regularity theorem, Theorem 1.1, the map \tilde{P} of Equation 65 above is surjective. Since this map is also continuous (Proposition 6.1) and injective

(because $h \in \mathcal{W}_0(\Omega)$), it is an isomorphism by the open mapping theorem. This observation shows that $\mathcal{W}_0(\Omega) \subset \mathcal{W}_\mu(\Omega)$, for any $\mu \in \mathbb{Z}_+$.

Since we have already proven that $\mathcal{W}_\mu(\Omega)$ is open, the proof is complete. \square

Remark 6.11. It seems that it would be more natural to work in the framework of stratified spaces than in the framework of polyhedral domains. For example, if we consider a smooth, bounded domain $\Omega \subset \mathbb{R}^n$ and a submanifold $X \subset \partial\Omega$ of lower dimension, then we can consider $\eta_{n-2}(x)$ to be the distance from x to X . Then Theorem 1.2 remains true, with essentially the same proof, by taking $\Omega^{(n-2)} := X$ in this case.

REFERENCES

- [1] B. Ammann, A. Ionescu, and V. Nistor. Sobolev spaces and regularity for polyhedral domains. Preprint, MSRI, February 2004. New title. Old title: *Sobolev spaces on Lie manifolds and on polyhedral domains*.
- [2] B. Ammann, R. Lauter, and V. Nistor. Pseudodifferential operators on manifolds with a Lie structure at infinity. Preprint math.AP/0304044, December 2002.
- [3] B. Ammann, R. Lauter, and V. Nistor. On the geometry of Riemannian manifolds with a Lie structure at infinity. *Int. J. Math. Math. Sci.*, 2004(1-4):161–193, 2004.
- [4] I. Babuška. Finite element method for domains with corners. *Computing (Arch. Elektron. Rechnen)*, 6:264–273, 1970.
- [5] I. Babuška and B. Q. Guo. The h - p version of the finite element method for domains with curved boundaries. *SIAM J. Numer. Anal.*, 25(4):837–861, 1988.
- [6] Constantin Bacuta. Subspace interpolation with applications to elliptic regularity. *Numer. Funct. Anal. Optim.*, 29(1-2):88–114, 2008.
- [7] C. Băcuță, J.H. Bramble, and J Xu. Regularity estimates for elliptic boundary value problems in Besov spaces. *Math. Comp.*, 72:1577–1595, 2003.
- [8] C. Băcuță, V. Nistor, and L. Zikatanov. A note on improving the rate of convergence of ‘high order finite elements’ on polygons. *Numerische Mathematik*, 100:165–184, 2005.
- [9] M. Cheeger, J. Gromov and M. Taylor. Finite propagation speed, kernel estimates for functions of the Laplace operator, and the geometry of complete Riemannian manifolds. *J. Differential Geom.*, 17(1):15–53, 1982.
- [10] M. Costabel. Boundary integral operators on curved polygons. *Ann. Mat. Pura Appl. (4)*, 133:305–326, 1983.
- [11] M. Dauge. *Elliptic boundary value problems on corner domains*, volume 1341 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1988. Smoothness and asymptotics of solutions.
- [12] M. Dauge. Neumann and mixed problems on curvilinear polyhedra. *Integral Equations Operator Theory*, 15(2):227–261, 1992.
- [13] J. Elschner. The double layer potential operator over polyhedral domains. I. Solvability in weighted Sobolev spaces. *Appl. Anal.*, 45(1-4):117–134, 1992.
- [14] J. Elschner. On the double layer potential operator over polyhedral domains: solvability in weighted Sobolev spaces and spline approximation. In *Symposium “Analysis on Manifolds with Singularities” (Breitenbrunn, 1990)*, volume 131 of *Teubner-Texte Math.*, pages 57–64. Teubner, Stuttgart, 1992.
- [15] A.K. Erkip and E. Schrohe. Normal solvability of elliptic boundary value problems on asymptotically flat manifolds. *J. Funct. Anal.*, 109:22–51, 1992.
- [16] L.C. Evans. *Partial differential equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1998.
- [17] G. Folland. *Introduction to partial differential equations*. Princeton University Press, Princeton, NJ, second edition, 1995.
- [18] P. Grisvard. *Elliptic problems in nonsmooth domains*, volume 24 of *Monographs and Studies in Mathematics*. Pitman (Advanced Publishing Program), Boston, MA, 1985.
- [19] P. Grisvard. *Singularities in boundary value problems*, volume 22 of *Research in Applied Mathematics*. Masson, Paris, 1992.
- [20] G. Grubb. *Functional calculus of pseudodifferential boundary problems*, volume 65 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, second edition, 1996.

- [21] B. Guo and I. Babuška. Regularity of the solutions for elliptic problems on nonsmooth domains in \mathbf{R}^3 . I. Countably normed spaces on polyhedral domains. *Proc. Roy. Soc. Edinburgh Sect. A*, 127(1):77–126, 1997.
- [22] D. Jerison and C. Kenig. The Dirichlet problem in nonsmooth domains. *Ann. of Math. (2)*, 113(2):367–382, 1981.
- [23] D. Jerison and C. Kenig. The Neumann problem on Lipschitz domains. *Bull. Amer. Math. Soc. (N.S.)*, 4(2):203–207, 1981.
- [24] D. Jerison and C.E. Kenig. The inhomogeneous Dirichlet problem in Lipschitz domains. *J. Funct. Anal.*, 130(1):161–219, 1995.
- [25] R. B. Kellogg. Singularities in interface problems. In *Numerical Solution of Partial Differential Equations, II (SYNSPADE 1970) (Proc. Sympos., Univ. of Maryland, College Park, Md., 1970)*, pages 351–400. Academic Press, New York, 1971.
- [26] R. B. Kellogg. On the Poisson equation with intersecting interfaces. *Applicable Anal.*, 4:101–129, 1974/75. Collection of articles dedicated to Nikolai Ivanovich Muskhelishvili.
- [27] C. Kenig. Recent progress on boundary value problems on Lipschitz domains. In *Pseudodifferential operators and applications (Notre Dame, Ind., 1984)*, volume 43 of *Proc. Sympos. Pure Math.*, pages 175–205. Amer. Math. Soc., Providence, RI, 1985.
- [28] C.E. Kenig and T. Toro. Harmonic measure on locally flat domains. *Duke Math. J.*, 87(3):509–551, 1997.
- [29] V. A. Kondrat'ev. Boundary-value problems for elliptic equations in conical regions. *Dokl. Akad. Nauk SSSR*, 153:27–29, 1963.
- [30] V. A. Kondrat'ev. Boundary value problems for elliptic equations in domains with conical or angular points. *Transl. Moscow Math. Soc.*, 16:227–313, 1967.
- [31] V. A. Kondrat'ev. The smoothness of the solution of the Dirichlet problem for second order elliptic equations in a piecewise smooth domain. *Differencial'nye Uravnenija*, 6:1831–1843, 1970.
- [32] V. Kozlov, V. Maz'ya, and J. Rossmann. *Elliptic boundary value problems in domains with point singularities*, volume 52 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1997.
- [33] V. Kozlov, V. Maz'ya, and J. Rossmann. *Spectral problems associated with corner singularities of solutions to elliptic equations*, volume 85 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2001.
- [34] R. Lauter and V. Nistor. Analysis of geometric operators on open manifolds: a groupoid approach. In N.P. Landsman, M. Pflaum, and M. Schlichenmaier, editors, *Quantization of Singular Symplectic Quotients*, volume 198 of *Progress in Mathematics*, pages 181–229. Birkhäuser, Basel - Boston - Berlin, 2001.
- [35] H. Li, A. L. Mazzucato, and V. Nistor. Well-posedness and regularity for the elasticity equation with mixed boundary conditions in polyhedral domains and domains with cracks. Submitted.
- [36] J.-L. Lions and E. Magenes. *Non-homogeneous boundary value problems and applications. Vol. I*. Springer-Verlag, New York, 1972. Translated from the French by P. Kenneth, Die Grundlehren der mathematischen Wissenschaften, Band 181.
- [37] V. Maz'ya, S. Nazarov, and B. Plamenevskij. *Asymptotic theory of elliptic boundary value problems in singularly perturbed domains. Vol. I*, volume 111 of *Operator Theory: Advances and Applications*. Birkhäuser Verlag, Basel, 2000. Translated from the German by Georg Heinig and Christian Posthoff.
- [38] V. Maz'ya, S. Nazarov, and B. Plamenevskij. *Asymptotic theory of elliptic boundary value problems in singularly perturbed domains. Vol. II*, volume 112 of *Operator Theory: Advances and Applications*. Birkhäuser Verlag, Basel, 2000. Translated from the German by Plamenevskij.
- [39] V. Maz'ya and J. Roßmann. Weighted L_p estimates of solutions to boundary value problems for second order elliptic systems in polyhedral domains. *ZAMM Z. Angew. Math. Mech.*, 83(7):435–467, 2003.
- [40] A. L. Mazzucato and V. Nistor. Well-posedness and regularity for the elasticity equation with mixed boundary conditions in polyhedral domains and domains with cracks. Submitted.
- [41] R.B. Melrose. *The Atiyah-Patodi-Singer index theorem*. Research Notes in Mathematics (Boston, Mass.). 4. Wellesley, MA: A. K. Peters, Ltd.. xiv, 377 p. , 1993.

- [42] R.B. Melrose. *Geometric scattering theory*. Stanford Lectures. Cambridge University Press, Cambridge, 1995.
- [43] D. Mitrea, M. Mitrea, and M. Taylor. Layer potentials, the Hodge Laplacian, and global boundary problems in nonsmooth Riemannian manifolds. *Mem. Amer. Math. Soc.*, 150(713):x+120, 2001.
- [44] M. Mitrea and V. Nistor. Boundary layer potentials on manifolds with cylindrical ends. ESI Preprint.
- [45] M. Mitrea and M. Taylor. Boundary layer methods for Lipschitz domains in Riemannian manifolds. *J. Funct. Anal.*, 163(2):181–251, 1999.
- [46] M. Mitrea and M. Taylor. Potential theory on Lipschitz domains in Riemannian manifolds: L^p Hardy, and Hölder space results. *Comm. Anal. Geom.*, 9(2):369–421, 2001.
- [47] S. Nazarov and B. Plamenevsky. *Elliptic problems in domains with piecewise smooth boundaries*, volume 13 of *de Gruyter Expositions in Mathematics*. Walter de Gruyter & Co., Berlin, 1994.
- [48] J. Rossmann. The asymptotics of the solutions of linear elliptic variational problems in domains with edges. *Z. Anal. Anwendungen*, 9(6):565–578, 1990.
- [49] E. Schrohe. Fréchet algebra techniques for boundary value problems: Fredholm criteria and functional calculus via spectral invariance. *Math. Nachr.*, 199:145–185, 1999.
- [50] E. Schrohe and B.-W. Schulze. Boundary value problems in Boutet de Monvel’s algebra for manifolds with conical singularities. I. In *Pseudo-differential calculus and mathematical physics*, volume 5 of *Math. Top.*, pages 97–209. Akademie Verlag, Berlin, 1994.
- [51] E. Schrohe and B.-W. Schulze. Boundary value problems in Boutet de Monvel’s algebra for manifolds with conical singularities II. In *Boundary value problems, Schrödinger operators, deformation quantization*, volume 8 of *Math. Top.* Akademie Verlag, Berlin, 1995.
- [52] M.A. Shubin. Spectral theory of elliptic operators on noncompact manifolds. *Astérisque*, 207:5, 35–108, 1992. Méthodes semi-classiques, Vol. 1 (Nantes, 1991).
- [53] D. A. Stone. *Stratified polyhedra*. Lecture Notes in Mathematics, Vol. 252. Springer-Verlag, Berlin, 1972.
- [54] M. Taylor. *Partial differential equations I, Basic theory*, volume 115 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1995.
- [55] G. Verchota. Layer potentials and regularity for the Dirichlet problem for Laplace’s equation in Lipschitz domains. *J. Funct. Anal.*, 59(3):572–611, 1984.
- [56] G. Verchota and A. Vogel. The multidirectional Neumann problem in \mathbb{R}^4 . Preprint, 2003.
- [57] G.C. Verchota and A.L. Vogel. A multidirectional Dirichlet problem. *J. Geom. Anal.*, 13(3):495–520, 2003.
- [58] H. Whitney. Tangents to an analytic variety. *Ann. of Math. (2)*, 81:496–549, 1965.

UNIVERSITY OF DELAWARE, DEPARTMENT OF MATHEMATICAL SCIENCES, 501 EWING HALL,
NEWARK, DE 19716-2553

E-mail address, C. Băcuță: bacuta@math.udel.edu

PENNSYLVANIA STATE UNIVERSITY, DEPARTMENT OF MATHEMATICS, UNIVERSITY PARK, PA
16802

E-mail address, A. Mazzucato: alm24@psu.edu

E-mail address, V. Nistor: nistor@math.psu.edu

E-mail address, L. Zikatanov: ltz@math.psu.edu