



A Four Vertex Theorem for Polygons

Serge Tabachnikov

The American Mathematical Monthly, Vol. 107, No. 9. (Nov., 2000), pp. 830-833.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9890%28200011%29107%3A9%3C830%3AAAFVTFP%3E2.0.CO%3B2-D>

The American Mathematical Monthly is currently published by Mathematical Association of America.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/maa.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is an independent not-for-profit organization dedicated to creating and preserving a digital archive of scholarly journals. For more information regarding JSTOR, please contact support@jstor.org.

NOTES

Edited by Jimmie D. Lawson and William Adkins

A Four Vertex Theorem for Polygons

Serge Tabachnikov

The classical 4-vertex theorem states that a closed smooth convex plane curve has at least 4 curvature extrema. Published in 1909 [4] this theorem and its various generalizations continue to attract interest; see [1] for a contemporary point of view on the subject.

One can formulate the 4-vertex theorem in terms of the 1-parameter family of normal lines to the curve; the envelope of these lines is called the *caustic* (or the *evolute*) of the curve. The caustic is not smooth: generically, it has semi-cubical cusp singularities, and these cusps correspond to the extrema of the curvature. One gives the smooth pieces of the caustic the orientation induced by the inward orientation of the normals. As one traverses the caustic, the cusps are the points at which the orientation changes to the opposite; see Figure 1. The next two results, due to O. Musin and B. Wegner, respectively, extend the 4-vertex theorem from smooth curves to polygons. Let A_1, \dots, A_n be consecutive vertices of a plane convex polygon P . Consider the circle that is circumscribed around the triangle $A_{i-1}A_iA_{i+1}$, and let R_i be its radius. Assume, in addition, that for every i the center of the circumscribed circle lies inside the angle $A_{i-1}A_iA_{i+1}$. Likewise, let r_i be the radius of the circle touching the lines $A_{i-1}A_i$, A_iA_{i+1} and $A_{i+1}A_{i+2}$ and lying on the same sides of these lines as P ; in other words, the center of this circle is the intersection of the bisectors of the i -th and $(i + 1)$ -st interior angles of P (here and elsewhere we understand the index i cyclically, that is, $n + 1 = 1$, etc.). Then each of the two cyclic difference sequences

$$(\Delta R)_i = R_{i+1} - R_i \quad \text{and} \quad (\Delta r)_i = r_{i+1} - r_i, \quad i = 1, \dots, n$$

changes sign at least 4 times; see [7], [8], and [11]. In the limit $n \rightarrow \infty$, when P approximates a smooth curve, both sequences R_i and r_i approximate the radius of




Figure 1

curvature function, and the places where $(\Delta R)_i$ or $(\Delta r)_i$ change signs become critical points of this function, i.e., vertices of the curve.

We propose the following generalization. Consider a convex polygon P with vertices A_1, \dots, A_n , and let $l_i, i = 1, \dots, n$ be a line through the vertex A_i that intersect the interior of P and is oriented inwards. Denote by α_i and β_i the angles made by l_i with the sides $A_i A_{i+1}$ and $A_i A_{i-1}$, respectively, and assume that $\alpha_i + \beta_{i+1} < \pi$ for all i .

We think of the polygon P as an analog of a smooth curve, so the lines l_i play the role of the normals. Let $B_i = l_i \cap l_{i+1}$; the closed (possibly self-intersecting) polygon $Q = (B_1, \dots, B_n)$ is the analog of the caustic. Each side of Q lies on a line l_i and gets orientation from it. Traverse Q and mark the vertices at which the orientations of the sides change; by analogy with the smooth case, call them *cuspl vertices*; the four cuspl vertices are marked in Figure 2. Of course, the number of cuspl vertices is even.




Figure 2

Call a collection of lines l_i *exact* if $\prod_1^n \sin \alpha_i = \prod_1^n \sin \beta_i$. For example, the collection of the bisectors of the interior angles of P is exact. A collection of lines l_i is called *generic* if no three consecutive lines intersect at one point.

Theorem 1. *For every exact generic collection of lines l_1, \dots, l_n , the polygon Q has at least 4 cuspl vertices.*

If l_i are the bisectors of the angles of P , one obtains Wegner's theorem. Note that without the exactness assumption one can easily do without cuspl vertices at all; see Figure 3.




Figure 3

Proof: Start with a geometrical interpretation of the exactness condition. Pick a point $X_1 \in l_1$, draw a line through X_1 parallel to A_1A_2 until it intersects l_2 at point X_2 , draw a line through X_2 parallel to A_2A_3 until it intersects l_3 at point X_3, \dots ; see Figure 4.




Figure 4

We claim that the polygonal line X_1, X_2, \dots closes (that is, $X_{n+1} = X_1$) if and only if the collection of lines l_i is exact. Indeed, one has: $|X_{i+1}A_{i+1}|/|X_iA_i| = \sin \alpha_i/\sin \beta_{i+1}$; therefore $X_{n+1} = X_1$ if and only if $\prod \sin \alpha_i/\prod \sin \beta_i = 1$.

Translate the vectors $A_i X_i$ to an origin O to obtain a polygon $V_1 \cdots V_n$; see Figure 4. It follows from the argument in the previous paragraph that the sides of this polygon are parallel to those of P . For every i the triangle $B_i A_i A_{i+1}$ is similar to $OV_i V_{i+1}$; denote by t_i the similarity coefficient. As long as the t_i monotonically increase or decrease, the orientation of the sides of the “caustic” Q do not change, and the cusp vertices of Q correspond to the change of sign of the difference sequence $(\Delta t)_i = t_{i+1} - t_i$; more precisely, B_i is a cusp vertex if and only if $(\Delta t)_{i-1}$ and $(\Delta t)_i$ have opposite signs.

It remains to show that the cyclic difference sequence $(\Delta t)_i$ changes sign at least 4 times. Since $t_i(A_{i+1} - A_i) = V_{i+1} - V_i$, we have

$$\sum_1^n (\Delta t)_i A_{i+1} = - \sum_1^n t_i (A_{i+1} - A_i) = - \sum_1^n (V_{i+1} - V_i) = 0,$$

(the first equality is “integration by parts”). Since $\sum (\Delta t)_i = 0$, the cyclic sequence $(\Delta t)_i$ changes sign at least twice. Assume that it happens exactly twice, say, $(\Delta t)_i > 0$ for $i = 1, \dots, k$ and $(\Delta t)_i < 0$ for $i = k + 1, \dots, n$. Choose an origin on a line m that intersects the segments $A_{k+1}A_{k+2}$ and A_1A_2 . With this choice, all of the vectors $(\Delta t)_i A_{i+1}$ lie on the same side of m , and $\sum_1^n (\Delta t)_i A_{i+1}$ cannot vanish. This contradiction proves the theorem. ■

Remarks. 1. The continuous case $n \rightarrow \infty$ of Theorem 1 is discussed in [9] and [10]. For another aspect of exact systems of lines see [6].

2. The argument in the proof of Theorem 1 is a particular case of a discrete version of the following theorem (Sturm, Hurwitz, Kellogg, ...): if a function on a circle is L_2 -orthogonal to a k -dimensional Chebyshev system, then this function

has at least $k + 1$ distinct zeroes; see [1] for a continuous and [3] for a discrete version.

3. One may argue that the oldest 4-vertex type result is the following Cauchy lemma (1813), which plays a crucial role in the proof of convex polyhedra rigidity: given two convex n -gons whose i -th sides have equal lengths for all i , the cyclic sequence of the differences of the corresponding angles of the polygons changes sign at least 4 times; see [2]. Our theorem concerns a dual configuration: two polygons whose corresponding angles are equal; see Figure 4.

4. A general approach to discrete versions of various 4 and 6-vertex theorems can be found in [5].

We conclude with a curious and not immediately obvious property of exact systems of lines.

Lemma . *Exactness is invariant under projective transformations of the plane.*

Proof: Choose a point O inside the polygon P , let l'_i be the line A_iO and denote by α'_i and β'_i the angles made by l'_i with sides A_iA_{i+1} and A_iA_{i-1} , respectively. The new lines l'_i form an exact system: consider the Sine Rule for the triangle $A_iA_{i+1}O$ and take the product over all i . The cross-ratio of the four lines through vertex A_i , namely, the two adjacent sides of P and the lines l_i and l'_i , equals $(\sin \alpha'_i / \sin \alpha_i) / (\sin \beta'_i / \sin \beta_i)$, and the product of these ratios is equal to one.

Apply a projective transformation F to this configuration of lines and points. The lines $F(l'_i)$ are still exact since they pass through a point $F(O)$. The cross-ratios of each quadruple of lines remains the same, and so their product still equals one. It follows that $F(l_i)$ form an exact collection of lines as well. ■

ACKNOWLEDGMENTS. This work was done during the author's visit at Max-Planck-Institute in Bonn; it is a pleasure to thank the Institute for its hospitality.

REFERENCES

1. V. Arnold, Topological problems of wave propagation, *Russian Math. Surveys*, **51** (1996) 3–50.
2. R. Connelly, Rigidity in *Handbook of Convex Geometry*, North-Holland, 1993, pp. 223–272.
3. F. Gantmakher and M. Krein, Sur les matrices complètement non négative et oscillatoires, *Compositio Math.* **4** (1937) 445–476.
4. S. Mukhopadhyaya, New methods in the geometry of a plane arc, *Bull. Calcutta Math. Soc.* **1** (1909) 31–37.
5. V. Ovsienko and S. Tabachnikov, Projective geometry of polygons and discrete 4-vertex and 6-vertex theorems, preprint, 1999.
6. Problem No. 10724, *Amer. Math. Monthly* **106** (1999) 265.
7. V. Sedykh, Theorem of four support vertices of a polygonal line, *Functional Anal. Appl.* **30** (1996) 216–218.
8. V. Sedykh, Discrete versions of four-vertex theorem, *Amer. Math. Soc. Transl. Ser. 2* **180** (1997) 197–207.
9. S. Tabachnikov, The four vertex theorem revisited—two variations on the old theme, *Amer. Math. Monthly* **102** (1995) 912–916.
10. S. Tabachnikov, Parametrized plane curves, Minkowski caustics, Minkowski vertices and conservative line fields, *Enseign. Math.* **43** (1997) 3–26.
11. B. Wegner, On the evolutes of piecewise linear curves in the plane, *Rad. Hrvat. Acad. Znan. Umjet.* **467** (1994) 1–16.

Pennsylvania State University, University Park, PA 16802
tabachni@psu.edu