

On Zeroes of the Schwarzian Derivative

Serge Tabachnikov

MPIM, 53225 Bonn, Germany (tabachnikov@mpim-bonn.mpg.de), and
Department of Mathematics, University of Arkansas, Fayetteville AR 72701, USA

1. The Schwarzian derivative

The main character of the present note is the Schwarzian derivative, and we start with a brief reminder of its definition and main properties.

Let $f : \mathbf{RP}^1 \rightarrow \mathbf{RP}^1$ be a projective line diffeomorphism. For every point $x \in \mathbf{RP}^1$ there exists a unique projective transformation $g_x : \mathbf{RP}^1 \rightarrow \mathbf{RP}^1$ whose 2-jet at x coincides with that of f . The Schwarzian derivative $S(f)$ measures the deviation of the 3-jet $j^3 f$ from $j^3 g_x$.

More specifically, let $x \in \mathbf{RP}^1$ and v be a tangent vector to \mathbf{RP}^1 at x . Extend v to a vector field in a vicinity of x and denote by ϕ_t the corresponding local one-parameter group of diffeomorphisms.

Consider 4 points:

$$x, x_1 = \phi_\epsilon(x), x_2 = \phi_{2\epsilon}(x), x_3 = \phi_{3\epsilon}(x)$$

(ϵ is small) and compare their cross-ratio with that of their images under f . It turns out that the cross-ratio does not change in the first order in ϵ :

$$[f(x), f(x_1), f(x_2), f(x_3)] = [x, x_1, x_2, x_3] + \epsilon^2 S(f)(x) + O(\epsilon^3).$$

The ϵ^2 -coefficient depends on the diffeomorphism f , the point x and the tangent vector v (but not on its extension to a vector field); it is a quadratic function in v . That is to say, $S(f)$, the Schwarzian derivative of a diffeomorphism f , is a quadratic differential on \mathbf{RP}^1 .

By the very construction $S(g) = 0$ if g is a projective transformation, and $S(g \circ f) = S(f)$ if g is a projective transformation and f is an arbitrary diffeomorphism.

Choose a projective coordinate $x \in \mathbf{R}^1 \cup \{\infty\} = \mathbf{RP}^1$. Then a diffeomorphism can be considered as a function of x , and the projective transformations are identified with fraction-linear functions. The Schwarzian derivative is given by the formula

$$S(f) = \frac{f'''}{f'} - \frac{3}{2} \left(\frac{f''}{f'} \right)^2.$$

The Schwarzian derivative enjoys the following cocycle property:

$$S(g \circ f) = S(g)(f')^2 + S(f),$$

which means that S is a 1-cocycle of the group of the projective line diffeomorphisms with the values in quadratic differentials.

Let α be another parameter on \mathbf{RP}^1 , that is, $x = g(\alpha)$ for some function g , and let $f : \mathbf{RP}^1 \rightarrow \mathbf{RP}^1$ be a diffeomorphism. The Schwarzian derivative of f is given, in terms of α , by the formula $S(g \circ f) - S(g)$ where, as before,

$$S(h) = \frac{h'''}{h'} - \frac{3}{2} \left(\frac{h''}{h'} \right)^2,$$

and where prime means now $d/d\alpha$. By the cocycle property,

$$S(g \circ f) - S(g) = ((f')^2 - 1)S(g) + S(f).$$

A particular case of interest is $x = \tan(\alpha/2)$ where $\alpha \in \mathbf{T}^1 = \mathbf{R}/2\pi\mathbf{Z}$ is an angle parameter. Then $S(g) = 1/2$ and the formula for the Schwarzian derivative reads:

$$S(f) = \frac{f'''}{f'} - \frac{3}{2} \left(\frac{f''}{f'} \right)^2 + \frac{1}{2} \left((f')^2 - 1 \right).$$

2. Theorem on zeroes of the Schwarzian derivative

Recently E. Ghys discovered a beautiful theorem ([Gh]; see also [O-T]):

Theorem 2.1. *The Schwarzian derivative of every projective line diffeomorphism has at least 4 distinct zeroes.*

The proof outlined by Ghys is based on the similarity with the classical 4 vertex theorem:

Theorem 2.2.

A smooth closed convex plane curve has at least 4 distinct vertices (curvature extrema).

The latter theorem has the following refinement (see [Bl]):

Theorem 2.3. *If a smooth closed convex plane curve transversally intersects a circle in N points then it has at least N distinct vertices (N is even).*

The analogy between zeroes of the Schwarzian derivative and curvature extrema suggests a refinement of the Ghys theorem. Call a fixed point x of a projective line diffeomorphism f *simple* if the graph of f transversally intersects the diagonal at point (x, x) and $S(f)(x) \neq 0$. A generic diffeomorphism has only simple fixed points. The next theorem is the main observation of the present note.

Theorem 2.4. *If a projective line diffeomorphism has N fixed points of which at least one is simple then its Schwarzian derivative has at least N distinct zeroes.*

Notice that the number of fixed points of a projective line diffeomorphism, counted with multiplicities, is even.

Theorem 2.4 implies the existence of 4 zeroes of the Schwarzian derivative. Given a diffeomorphism f pick 3 generic points $x_1, x_2, x_3 \in \mathbf{RP}^1$, and let g be the projective

transformation that takes each x_i to $f(x_i)$; $i = 1, 2, 3$. Then $g^{-1} \circ f$ has 3 fixed points. The number of fixed points being even, there exists a fourth one. By Theorem 2.4 $S(g^{-1} \circ f)$ has at least 4 zeroes, and so does $S(f) = S(g^{-1} \circ f)$.

3. Proof of Theorem 2.4

The proof is elementary in that it boils down to a (somewhat messy) successive application of Rolle's theorem.

We assume that f is orientation-preserving; otherwise f has exactly 2 fixed points and the assertion of Theorem 2.4 is weaker than that of Theorem 2.1.

Without loss of generality, assume that $\infty \in \mathbf{RP}^1$ is a simple fixed point. The graph G of $f(x)$ is a smooth curve with everywhere positive slope that intersects the diagonal at $(N - 1)$ points.

The following identity is straightforward:

$$((f')^{-1/2})'' = -\frac{1}{2}(f')^{-1/2}S(f).$$

Thus one wants to show that $((f')^{-1/2})''$ has N zeroes.

Rolle's theorem provides quite a few. Between every two consecutive intersections of G with the diagonal there is a point at which $f' = 1$. Hence $(f')^{-1/2}$ assumes the value 1 at least $(N - 2)$ times, and therefore $((f')^{-1/2})''$ vanishes at least $(N - 4)$ times. To account for the missing 4 zeroes we analyse the behaviour of f at infinity.

Let $t = 1/x$ be a local parameter at ∞ ; set $g(t) = 1/f(1/t)$. Then $g(0) = 0$; consider the Taylor expansion

$$g(t) = pt + qt^2 + rt^3 + O(t^4).$$

Since ∞ is a simple fixed point, $p \neq 1$ and $S(g)(0) \neq 0$. It is straightforward that $S(g)(0) = 6(pr - q^2)/p^2$; thus $pr - q^2 \neq 0$.

One obtains the expansion of f at ∞ :

$$f(x) = 1/g(1/x) = ux + v + w\frac{1}{x} + O\left(\frac{1}{x^2}\right)$$

where $u = 1/p$, $v = -q/p^2$ and $w = (q^2 - pr)/p^3$. Thus $u \neq 1$, $w \neq 0$ and G has the asymptotic line $y = ux + v$ - see fig. 1.

Without loss of generality, assume that $u > 1$; otherwise consider f^{-1} instead of f . Let a and b be the x -coordinates of the left and rightmost intersection points of G with the diagonal.

Case 1: $w < 0$ (fig. 1 a).

By the Taylor expansion of f at infinity, $f'(x) > u$, and so $f'(x)^{-1/2} < u^{-1/2}$, for sufficiently great $|x|$. Consider the graph G' of $f'(x)^{-1/2}$.

By Rolle's theorem G' intersects the line $y = 1$ at least $(N - 2)$ times, and this provides $(N - 3)$ extrema of $(f')^{-1/2}$. Also G' lies below the line $y = u^{-1/2}$ for sufficiently great $|x|$, therefore $(f')^{-1/2}$ has a local minimum to the right of the rightmost intersection of G' with $y = 1$ and to the left of the leftmost such intersection - see fig. 2 a. The points

$x = +\infty$ and $x = -\infty$ also qualify as extrema of $(f')^{-1/2}$. Altogether $(f')^{-1/2}$ has $(N + 1)$ critical points, and by Rolle's theorem $((f')^{-1/2})''$ has N zeroes.

Case 2: $w > 0$ (fig. 1 b).

In this case $f'(x) < u$, and so $f'(x)^{-1/2} > u^{-1/2}$, for sufficiently great $|x|$. Either $a < v/(1 - u)$ or $b > v/(1 - u)$ (or both). Consider the former case, the latter being completely analogous.

For x sufficiently close to $-\infty$ the graph G lies below the asymptotic line $y = ux + v$. Therefore G must intersect the asymptotic line to the left of the vertical line $x = a$. By Rolle's theorem there exists $c < a$ such that $f'(c) = u$, and so $f'(c)^{-1/2} = u^{-1/2}$ – see fig. 2 b.

We do a similar bookkeeping of critical points of $(f')^{-1/2}$. The graph G' intersects the line $y = 1$ at least $(N - 2)$ times on the interval $a \leq x \leq b$; let d be the x -coordinate of the leftmost such intersection. By Rolle's theorem $(f')^{-1/2}$ has $(N - 3)$ critical points on the interval $[d, b]$.

In addition $(f')^{-1/2}$ has a local maximum on the interval $(-\infty, c)$ and a local minimum on the interval $[c, d]$. The points $\pm\infty$ are critical points of $(f')^{-1/2}$ as well. Altogether $(f')^{-1/2}$ has $(N + 1)$ extrema, thus $((f')^{-1/2})''$ has N zeroes.

4. Infinitesimal version of Theorem 2.4

Let α be the angle parameter on \mathbf{RP}^1 introduced at the end of Section 1. Consider a diffeomorphism close to identity: $f(\alpha) = \alpha + \epsilon g(\alpha)$ where g is a 2π -periodic function and ϵ is small. Then, according to the last formula of Section 1,

$$S(f) = \epsilon (g''' + g') + O(\epsilon^2).$$

Thus the differential operator $(d/d\alpha)^3 + (d/d\alpha)$ is a linearization of the Schwarzian derivative.

Fixed points of f are zeroes of g , and zeroes of $S(f)$ are, in the first approximation, zeroes of $g''' + g'$.

Let $g(\alpha) = 0$. Call α a *nondegenerate* zero if $g'(\alpha) \neq 0$ and $g'''(\alpha) + g'(\alpha) \neq 0$. The next statement is an infinitesimal version of Theorem 2.4.

Theorem 4.1. *If a smooth function g on the circle \mathbf{T}^1 has N zeroes of which at least one is nondegenerate then $g''' + g'$ has at least N distinct zeroes.*

We only outline the proof. Consider g as a function of $x = \tan(\alpha/2)$. Make use of the following identity between differential operators:

$$\left(\frac{d}{d\alpha}\right)^3 + \left(\frac{d}{d\alpha}\right) = \frac{1}{8}(1+x^2)^2 \left(\frac{d}{dx}\right)^3 (1+x^2).$$

Thus one wants to show that $((1+x^2)g(x))'''$ has N zeroes. Assume that π is a simple zero of $g(\alpha)$. One shows that an expansion at infinity holds:

$$(x^2 + 1)g(x) = ux + v + w\frac{1}{x} + O\left(\frac{1}{x^2}\right).$$

where $u \neq 0$ and $w \neq 0$. Therefore the graph of $(1 + x^2)g(x)$ has an asymptotic line and intersects the x -axis $(N - 1)$ times.

The rest of the argument repeats the proof of Theorem 1 and we do not dwell on it.

Remark. $(d/d\alpha)^3 + (d/d\alpha)$ is a *disconjugate* differential operator on the circle. For an operator of odd degree k this means that every function in its kernel is 2π -periodic and has at most $(k - 1)$ zeroes on \mathbf{T}^1 , multiplicities counted. Could Theorem 2 be generalized to other disconjugate operators? For a disconjugate differential operator L of degree k on an interval I the following holds: *if f has N zeroes on I then $L(f)$ has at least $N - k$ zeroes on I* (see [Po]). The proof consists in factorizing L into a product

$$v_0(x) \frac{d}{dx} v_1(x) \frac{d}{dx} v_2(x) \dots v_{k-1}(x) \frac{d}{dx} v_k(x),$$

where each function $v_i(x)$ is positive on I , and successively applying Rolle's theorem. Such a factorization is not available on the circle.

5. Application to vertices of plane curves

The relation between Theorems 2.4 and 4.1 is another manifestation of the relation between zeroes of the Schwarzian derivative and curvature extrema of plane curves. We show here that Theorem 4.1 implies Theorem 2.3.

Let C be the circle that transversely intersects a smooth convex closed plane curve γ at N points. Denote by O the center and by r the radius of C . Let $f(\alpha)$ be the *support function* of γ , that is, the (signed) distance from O to an oriented tangent line to γ as a function of the direction of this line.

Support functions of circles are linear combinations of first harmonics: $a + b \cos \alpha + c \sin \alpha$. These functions generate the kernel of the differential operator $(d/d\alpha)^3 + (d/d\alpha)$. A vertex of a curve is its third order tangency with a circle. Therefore vertices of γ are zeroes of the function $f'''(\alpha) + f'(\alpha)$.

Lemma 5.1. *There exists at least N common support lines to γ and C .*

Proof. Consider an arc δ of the curve that lies outside C and is bounded by two consecutive intersection points $x, y \in \gamma \cap C$ – see fig.3. There are at least $N/2$ such arcs.

The distance function from O assumes maximum on δ at some point z . The support line to γ at z does not intersect C , while the support line at x does. It follows that there is a point on the arc xz such that the support line at this point is tangent to C . Likewise such a point exists on the arc yz . Altogether one gets $2 \cdot N/2 = N$ common support lines.

Lemma 5.1 implies that the function $f(\alpha)$ assumes the value r at least N times, so the function $g(\alpha) = f(\alpha) - r$ has N zeroes. Making a small perturbation of the circle C , if necessary, one may assume that these zeroes are nondegenerate. By Theorem 4.1 $g''' + g'$ has N zeroes, and so does $f''' + f'$. Thus γ has at least N vertices.

6. Lorentz caustics and wave fronts

According to E. Ghys ([Gh]), the theorem on 4 zeroes of the Schwarzian derivative is a 4-vertex theorem in *Lorentzian geometry*. In this last section we discuss this viewpoint.

As before, G denotes the graph of an orientation-preserving projective line diffeomorphism $f(x)$. The graphs of (orientation-preserving) fraction-linear functions are either straight lines or hyperbolas $(x - a)(y - b) = -s^2$; in the latter case $|s|$ is called the *radius* of the hyperbola. Abusing the language we will call these curves simply "hyperbolas".

For every point of G there exists a unique hyperbola second order tangent to G at this point; call it the *osculating hyperbola*. Zeroes of $S(f)$ are moments of third order tangency between G and its osculating hyperbola.

Following Ghys, introduce the Lorentz metric $g = dx dy$ in the (x, y) plane. Hyperbolas are circles in this metric. The graph G is a space-like curve. All geometrical terms such as "length", "orthogonality", etc, refer to the metric g . Let J be the linear operator $(x, y) \rightarrow (-x, y)$. Then $J(v)$ is orthogonal to v , and $g(J(u), J(v)) = -g(u, v)$ for every vectors u and v .

Let t be a (Lorentz) arc-length parameter along G . Denote d/dt by dot and d/dx by prime. One easily finds that

$$\ddot{G}(t) = \frac{1}{2} f''(x(t)) f'(x(t))^{-3/2} J(\dot{G}(t)).$$

Call the coefficient $k = f''(f')^{-3/2}/2$ the (Lorentz) *curvature* of G . Then $\dot{k} = (f')^{-1/2}S(f)$ (the observation due to L. Guieu and V. Ovsienko). Thus curvature extrema are zeroes of the Schwarzian derivative.

Definition. The envelope Γ of g -orthogonal lines to G is called its *Lorentz caustic*.

Note that Γ is a time-like curve. Note also that at inflection points of G the caustic goes to infinity. If G is the graph of a projective line diffeomorphism with a fixed point at infinity then, according to the Moebius theorem, G has at least 3 inflections.

Definition. A *wave front* of G is obtained by moving every point $x \in G$ the same distance in the direction g -orthogonal to G at x .

If $G(t)$ is an arc-length parameterized curve then its s -front is the curve

$$G_s(t) = G(t) - s J(\dot{G}(t)).$$

Example. Let G be the upper branch of the hyperbola $xy = -s^2, s > 0$. The arc-length parameterization is

$$G(t) = (-e^{-t/s}, s^2 e^{t/s}).$$

The curvature is constant: $k = 1/s$ (and $k = -1/s$ on the lower branch of the same hyperbola). The acceleration vector $\ddot{G}(t)$ is collinear to $G(t)$ for all t , so all normals pass through the origin. The caustic degenerates to a point, the center of the hyperbola. The wave fronts are concentric hyperbolas, and in particular, G_s degenerates to the center of the original hyperbola G .

Lorentz caustics and fronts enjoy the familiar properties of their Euclidean counterparts. We summarize them in the next theorem.

Theorem 6.1. a). The Lorentz caustic Γ of a curve G is the locus of its osculating hyperbolas' centers. Γ is also the locus of the singularities of the fronts G_s .
b). Singularities of Γ correspond to curvature extrema of G . If G is the graph of a projective line diffeomorphism f then singularities of Γ correspond to zeroes of $S(f)$.
c). (Huygens' Principle). The front G_s is the envelope of the hyperbolas of radii s with the centers at points of G . The caustic of each front G_s coincides with the caustic of G . The fronts enjoy the evolution property: the s -front of the t -front of G coincides with the $(s + t)$ -front G_{s+t} .
d). Each front G_s is the locus of free ends of a stretched string of constant Lorentz length unwinding from Γ .

Proof. To prove the assertions a) and b) approximate G by its osculating hyperbolas and use the properties of hyperbolas described in the Example.

To prove c) consider 3-dimensional contact space of space-like contact elements in the Lorentz plane with coordinates (x, y, u) and the contact 1-form $\lambda = (e^u dx - e^{-u} dy)/2$. A space-like curve in the (x, y) plane lifts to contact space as a Legendrian curve satisfying $2u = \log(dy/dx)$.

Let $\xi = e^{-u} d/dx - e^u d/dy$ be the Reeb vector field, that is, $\lambda(\xi) = 1$ and $i_\xi d\lambda = 0$. The flow of ξ parallel translates a contact element with a constant velocity in the direction of its Lorentz normal. Denote by $\{\phi_s\}$ the corresponding 1-parameter group of diffeomorphisms. Each ϕ_s preserves the contact structure (and even the contact form).

The s -front of a curve G is obtained as follows: lift G to contact space, apply ϕ_s to the lifted curve and project back to the plane. The evolution property follows. It follows also that all the fronts have the same family of orthogonal lines, therefore they share the same caustic.

To prove that the front G_s is the envelope of the hyperbolas of radii s centered at points of G , consider a point $x \in G$. Its lift to contact space is the curve γ of all space-like contact elements with the base-point x . The lift \tilde{G} of G intersects γ , so $\phi_s(\tilde{G})$ intersects $\phi_s(\gamma)$. The projection of $\phi_s(\gamma)$ to the plane is the hyperbola of radius s with the center at x . Since $\phi_s(\tilde{G})$ intersects $\phi_s(\gamma)$ this hyperbola is tangent to G_s .

To prove d) let t be the arc-length parameter along Γ , i.e., $g(\dot{\Gamma}(t), \dot{\Gamma}(t)) = -1$. Then $\ddot{\Gamma}(t)$ is orthogonal to $\dot{\Gamma}(t)$. The free end of a string unwinding from Γ is the point

$$X(t) = \Gamma(t) + (c - t)\dot{\Gamma}(t)$$

where c is a constant, "length of the string". Then $\dot{X}(t) = (c - t)\ddot{\Gamma}(t)$, which is orthogonal to $\dot{\Gamma}(t)$. Therefore the normals to the curve X are tangent to Γ , that is, X is a front of G .

Remark. The above contact space is contactomorphic to the standard contact space of 1-jets $J^1\mathbf{R}^1$; indeed $\lambda = dq - pdu$ where $p = e^u x + e^{-u} y$ and $q = e^u x - e^{-u} y$.

Acknowledgements. I am grateful to S. Lvovsky, V. Ovsienko, G. Thorbergsson and M. Umehara for numerous stimulating discussions. It is a pleasure to acknowledge the hospitality of the Max-Planck-Institut in Bonn. The research was supported in part by an NSF grant.

7. Appendix: Barner's theorem with applications

After the completion of the present paper G. Thorbergsson attracted my attention to the paper by M. Barner [Ba]. The main result of this paper implies Theorem 2.4, answers in the affirmative the question asked in Remark in Section 4 and provides alternative proofs of some results on flattening points of space curves obtained in [A 1]. Since the paper [Ba] is not sufficiently well known and the journal, in which it was published, is not easily accessible, I reproduce it's main theorem with a (somewhat shortened) proof and discuss some applications here.

Definition. A curve $\gamma \subset \mathbf{RP}^n$ is called *strictly convex* if for every $(n - 1)$ -tuple of points of γ there exists a hyperplane through these points that does not intersect γ anymore.

Remarks. 1). This definition does not exclude infinitely close points. In particular, for every point $x \in \gamma$ there exists a hyperplane through x that contains the osculating codimension 2 space of γ at x and does not intersect the curve anymore.

2). A closed strictly convex curve is contractible if n is odd and noncontractible if n is even.

3). The following definition is given in [A 1]: a curve $\gamma \subset \mathbf{RP}^n$ is called *convex* if every hyperplane intersects γ at at most n points, multiplicities counted. The relation between these two notions of convexity is as follows: a curve in \mathbf{RP}^n , whose projection to \mathbf{RP}^{n-1} from a point convex, is strictly convex.

Definition. An inflection (or flattening) point of a space curve is a stationary point of the osculating hyperplane.

Theorem 7.1. *If a strictly convex closed curve $\gamma \subset \mathbf{RP}^n$ transversely intersects a hyperplane in k points then it has at least k distinct inflections.*

Proof. Induction in n . The case of $n = 1$ is the following statement: if γ is a curve in \mathbf{RP}^1 that does not pass through some point and passes through another point k times then γ has k singularities. This follows from the Rolle theorem.

Consider the induction step from $n - 1$ to n . Let x be one of the intersection points with the hyperplane, and let π be the projection $\mathbf{RP}^n \rightarrow \mathbf{RP}^{n-1}$ from x . Let $\Gamma = \pi(\gamma)$.

Claim 1: Γ is strictly convex.

Indeed, given $y_1, \dots, y_{n-2} \in \Gamma$, consider their preimages $x_i = \pi^{-1}(y_i)$. There exists a hyperplane in \mathbf{RP}^n through points x, x_1, \dots, x_{n-2} that does not intersect γ anymore, and its projection to \mathbf{RP}^{n-1} is the desired hyperplane therein.

Since Γ intersects a hyperplane $(k - 1)$ times, it has $(k - 1)$ inflections.

Claim 2: An inflection of Γ corresponds to an osculating hyperplane of γ passing through x .

Indeed, an inflection of Γ means that n its infinitely close points belong to a hyperplane. Upstairs one has n infinitely close points of γ generating a hyperplane that contains x . This hyperplane is an osculating hyperplane of γ .

Let $0 \leq t \leq 1$ be a parameter on γ , such that $x = \gamma(0) = \gamma(1)$ and $0 \leq t_1 < \dots < t_{k-1} < 1$ are the inflection times of Γ .

Claim 3: If x is not an inflection point of γ then $t_1 \neq 0$.

Indeed, if x is not an inflection, the germ of γ at x is $(1 : t : t^2 : \dots : t^n)$, therefore the corresponding germ of Γ is $(1 : t : t^2 : \dots : t^{n-1})$.

Claim 4: Each interval $(0, t_1), (t_1, t_2), \dots, (t_{k-2}, t_{k-1}), (t_{k-1}, 1)$ contains an inflection of γ .

Let $C(t)$ be the osculating codimension 2 space of γ . Fix one of the above intervals and consider the 2-dimensional vector bundle E over the segment of γ which is the quotient of its normal bundle by $C(t)$. The bundle E has two line subbundles: L_1 , generated by the hyperplane through $C(t)$ that does not meet γ anymore (see Remark 1 above), and L_2 , generated by the hyperplane through x and $C(t)$. By strict convexity L_1 is complimentary to L_2 everywhere.

One also has the line subbundle L_0 in E , generated by the osculating hyperplane of γ . By strict convexity L_0 is complimentary to L_1 everywhere, and at the end points of the interval one has $L_0 = L_2$. This was stated in Claim 2 for each t_i ; it also holds at 0: the osculating hyperplane at x is generated by $C(\epsilon)$ and x as ϵ goes to zero. It follows that somewhere on the interval L_0 is stationary, and this is the desired inflection of γ .

Corollary 7.2. *A strictly convex curve $\gamma \subset \mathbf{RP}^n$ has at least $(n + 1)$ inflections.*

Proof. Consider the hyperplane through n generic points of γ . If n is odd γ is contractible (see Remark 2 above), so there exists another intersection point with this hyperplane. The same holds when n is even, and Theorem 7.1 applies.

Barner gives some applications to vertices of convex curves. One concerns (equi)affine geometry. Recall that a point of a plane curve is called *sextactic* (or an *affine vertex*) if there exists a conic which is fifth order tangent to the curve at this point.

Corollary 7.3. *If a convex closed plane curve γ transversely intersects a conic in $2k$ points then it has $2k$ sextactic points.*

Proof. Consider the Veronese map $V : \mathbf{RP}^2 \rightarrow \mathbf{RP}^5$ of degree 2:

$$V : (x_0 : x_1 : x_2) \rightarrow (x_0^2 : x_1^2 : x_2^2 : x_0x_1 : x_1x_2 : x_2x_0).$$

First note that inflections of the curve $V(\gamma)$ correspond to sextactic points of γ (cf. [A 2]). Next, $V(\gamma)$ is strictly convex: 4 points on γ determine 2 lines, that is, a degenerate conic, which do not intersect γ anymore. If γ intersects a conic $2k$ times then $V(\gamma)$ intersects the corresponding hyperplane $2k$ times, and the result follows.

Another application mentioned in [Ba] is the above Theorem 2.3: its proof repeats the previous argument with the Veronese map $V : \mathbf{RP}^2 \rightarrow \mathbf{RP}^3$:

$$V : (x_0 : x_1 : x_2) \rightarrow (x_0^2 : x_0x_1 : x_0x_2 : x_1^2 + x_2^2).$$

Next, Barner's theorem provides an affirmative answer to the question in remark in Section 4.

Corollary 7.4. *Let L be a disconjugate differential operator on the circle and f be a function. If f has N simple zeroes then $L(f)$ has N zeroes as well.*

Proof. Choose a basis ϕ_1, \dots, ϕ_k in the kernel of L , and consider the curve $\gamma = (f : \phi_1 : \dots : \phi_k) \subset \mathbf{RP}^k$. Since L is disconjugate, the projection of γ from the point $(1 : 0 : \dots : 0)$

is a convex curve. Therefore γ is strictly convex (see Remark 3 above), and it has N intersections with the hyperplane $x_0 = 0$. Inflections of γ correspond to zeroes of $L(f)$ (cf. [A 1]), and the result follows.

In particular, Corollary 7.2 implies that for every function f on the circle and a disconjugate differential operator L of order k the function $L(f)$ has at least $(k + 1)$ zeroes – the result proved differently in [A 1] and [G-M-O].

Finally we show how Barner’s theorem implies our Theorem 2.4 (this application is not in [Ba]).

Let $\gamma \subset \mathbf{RP}^1 \times \mathbf{RP}^1$ be the graph of a projective line diffeomorphism f . Consider the Segre map $S : \mathbf{RP}^1 \times \mathbf{RP}^1 \rightarrow \mathbf{RP}^3$ of degree 2 (cf. [A 2]):

$$S : ((x_0 : x_1), (y_0 : y_1)) \rightarrow (x_0y_0 : x_1y_0 : x_0y_1 : x_1y_1).$$

A plane in \mathbf{RP}^3 corresponds to the graph of a projective transformation, to a vertical, to a horizontal, or to the wedge of a vertical and a horizontal projective lines in $\mathbf{RP}^1 \times \mathbf{RP}^1$. Therefore inflection points of the curve $S(\gamma)$ correspond to zeroes of the Schwarzian derivative $S(f)$.

Next, the curve $S(\gamma)$ is strictly convex. Indeed, given two points A and B on γ , consider the union of the vertical projective line through A and the horizontal projective line through B . This ”cross” does not intersect γ anymore, and strict convexity follows.

Finally, if the diffeomorphism f has N fixed points then its graph intersects the graph of the identity N times. Thus the curve $S(\gamma)$ intersects the corresponding plane N times, and by Barner’s theorem, has N inflections.

References

- [A 1] V. Arnold. On the number of flattening points of space curves. AMS Transl., v. 171 (1996), 11-22.
- [A 2] V. Arnold. Remarks on the extactic points of plane curves. Preprint.
- [Ba] M. Barner. Über die Mindestanzahl stationärer Schmiegebene bei geschlossenen streng-konvexen Raumkurven. Abh. Math. Sem. Univ. Hamburg, 20 (1956), 196-215.
- [Bl] W. Blaschke. Kreis und Kugel, Leipzig 1916.
- [Gh] E. Ghys. Circles osculateurs et géométrie lorentzienne. Talk at journée inaugurale du CMI, Marseille, February 1995.
- [G-M-O] L. Guieu, E. Murre, V. Ovsienko. Theorem on six vertices of a plane curve via the Sturm theory. To appear in Arnold-Gelfand Mathematical Seminars, Birkhauser, 1996.
- [O-T] V. Ovsienko, S. Tabachnikov. Sturm theory, Ghys theorem on zeroes of the Schwarzian derivative and flattening of Legendrian curves. Selecta Mathematica, New Series, to appear.
- [Po] G. Polya. On the mean-value theorem corresponding to a given linear homogeneous differential equation. Trans. AMS, 24 (1922), 312–324.