

Existence and non-existence of skew branes

Serge Tabachnikov

*Department of Mathematics, Penn State
University Park, PA 16802, USA*

Abstract

A skew brane is a codimension 2 submanifold in affine space such that the tangent spaces at any two distinct points are not parallel. We prove that if an oriented closed manifold has a non-zero Euler characteristic χ then it is not a skew brane; generically, the number of oppositely oriented pairs of parallel tangent spaces is not less than $\chi^2/4$. We give a version of this result for immersed submanifolds in codimension 2. We construct examples of skew spheres of arbitrary odd dimensions, generalizing the construction of skew loops in 3-dimensional space due to Ghomi and Solomon. We conclude with two conjectures that are theorems in 1-dimensional case.

1 Introduction

In mid-1960s, H. Steinhaus conjectured that every closed smooth curve in 3-dimensional space has parallel tangent lines. Shortly after that, B. Segre [14, 15] constructed examples of curves without parallel tangent lines but showed that no such curve can lie on the sphere (and therefore, due to the affine invariance of the problem, on an ellipsoid). Call a curve without parallel tangents a *skew loop*. Geometrical and topological study of skew loops and their multi-dimensional analogs has become an active research topic.

Let us briefly survey known results. The supply of skew loops is abundant: for example, every knot type in \mathbf{R}^3 can be realized by a skew loop [24], and so can every homotopy class of loops in a flat 3-torus [17]. The aversion of skew loops to ellipsoids was extended to convex quadratic surfaces in [6] and to non-convex ones in [22], see also [5, 16] for C^1 (as opposed to C^2) versions of these results. Ghomi and Solomon [6] proved a converse statement: if a convex surface is not quadratic then it carries a skew loop. See [18, 19] for generalizations of the Blaschke theorem that convex surfaces whose plane sections are centrally-symmetric must be quadratic. Another non-existence result is proved in [22]: no skew loops lie on a ruled developable disc.

A multi-dimensional version of a skew loop is called a *skew brane*.¹ A skew brane $f : M^n \rightarrow \mathbf{R}^{n+2}$ is an immersion such that the tangent spaces

¹The terms “skew loop” and “skew brane” were coined by M. Ghomi and B. Solomon; see

$df(T_x M)$ and $df(T_y M)$ are not parallel for all $x \neq y$. Pairs of parallel tangent spaces correspond to self-intersections of the image of the tangent Gauss map $M \rightarrow G_n(n+2)$ in the Grassman manifold of n -dimensional subspaces in \mathbf{R}^{n+2} . Since $\dim G_n(n+2) = 2n$, these self-intersections generically occur in isolated points and cannot be destroyed by small perturbations of f . If M is oriented, one distinguishes the cases when parallel tangent spaces have the same or the opposite orientations; we refer to them as positively and negatively parallel tangent spaces.

The aversion of skew loops to quadric surfaces extends to skew branes: no skew brane can lie on a quadratic hypersurface of any signature [22, 16]; for spheres, this was proved in [23]. The paper by Lai [10] has been overlooked in the more recent literature on the subject. Apparently unaware of the work of Segre, Lai provides an example of a skew loop in \mathbf{R}^3 and of a torus $T^2 \subset \mathbf{R}^4$ which is free from pairs of negatively parallel tangent planes. The main result of [10] is the following theorem on non-existence of skew branes.

Theorem 1 *Let $M^{2n} \subset \mathbf{R}^{2n+2}$ be a closed oriented embedded submanifold with non-zero Euler characteristic χ . Then there exists a pair of distinct points $x, y \in M$ such that the tangent spaces $T_x M$ and $T_y M$ are negatively parallel. For a generic submanifold M , the number of such (unordered) pairs (x, y) is not less than $\chi^2/4$.*

We give a streamlined proof of this theorem in Section 2. By a generic submanifold we mean the one whose Gauss map is an immersion with transversal self-intersections; see Section 2 for a justification.

The papers by Lai [9, 10] continue the work by Blaschke [2] and Chern and Spanier [3] concerning embedded surfaces in \mathbf{R}^4 . In this case, we add to Theorem 1 the following result. Let $M \subset \mathbf{R}^4$ be a closed oriented immersed surface of genus g . Assume that M is generic in the sense that self-intersection of M are transversal. Then each double point is assigned a sign; denote by d the algebraic number of double points.

Theorem 2 *M has at least $|d^2 - (1 - g)^2|$ pairs of negatively parallel tangent planes.*

In Section 3 we construct examples of skew branes. Let \mathbf{R}^{2n} be a subspace in \mathbf{R}^{2n+1} and $S^{2n-1} \subset \mathbf{R}^{2n}$ the unit sphere. We consider this sphere as a codimension 2 submanifold in $2n + 1$ -dimensional space.

Theorem 3 *There exists a small perturbation of S^{2n-1} in \mathbf{R}^{2n+1} that makes it into a skew brane.*

For $n = 1$, a much stronger result is proved in [6]: given a non-centrally symmetric smooth closed curve γ in the horizontal plane, there exists a skew loop

[5, 7, 20, 21] for recent study of other classes of non-degenerate embeddings of manifolds into affine and projective spaces.

on the cylinder over γ that projects diffeomorphically on γ (“cylinder lemma”). We believe that an analog of the cylinder lemma holds for odd-dimensional spheres; see Conjecture 4.1 and a more general Conjecture 4.2 in Section 4.

A preliminary version of this paper was posted as preprint math.DG/0504484. More details can be found in Yu. Tyurina’s Ph.D. thesis (Penn State, 2005).

Acknowledgments. I am grateful to D. Fuchs, M. Ghomi and especially B. Solomon for their interest, help and criticism. The author was partially supported by an NSF grant.

2 Topological obstructions to the existence of skew branes

In this section we prove Theorems 1 and 2. The proof of the former in [10] is based on a detailed analysis of the Schubert cell decomposition of Grassmanian manifolds made in [9]; our proofs use characteristic classes of vector bundles and are more straightforward.

Consider the tangent Gauss map $F : M^{2n} \rightarrow G_{2n}^+(2n+2)$ to the Grassman manifold of oriented $2n$ -dimensional subspaces in \mathbf{R}^{2n+2} . Let σ be the involution of the Grassmanian that inverts the orientation of every $2n$ -dimensional subspace. The proofs consist of computing the homology class $F_*[M]$ in the group $H_{2n}(G_{2n}^+(2n+2), \mathbf{Z})$ where $[M]$ is the fundamental class of M , and of the homology intersection number $F_*[M] \cap \sigma_* F_*[M]$. We suppress the coefficients from the notation of homology and cohomology.

Let p and q be even, and consider the Grassmanian $G_p^+(p+q)$. Denote by ξ^p and ν^q the tautological vector bundles: the former has the oriented p -plane E , and the latter its orthogonal complement, as the fiber over E , considered as a point in $G_p^+(p+q)$. The bundles are oriented and so is the Grassmanian. The tangent bundle is as follows: $TG_p^+(p+q) = \text{Hom}(\xi, \nu)$. Denote the fundamental class by $c \in H_{pq}(G_p^+(p+q))$ and let $x = e(\xi) \in H_p(G_p^+(p+q))$ and $y = e(\nu) \in H_q(G_p^+(p+q))$ be the Euler classes. Denote by dot the pairing between homology and cohomology classes.

For a vector $e \in \mathbf{R}^{p+q}$, let $N_e \subset G_p^+(p+q)$ be the set of p -planes orthogonal to e , and let $P_e \subset G_p^+(p+q)$ be the set of p -planes that contain e .

Remark 2.1 Let $M^m \subset \mathbf{R}^{m+2}$ be a closed immersed manifold and $e \in \mathbf{R}^{m+2}$ an arbitrary non-zero vector. Then the Gauss image $F(M) \subset G_m^+(m+2)$ intersects N_e , and for a generic e , in at least $\text{rk } H_*(M)$ points. Indeed, the tangent space $T_x M$ is orthogonal to e if and only if the restriction of the “height” function $f(y) = y \cdot e$ on M has a critical point at x . Then Morse theory implies the above claim. In contrast, the Gauss image $F(M)$ can be disjoint from the set $P_e \subset G_m^+(m+2)$: for example, this is the case when M lies in a hyperplane and the vector e is orthogonal to this hyperplane.

Now we start the proofs of Theorems 1 and 2.

Lemma 2.1 *One has: $c \cdot x^q = 2$, $c \cdot y^p = 2(-1)^{pq/4}$, $xy = 0$.*

Proof. The homological Euler class of an oriented vector bundle η over an oriented manifold M can be found as follows: choose a generic section of η and let $N \subset M$ be the zero locus of this section. Then N is a submanifold, oriented as the intersection of this section and the zero section. The Euler class $e(\eta)$ is dual to the homology class $[N]$.

Fix a vector $e \in R^{p+q}$ and project it to each p -plane. This gives a section of ξ whose zero locus is N_e ; one has: $N_e = G_p^+(p+q-1)$. The class $[N_e] \cap \dots \cap [N_e]$ ($q-1$ times) is dual to x^{q-1} ; it is represented by the set Q consisting of p -planes that are orthogonal to fixed $q-1$ vectors. Hence $Q = G_p^+(p+1) = S^p$. One has: $c \cdot x^q = [Q] \cdot x$, which is the Euler number of the restriction of ξ on Q . The latter is the tangent bundle of the sphere, and its Euler number is 2.

Consider the map $d : G_p^+(p+q) \rightarrow G_q^+(p+q)$ that takes an oriented p -plane to its orthogonal complement. Denote by ξ' and ν' the tautological bundles over $G_q^+(p+q)$ and x', y' their Euler classes. Then $d^*(\xi') = \nu$, $d^*(\nu') = \xi$ and $d^*(x') = y$, $d^*(y') = x$. Let c' be the fundamental class of $G_q^+(p+q)$.

Claim: $d_(c) = (-1)^{pq/4}c'$.*

Assuming this claim and interchanging p and q , one has:

$$2 = c' \cdot (x')^p = (-1)^{pq/4}d_*(c) \cdot (x')^p = (-1)^{pq/4}c \cdot d^*(x')^p = (-1)^{pq/4}c \cdot y^p,$$

as stated.

To prove that $xy = 0$, fix a vector $f \in R^{p+q}$ and, for every p -plane E , project f to E^\perp . This gives a section of ν whose zero locus is P_f . The homology class, dual to xy , is $[N_e \cap P_f]$. If f is not orthogonal to e then this intersection is empty, and hence $xy = 0$.

It remains to prove the above claim. One has:

$$d^*(TG_q^+(p+q)) = d^*(\text{Hom}(\xi', \nu')) = \text{Hom}(\nu, \xi).$$

The bundles $\text{Hom}(\xi, \nu)$ and $\text{Hom}(\nu, \xi)$ are naturally isomorphic, and we need to find whether this isomorphism preserves orientations.

The orientation of the former bundle is determined by the identification of a $p \times q$ matrix with the pq -vector formed by its first, second, etc., rows. Likewise, for the latter bundle, one considers the pq -vector formed by its first, second, etc., columns. A matrix element (i, j) has positions $(i-1)p + j$ in the first and $(j-1)q + i$ in the second vector. Thus we need to find the sign of the permutation of pq elements

$$\tau : (i-1)p + j \mapsto (j-1)q + i; \quad i = 1, \dots, q, \quad j = 1, \dots, p.$$

Let us find the number of inversions in τ : this is the number of pairs (i_1, j_1) and (i_2, j_2) satisfying

$$(i_1 - 1)p + j_1 > (i_2 - 1)p + j_2, \quad (j_1 - 1)q + i_1 < (j_2 - 1)q + i_2$$

or, equivalently,

$$(i_1 - i_2)p > j_2 - j_1, \quad (j_2 - j_1)q > i_1 - i_2. \quad (1)$$

It follows from (1) that $(i_1 - i_2)pq > i_1 - i_2$ and hence

$$i_1 > i_2, \quad j_2 > j_1. \quad (2)$$

Since $j \leq p$ and $i \leq q$, inequalities (2) imply (1). The number of pairs satisfying (2) equals

$$\binom{p}{2} \binom{q}{2} \equiv \frac{pq}{4} \pmod{2},$$

as claimed. \square

Let us now set $p = 2n, q = 2$. As before, let x and y be the Euler classes of the tautological bundles, and denote by $u \in H_{2n}(G_{2n}^+(2n+2))$ and $v \in H_{4n-2}(G_{2n}^+(2n+2))$ the respective homological Euler classes. Set $w = v \cap \dots \cap v \in H_{2n}(G_{2n}^+(2n+2))$. It follows from Lemma 2.1 that

$$u \cap u = 2, w \cap w = 2(-1)^n, u \cap w = 0.$$

As a consequence, u and w are linearly independent.

It is known that $H_{2n}(G_{2n}^+(2n+2)) = \mathbf{Z}^2$, see, e.g., [9]. It follows that one may take u and w for a basis in $H_{2n}(G_{2n}^+(2n+2); \mathbf{Q})$.

Lemma 2.2 *One has: $F_*[M] = (\chi/2)u$.*

Proof. Note that the induced bundles $F^*(\xi)$ and $F^*(\nu)$ are the tangent and the normal bundles of M . One has:

$$F_*([M]) \cap u = F_*([M]) \cdot x = [M] \cdot F^*(x) = [M] \cdot F^*(e(\xi)) = [M] \cdot e(TM) = \chi.$$

It is well known that the Euler class of the normal bundle $\nu(M)$ of an embedded manifold M vanishes. Hence

$$F_*([M]) \cap w = F_*([M]) \cdot y^n = [M] \cdot F^*(y^n) = [M] \cdot e(\nu(M))^n = 0.$$

One can write: $F_*[M] = au + bw$ with $a, b \in \mathbf{Q}$, and Lemma 2.1 implies that $a = \chi/2$ and $b = 0$. \square

It follows that

$$\sigma_* F_*[M] \cap F_*[M] = -\left(\frac{\chi}{2}\right) u \cap \left(\frac{\chi}{2}\right) u = -\frac{\chi^2}{2}. \quad (3)$$

If $\chi \neq 0$ then $F(M)$ and $\sigma F(M)$ must intersect, therefore M has negatively parallel tangent spaces. This proves the first statement of Theorem 1.

Let us now prove Theorem 2.

Lemma 2.3 *One has: $e(\nu(M)) = -2d$.*

Proof. Since \mathbf{R}^4 is contractible, one has: $[M] \cap [M] = 0$. On the other hand, this homological self-intersection can be computed as follows. Choose a generic section γ of the normal bundle $\nu(M)$ and let M_ε be the result of pushing M slightly along this section. Then every double point of M contributes two points (with the same sign) to $M \cap M_\varepsilon$, and each zero of γ contributes one point to this intersection. It follows that $e(\nu(M)) + 2d = 0$. \square

Arguing as in Lemma 2.2, one finds that $F_*([M]) = (1 - g)u + dw$, and therefore

$$\sigma_* F_* [M] \cap F_* [M] = 2(d^2 - (1 - g)^2).$$

This implies Theorem 2.

Let us return to the general position statement of Theorem 1. If the Gauss image $F(M^{2n}) \subset G_{2n}^+(2n + 2)$ is an immersed submanifold which intersects $\sigma(F(M^{2n}))$ transversally then their homological intersection equals the algebraic number of the intersection points. In particular, there are no fewer intersection points than the absolute value of the homological intersection number, see (3).

Thus we need the following lemma in which “generic” is understood as belonging to an open and dense subset in the space of smooth maps with an appropriate topology; see, e.g., [8].

Lemma 2.4 *For a generic immersion $M^n \rightarrow \mathbf{R}^{n+2}$, the tangent Gauss map $G : M \rightarrow G_n(n + 2)$ is an immersion with transverse self-intersections.*

Proof. Locally, M is represented as the graph of a smooth map $U^n \rightarrow \mathbf{R}^2$ where U is a domain in \mathbf{R}^n . Thus one has two functions, say, u and v , of variables $x = (x_1, \dots, x_n)$. The tangent space to M at the point corresponding to $x \in U$ is parallel to the graph of the linear map $\mathbf{R}^n \rightarrow \mathbf{R}^2$ with the $n \times 2$ matrix $A(x) = (u_{x_i}, v_{x_i})$, $i = 1, \dots, n$. Hence, in an appropriate local chart of the Grassmanian, the Gauss map $G : U \rightarrow G_n(n + 2)$ is given by $x \mapsto A(x)$.

The Gauss map is an immersion if the $2n \times n$ matrix $B = (u_{x_i x_j}, v_{x_i x_j})$, $i, j = 1, \dots, n$ has full rank n . This matrix is formed by two symmetric $n \times n$ matrices of second partial derivatives. Consider the $n(n + 1)$ -dimensional space of $2n \times n$ matrices formed by two symmetric $n \times n$ matrices, and let Δ be its algebraic subvariety that consists of matrices of rank $n - 1$ or less.

Claim: $\text{codim } \Delta \geq n + 1$.

Assuming this claim, one proceeds as follows. In the space of 2-jets of smooth maps $U^n \rightarrow \mathbf{R}^2$, consider the algebraic subvariety Σ of the maps for which the matrix B is not of full rank. By the above claim, this subvariety has codimension $n + 1$. Then the Thom transversality theorem (see, e.g., [8]) implies that the 2-jet extension of a generic map $U^n \rightarrow \mathbf{R}^2$ avoids Σ , and Lemma 2.4 follows.

It remain to prove the claim. Consider a $2n \times n$ matrix $C = (S, T)$ where S and T are symmetric $n \times n$ matrices. By choosing an appropriate basis in \mathbf{R}^n , one can diagonalize both matrices. Assume that $\text{rank } C = n - 1$.

Assume that the first $k \leq n - 1$ diagonal entries of S are non-zero, and the remaining $n - k$ are equal to zero. Since the rank of C is $n - 1$, there are $n - k - 1$ non-zero entries among $t_{k+1,k+1}, \dots, t_{n,n}$; assume that $t_{i,i} \neq 0$ for $i = k + 2, \dots, n$.

Every matrix, close to C , can be written as $C_\varepsilon = (S + \varepsilon U, T + \varepsilon V)$ where U and V are symmetric. Then u_{ij} and v_{ij} are local coordinates in a neighborhood of C . For small ε , the rank of C_ε is not less than $n - 1$. Denote the columns of C_ε by η_1, \dots, η_{2n} .

If $\text{rank } C_\varepsilon = n - 1$ then, for every $j = k + 1, k + 2, \dots, n + k + 1$, the rank of the n -tuple of n -dimensional vectors $\eta_1, \dots, \eta_k, \eta_j, \eta_{n+k+2}, \dots, \eta_{2n}$ equals $n - 1$. Hence the $k + 1$ -st components of the vectors η_j vanish for $j = k + 1, \dots, n + k + 1$, that is, $u_{k+1,j} = 0$ for $j = k + 1, \dots, n$ and $v_{k+1,j} = 0$ for $j = 1, \dots, k + 1$. This gives $n + 1$ equations on the variables u_{ij} and v_{ij} , and these equations are independent of each other. This completes the proof. \square

Remarks 2.1 1. In the case $n = 1$, the Grassmanian is the product of spheres: $G_2^+(4) = S^2 \times S^2$. Denote the two factors by S_1 and S_2 . The main result of [2, 3] is that $F_*[M] = (\chi(M)/2)([S_1] + [S_2])$. The relation with classes u and v is as follows: $u = [S_1] + [S_2], v = [S_2] - [S_1]$. The map d that takes a plane to its orthogonal complement is the identity on S_1 and the antipodal involution on S_2 . Under the identification $\mathbf{R}^4 = \mathbf{C}^2$, the space of complex lines in \mathbf{C}^2 identifies with $\{N, S\} \times S_2$ where N and S are the poles of S_1 .

2. Suppose that an immersed surface M in the statement of Theorem 2 is totally real, that is, the tangent plane to M is never a complex line. Then $F_*[M] = a[S_2]$ for some $a \in \mathbf{Z}$, and the algebraic number of negatively parallel tangent planes is zero. Alternatively, the normal bundle of a totally real immersed surface is isomorphic to the tangent bundle, which implies that $d^2 = (1 - g)^2$. Likewise, this number is zero for an immersed surface M with non-vanishing normal curvature, see [12]. Indeed, it is proved in [12] that the normal Euler class of M is $\pm 2\chi(M)$, that is, $d = \pm\chi(M)$.

3. It follows from the above proof that χ is even. Indeed, a theorem of Seifert states: a closed orientable embedded manifold $M^{2n} \subset \mathbf{R}^{2n+2} = \mathbf{C}^{n+1}$ has an even Euler characteristic. It is proved in [9] that $\chi(M)$ is twice the algebraic number of positively oriented complex points of M .

4. The Grassmanian $G_{2n}^+(2n + 2) = G_2^+(2n + 2)$ is the space of oriented great circles in the unit sphere S^{2n+1} . The space of oriented geodesics of a Riemannian manifold has a canonical symplectic structure (assuming it is a smooth manifold), see, e.g., [1]. Thus $G_{2n}^+(2n + 2)$ is a symplectic manifold, and $\omega^{2n} \neq 0$ where ω is the symplectic form. In the previous notation, the cohomology class of ω is y .

3 Odd-dimensional skew sphere

Now we shall construct an odd-dimensional skew sphere of Theorem 3.

Let $M^{m-1} \subset \mathbf{R}^m$ be a smooth strictly convex closed hypersurface containing the origin. Denote by $h : S^{m-1} \rightarrow \mathbf{R}$ the support function of M , that is, $h(x)$ is the distance from the origin to the tangent hyperplane to M for which the unit vector $x \in S^{m-1}$ is the outward normal. Extend h to \mathbf{R}^m as a homogeneous function of degree 1. By the Euler formula,

$$xh_x = h, \quad xh_{xx} = 0. \quad (4)$$

Here and elsewhere we use the following vector notation: for m -dimensional vectors x, y and a function g of m variables, one has:

$$xy = \sum_{i=1}^m x_i y_i, \quad xdy = \sum_{i=1}^m x_i dy_i, \quad g_x = (g_{x_1}, \dots, g_{x_m}),$$

$$xg_{xx} = \left(\sum_{i=1}^m x_i g_{x_i x_1}, \dots, \sum_{i=1}^m x_i g_{x_i x_m} \right),$$

etc.

The hypersurface M is parameterized by the unit sphere as follows.

Lemma 3.1 *Let $y(x) \in M$ be the point at which the outward unit normal to M is $x \in S^{m-1}$. Then $y = h_x$.*

Proof. Consider the hypersurface M' given by the formula $y(x) = h_x$. To prove that the tangent hyperplane to M' at y is orthogonal to x , one needs to check that the 1-form xdy vanishes on M' . Indeed, $xy = h$ by (4), and hence

$$xdy = d(xy) - ydx = dh - h_x dx = 0.$$

The distance to the tangent hyperplane with the normal vector x is $xy = xh_x$, which equals h according to (4). Thus h is the support function of M' and therefore $M' = M$. \square

It follows from (4) that, at point $x \in S^{m-1}$, the linear operator h_{xx} annihilates the normal direction x and preserves the tangent space $T_x S^{m-1}$. The convexity of M implies that, on the tangent space, h_{xx} is non-degenerate (for example, h_{xx} is the identity if $h = 1$ and hence M is the unit sphere). Denote the restriction of h_{xx}^{-1} on $T_x S^{m-1}$ by $A(x)$ and extend A to the normal direction, spanned by x , as the zero map.

We will construct a skew brane in \mathbf{R}^{m+1} as a section of the vertical cylinder over $M \subset \mathbf{R}^m \subset \mathbf{R}^{m+1}$. More specifically, let $f : S^{m-1} \rightarrow \mathbf{R}$ be a smooth function. Define

$$N = \{(h_x(x), f(x)) \mid x \in S^{m-1}\} \subset \mathbf{R}^m \times \mathbf{R} = \mathbf{R}^{m+1},$$

and let $\phi : S^{m-1} \rightarrow N$ be the parameterization map.

Let us find out when the tangent spaces to N are parallel. Assume that f is extended to \mathbf{R}^m as a homogeneous function of, say, degree 1 (the degree of homogeneity is of no importance).

Lemma 3.2 *The tangent spaces to N are parallel at distinct points $\phi(x_1)$ and $\phi(x_2)$ if and only if $x_2 = -x_1$ and $A(f_x)(x_2) = A(f_x)(x_1)$.*

Proof. Let us describe the normal 2-plane to N at point $\phi(x)$. This plane is generated by the vector $(x, 0)$ and a vector $(\xi, 1)$ where $\xi \in T_x S^{m-1}$. We claim that $\xi = -A(f_x)$.

Indeed, let $v \in T_x S^{m-1}$ be a test vector, $v = \sum_j v_j \partial_{x_j}$. Then

$$d\phi(v) = \sum_j (v_j h_{x_i x_j}, v_j f_{x_j}), \quad i = 1, \dots, m.$$

It follows that $d\phi(v)$ is orthogonal to $(\xi, 1)$ if and only if $v h_{xx} \xi + v f_x = 0$, or $v(h_{xx} \xi + f_x) = 0$. Since v is an arbitrary tangent vector to S^{m-1} , the projection of the vector $h_{xx} \xi + f_x$ to $T_x S^{m-1}$ is zero. This implies the claim.

Finally, the span of vectors $(\xi(x_2), 1)$ and $(x_2, 0)$ coincides with that of $(\xi(x_1), 1)$ and $(x_1, 0)$ if and only if $x_2 = -x_1$ and $\xi(x_2) = \xi(x_1) + t x_1$. Since $\xi(x_2)$ and $\xi(x_1)$ are orthogonal to x_1 , one has $\xi(x_2) = \xi(x_1)$. \square

Thus N is a skew brane if and only if

$$A(-x)(f_x(-x)) \neq A(x)(f_x(x)) \quad (5)$$

for all $x \in S^{m-1}$. Decompose the functions h and f into the even and odd components with respect to the antipodal involution of the sphere $x \mapsto -x$:

$$h = h_{ev} + h_{odd}, \quad f = f_{ev} + f_{odd}.$$

Then one has a similar decompositions of the operator A and the gradient vector field f_x . Note that

$$(f_x)_{ev} = (f_{odd})_x, \quad (f_x)_{odd} = (f_{ev})_x.$$

Decomposing (5) into even and odd parts yields:

$$A_{ev}((f_{ev})_x) + A_{odd}((f_{odd})_x) \neq 0 \quad (6)$$

for all $x \in S^{m-1}$.

If M is sufficiently close to the sphere then h is close to 1 and A is close to the identity. Thus we may assume that A_{ev} is invertible and rewrite (6) as

$$(f_{ev})_x + B((f_{odd})_x) \neq 0 \quad (7)$$

where $B(x) = A_{ev}^{-1} A_{odd}$ is an odd field of linear maps of the tangent spaces $T_x S^{m-1}$.

Note that (7) cannot hold for all x if h , and therefore A , is even: then $B = 0$ and the function f_{ev} must have critical points on the sphere. Note also that (7) cannot hold if $m - 1$ is even since there exist no non-vanishing vector fields on even-dimensional spheres. From now on, assume that $m = 2n$.

Lemma 3.3 *Let M^m be a connected manifold and B a smooth field of linear automorphisms of the cotangent spaces to M satisfying the following property: for every smooth function $f : M \rightarrow \mathbf{R}$ there exists a smooth function $g : M \rightarrow \mathbf{R}$ such that $B(df) = dg$. Then $B = c \text{Id}$ for some constant c .*

Proof. Being a local statement, we may assume that M is a contractible domain in \mathbf{R}^m . In local coordinates x_1, \dots, x_m , the linear map B is given by a matrix $b_{ij}(x)$. The condition is that, for every f , the 1-form $\sum_{ij} b_{ij} f_{x_j} dx_i$ is closed.

Choose $j \in \{1, \dots, m\}$ and let $f(x) = x_j$. Then the 1-form $\sum_i b_{ij}(x) dx_i$ is closed, and hence there exists a function $g^j(x)$ such that $b_{ij} = g^j_{x_i}$. The 1-form $\sum_{ij} g^j_{x_i} f_{x_j} dx_i = \sum_j f_{x_j} dg^j$ is closed for every f , therefore so is $\sum_j g^j d(f_{x_j})$.

Again fix $j \in \{1, \dots, m\}$ and let $f(x) = x_j^2/2$. Then the 1-form $g^j dx_j$ is closed, hence g^j depends only on x_j . Finally, fix $j, k \in \{1, \dots, m\}$, $j \neq k$, and let $f(x) = x_j x_k$. Then the 1-form $g^j(x_j) dx_k + g^k(x_k) dx_j$ is closed, and therefore $g^j_{x_j} = g^k_{x_k} = c$. It follows that $b_{ij} = c \delta_{ij}$, as claimed. \square

One has the following corollary.

Corollary 3.1 *If M is not centrally symmetric then there exists a function f_{odd} on S^{m-1} such that $B((f_{\text{odd}})_x)$ is not a gradient vector field.*

Proof. A parallel translation of the origin changes the support function h by addition of a linear function and does not affect h_{xx} . The hypersurface M is centrally symmetric if and only if, after a parallel translation, h is even, that is, A is even. Since M is not centrally symmetric, $A_{\text{odd}} \neq 0$ and $B \neq 0$.

Identify the tangent and cotangent spaces of the sphere by the Euclidean structure. Let U be a small domain on S^{m-1} . Then every function $f : U \rightarrow \mathbf{R}$ can be extended to S^{m-1} as an odd function. If $B(f_x)$ is a gradient for every such f then, by the preceding lemma, $B = c \text{Id}$ in U . This equality then must hold everywhere on the sphere. But B is odd, hence $B = 0$, a contradiction. \square

Our plan of proof of Theorem 3 is to construct functions h and f on S^{2n-1} so that (7) holds everywhere on S^{2n-1} . We set $h = 1 + \varepsilon g$ where g is an odd function and ε is a small parameter. Then $h_{xx} = \text{Id} + \varepsilon g_{xx} + O(\varepsilon^2)$ and $A = \text{Id} - \varepsilon g_{xx} + O(\varepsilon^2)$. It follows that $B = -\varepsilon g_{xx} + O(\varepsilon^2)$, and (7) can be rewritten as

$$(f_{ev})_x \neq \varepsilon g_{xx}((f_{\text{odd}})_x) + O(\varepsilon^2). \quad (8)$$

We wish to construct a function f and an odd function g so that, for all sufficiently small ε and a certain constant $c > 0$, depending on f and g , one has

$$|(f_{ev})_x - \varepsilon g_{xx}((f_{\text{odd}})_x)| > c\varepsilon \quad (9)$$

everywhere on the sphere. This will imply (8), and hence, for sufficiently small ε , the desired inequality (7) as well.

Let (z_1, \dots, z_n) be coordinates in \mathbf{C}^n . Choose distinct non-zero reals a_1, \dots, a_n , and let $f_{ev} = \sum_i a_i |z_i|^2$. Denote by ξ the unit Hopf vector field on S^{2n-1} . Then f_{ev} is an even, ξ -invariant Morse-Bott function on S^{2n-1} with exactly n critical Hopf circles C_1, \dots, C_n .

Lemma 3.4 *There exist odd smooth functions f_{odd} and g on S^{2n-1} such that $g_{xx}((f_{\text{odd}})_x) \cdot \xi = 2$ on each critical circle $C_i, i = 1, \dots, n$.*

Assuming this lemma, denote the vector field $g_{xx}((f_{\text{odd}})_x)$ by v . Let c be a constant such that $|v(x)| < c$ everywhere on the sphere (this constant exists by compactness of the sphere). Given $\varepsilon > 0$, let U_ε be a ξ -invariant neighborhood of the critical set $C_1 \cup \dots \cup C_n$ such that $|(f_{ev})_x| > 2\varepsilon c$ outside of U_ε . We claim that (9) holds outside of U_ε . Indeed, $|(f_{ev})_x - \varepsilon v| > 2\varepsilon c - \varepsilon c = \varepsilon c$, as claimed.

Next, consider the situation inside U_ε . Since $v \cdot \xi = 2$ on the critical circles, we can choose ε so small that $v \cdot \xi > 1$ inside U_ε . Note that $(f_{ev})_x$ is orthogonal to ξ . Therefore $|((f_{ev})_x - \varepsilon v) \cdot \xi| = \varepsilon v \cdot \xi > \varepsilon$, and hence $|((f_{ev})_x - \varepsilon v)| > \varepsilon$ inside U_ε . In particular, (9) holds for this choice of ε .

Thus, Theorem 3 will follow, once we prove Lemma 3.4.

Proof of Lemma 3.4. We construct the desired functions in a neighborhood of each critical circle, symmetric with respect to the antipodal involution, and then extend them to the sphere.

Let C be one such Hopf circle. Consider $\mathbf{C}^2 \subset \mathbf{C}^n$ such that C is a Hopf circle therein. Choose coordinates (z_1, z_2) in \mathbf{C}^2 so that $z_i = x_i + \sqrt{-1}y_i$, $i = 1, 2$, and C is given by $x_2 = y_2 = 0$. The Hopf field is given by the formula:

$$\xi = -y_1 \partial x_1 + x_1 \partial y_1 - y_2 \partial x_2 + x_2 \partial y_2.$$

The functions $f = f_{\text{odd}}$ and g will depend only on (x_1, y_1, x_2, y_2) .

It is straightforward to compute the operator g_{xx} , the gradient f_x and the dot product $g_{xx}(f_x) \cdot \xi$. Introduce the differential operator depending on f :

$$D = f_{x_1} \partial x_1 + f_{y_1} \partial y_1 + f_{x_2} \partial x_2 + f_{y_2} \partial y_2.$$

Then,

$$g_{xx}(f_x) \cdot \xi = x_1 D(g_{y_1}) - y_1 D(g_{x_1}) + x_2 D(g_{y_2}) - y_2 D(g_{x_2}). \quad (10)$$

Let us look for f and g in the following form:

$$f = a(x_1, y_1) + x_2 b(x_1, y_1) + y_2 d(x_1, y_1), \quad g = u(x_1, y_1) + x_2 v(x_1, y_1) + y_2 w(x_1, y_1)$$

where a, u are odd and homogeneous of degree 1, b, d, v, w are even and homogeneous of degree 0. Switch to polar coordinates $x_1 = r \cos \alpha, y_1 = r \sin \alpha$. On the circle C , one has $x_2 = y_2 = 0, r = 1$, and α is a coordinate. Then (10) becomes

$$bv' + dw' + (u + u'')a' \quad (11)$$

where prime is $d/d\alpha$. Set:

$$a = u = 0, \quad b = -w = \cos 2\alpha, \quad d = v = \sin 2\alpha,$$

and the expression (11) gets identically equal to 2. In terms of the Cartesian coordinates,

$$f = x_2(x_1^2 - y_1^2) - 2y_2 x_1 y_1, \quad g = 2x_2 x_1 y_1 - y_2(x_1^2 - y_1^2)$$

on the unit sphere.

It remains to extend the odd functions from a neighborhood of the union of the critical Hopf circles to the whole sphere as an odd function. Clearly, one can extend a smooth function φ from an open subset U to a function Φ on the whole sphere (for example, using a smooth cut-off function for the set U). If φ is odd then one may modify $\Phi(x)$ by replacing it with $(1/2)(\Phi(x) - \Phi(-x))$. The latter is an odd function on the sphere that coincides with φ on U . This completes the proof. \square

4 Two conjectures

Generalizing [6], we make the following conjecture.

Conjecture 4.1 *For every non-centrally symmetric $M^{2n-1} \subset \mathbf{R}^{2n}$ there exists a skew brane $N^{2n-1} \subset \mathbf{R}^{2n+1} = \mathbf{R}^{2n} \times \mathbf{R}$ which is a section of the vertical cylinder over M .*

Conjecture 4.1 follows from a more general conjecture of independent interest.

Conjecture 4.2 *Let M be a closed manifold with zero Euler characteristic, α a non-closed differential 1-form on it. Then there exists a smooth function $f : M \rightarrow \mathbf{R}$ such that the 1-form $df + \alpha$ has no zeroes on M .*

Deducing Conjecture 4.1 from Conjecture 4.2. One may rephrase (7) as

$$d(f_{ev}) + \alpha \neq 0 \tag{12}$$

everywhere on S^{2n-1} ; here α is the 1-form dual to the vector field $B((f_{odd})_x)$. By Corollary 3.1, we may choose $f_{odd} : S^{2n-1} \rightarrow \mathbf{R}$ so that α is not closed. Consider the quotient space $M = \mathbf{RP}^{2n-1} = S^{2n-1}/\mathbf{Z}_2$. Then α descends as a non-closed 1-form $\bar{\alpha}$ on M . Conjecture 4.2 implies that there exists a function $\bar{f} : \mathbf{RP}^{2n-1} \rightarrow \mathbf{R}$ such that $d\bar{f} + \bar{\alpha}$ is nowhere vanishing. Then \bar{f} lifts to an even function on S^{2n-1} for which (12) holds. \square

Remark 4.1 In dimension one, Conjecture 4.2 holds. Indeed, $\alpha = g(x)dx$ with $c := \int_0^1 g(x)dx \neq 0$; here x is the coordinate on the circle \mathbf{R}/\mathbf{Z} . Then $g - c$ is a derivative, $g(x) - c = -f'(x)$, and $df + \alpha = cdx$. It follows that, in dimension one, Conjecture 4.1 holds as well: this is precisely the Cylinder Lemma of [6].

We conclude with comments on Conjecture 4.2. If the 1-form is exact, i.e., $\alpha = dg$ for some function g , then zeros of $df + \alpha$ are the critical points of the function $f + g$. Morse theory implies that the number of these critical points is bounded below in terms of the topology of the manifold M , e.g., by the sum of its Betti numbers. Likewise, if α is a closed 1-form, one has similar lower

bounds for the number of zeros of a 1-form in the cohomology class of α , this time, due to the Morse-Novikov theory. Conjecture 4.2 states that if neither Morse nor Morse-Novikov theory applies then there is no topological reason for the existence of zeros of the form $df + \alpha$.

Conjecture 4.2 also resembles the following theorem of Polterovich [13] and Laudenbach-Sikorav [11]. It is known in symplectic topology that, under certain assumptions, the number of intersection points of a Lagrangian submanifold in a symplectic manifold with its translate under a Hamiltonian isotropy is at least the sum of its Betti numbers. This result of symplectic topology generalizes Morse inequalities. The theorem of Polterovich and Laudenbach-Sikorav asserts that if L is a middle-dimensional submanifold of a symplectic manifold which is not Lagrangian and which has zero Euler characteristic then L can be displaced from itself by a Hamiltonian vector field. The proof of this theorem makes use of Gromov's h -principle (the technique of convex integration). I believe that a similar approach will be instrumental in a proof of Conjecture 4.2.

References

- [1] V. Arnold, A. Givental. Symplectic geometry, 1–136. *Encycl. of Math. Sci., Dynamical Systems*, 4, Springer-Verlag, 1990.
- [2] W. Blaschke. Sulla geometria differenziale delle superficie S_2 nello spazio euclideo S_4 . *Ann. Mat. Pura Appl.* (4) 28 (1949), 205–209.
- [3] S.-S. Chern, E. Spanier. A theorem on orientable surfaces in four-dimensional space. *Comment. Math. Helv.* 25 (1951), 205–209.
- [4] M. Ghomi. Nonexistence of skew loops on ellipsoids. *Proc. Amer. Math. Soc.* 133 (2005), 3687–3690.
- [5] M. Ghomi. Tangent bundle embeddings of manifolds in Euclidean space. *Comment. Math. Helv.* 81 (2006), 259–270.
- [6] M. Ghomi, B. Solomon. Skew loops and quadratic surfaces. *Comment. Math. Helv.* 77 (2002), 767–782.
- [7] M. Ghomi, S. Tabachnikov. Totally skew embeddings of manifolds. *Math. Z.* 258 (2008), 499–512.
- [8] M. Golubitsky, V. Guillemin. *Stable mappings and their singularities*. Springer-Verlag, 1973.
- [9] H. F. Lai. Characteristic classes of real manifolds immersed in complex manifolds. *Trans. Amer. Math. Soc.* 172 (1972), 1–33.
- [10] H. F. Lai. On parallel tangents of embeddings in codimension 2 in Euclidean spaces. *Indiana Univ. Math. J.* 22 (1973), 1171–1181.

- [11] F. Laudenbach, J.-C. Sikorav. Hamiltonian disjunction and limits of Lagrangian submanifolds. *Internat. Math. Res. Notices*, 1994, no. 4, 161–168.
- [12] J. Little. On singularities of surfaces in E^4 . *Bull. Amer. Math. Soc.* 75 (1969), 368–369.
- [13] L. Polterovich. An obstacle to non-Lagrangian intersections. The Floer memorial volume, 575–586, Birkhauser, 1995.
- [14] B. Segre. Sulle coppie di tangenti fra loro parallele relative ad una curva chiusa sghemba. *Hommage au Professeur Lucien Godeaux*, pp. 141–167, Louvain 1968.
- [15] B. Segre. Global differential properties of closed twisted curves. *Rend. Sem. Mat. Fis., Milano*, 38 (1968), 256–263.
- [16] J.-P. Sha, B. Solomon. No skew branes on non-degenerate hyperquadrics. *Math. Z.* 257 (2007), 225–229.
- [17] B. Solomon. Skew loops in flat tori. *Geom. Dedicata* 128 (2007), 33–37.
- [18] B. Solomon. Central cross-sections make surfaces of revolution quadric. *Amer. Math. Monthly* 116 (2009), 351–355.
- [19] B. Solomon. Surfaces with central cross-sections. Preprint arXiv:0904.3493
- [20] G. Stojanovic. Embeddings with multiple regularity. *Geom. Dedicata* 123 (2006), 1–10.
- [21] G. Stojanovic, S. Tabachnikov. Non-existence of n -dimensional T -embedded discs in \mathbf{R}^{2n} . *Comment. Math. Helv.* 81 (2006), 877–882.
- [22] S. Tabachnikov. On skew loops, skew branes and quadratic hypersurfaces. *Moscow Math. J.* 3 (2003), 681–690.
- [23] J. White. Global properties of immersions into Euclidean spheres. *Indiana Univ. Math. J.* 20 (1971), 1187–1194.
- [24] Y.-Q. Wu. Knots and links without parallel tangents. *Bull. London Math. Soc.* 34 (2002), 681–690.